

# Plant Extracts, Isolated Phytochemicals, and Plant-Derived Agents Which Are Lethal to Arthropod Vectors of Human Tropical Diseases – A Review

## Authors

Adrian Martin Pohlit<sup>1,2</sup>, Alex Ribeiro Rezende<sup>2</sup>, Edson Luiz Lopes Baldin<sup>3</sup>, Norberto Peporine Lopes<sup>2</sup>, Valter Ferreira de Andrade Neto<sup>4</sup>

## Affiliations

<sup>1</sup> Instituto Nacional de Pesquisa da Amazônia, Manaus, Amazonas State, Brazil

<sup>2</sup> Universidade de São Paulo, Ribeirão Preto, São Paulo State, Brazil

<sup>3</sup> Universidade Estadual de São Paulo, Botucatu, São Paulo State, Brazil

<sup>4</sup> Universidade Federal de Rio Grande do Norte, Natal, Rio Grande do Norte State, Brazil

## Key words

- botanicals
- acaricide
- insecticidal and larvicidal plants
- plant extracts
- essential oils
- biotechnology
- natural products
- phytochemicals

received Sept. 23, 2010  
revised February 19, 2011  
accepted March 4, 2011

## Bibliography

DOI <http://dx.doi.org/10.1055/s-0030-1270949>  
Published online March 22, 2011  
Planta Med 2011; 77: 618–630  
© Georg Thieme Verlag KG  
Stuttgart · New York ·  
ISSN 0032-0943

## Correspondence

**Prof. Valter Ferreira de Andrade Neto, PhD**  
Departamento de Microbiologia e Parasitologia  
Laboratório de Biologia da Malária e Toxoplasmose  
Universidade Federal do Rio Grande do Norte – Campus Universitário  
Av. Senador Salgado Filho – Lagoa Nova  
CEP 69061-000 – Natal – RN  
Brazil  
Phone: + 55 84 32 15 34 37 ext. 226  
Fax: + 55 84 32 11 92 10  
aneto@cb.ufrn.br

## Abstract



The recent scientific literature on plant-derived agents with potential or effective use in the control of the arthropod vectors of human tropical diseases is reviewed. Arthropod-borne tropical diseases include: amebiasis, Chagas disease (American trypanosomiasis), cholera, cryptosporidiosis, dengue (hemorrhagic fever), epidemic typhus (Brill-Zinsser disease), filariasis (elephantiasis), giardia (giardiasis), human African trypanosomiasis (sleeping sickness), isosporiasis, leishmaniasis, Lyme disease (lyme borreliosis), malaria, onchocerciasis, plague, recurrent fever, sarcocystosis, scabies (mites as causal agents), spotted fever, toxoplasmosis, West Nile fever, and yellow fever. Thus, coverage was given to work describing plant-derived extracts, essential oils (EOs), and isolated chemicals with toxic or noxious effects on filth bugs (mechanical vectors), such as common houseflies (*Musca domestica* Linnaeus), American and German cockroaches (*Periplaneta americana* Linnaeus, *Blattella germanica* Linnaeus), and oriental latrine/blowflies (*Chrysomya megacephala* Fabricius) as well as biting,

blood-sucking arthropods such as blackflies (*Simulium* Latreille spp.), fleas (*Xenopsylla cheopis* Rothschild), kissing bugs (*Rhodnius* Stål spp., *Triatoma infestans* Klug), body and head lice (*Pediculus humanus humanus* Linnaeus, *P. humanus capitis* De Geer), mosquitoes (*Aedes* Meigen, *Anopheles* Meigen, *Culex* L., and *Ochlerotatus* Lynch Arribalzaga spp.), sandflies (*Lutzomyia longipalpis* Lutz & Neiva, *Phlebotomus* Loew spp.), scabies mites (*Sarcoptes scabiei* De Geer, *S. scabiei* var *hominis*, *S. scabiei* var *canis*, *S. scabiei* var *suis*), and ticks (*Ixodes* Latreille, *Amblyomma* Koch, *Dermacentor* Koch, and *Rhipicephalus* Koch spp.). Examples of plant extracts, EOs, and isolated chemicals exhibiting noxious or toxic activity comparable or superior to the synthetic control agents of choice (pyrethroids, organophosphorous compounds, etc.) are provided in the text for many arthropod-vectors of tropical diseases.

**Supporting information** available online at <http://www.thieme-connect.de/ejournals/toc/plantamedica>

## Introduction



### Arthropod vectors of human tropical disease

Arthropods are the vectors of a variety of human diseases which affect tropical countries around the world (● **Table 1**) [1–22]. Biting, blood-sucking arthropods are the most important vectors in terms of public health. These include mosquitoes, tsetse flies [7], kissing bugs, sandflies, ticks, etc., which are responsible for the transmission of malaria, dengue hemorrhagic fever, filariasis [7], American [3–6] and Human African [7] trypanosomiasis, leishmaniasis [7, 13], spotted [7, 18], West Nile, and yellow fevers [7], among other se-

vere tropical diseases. Importantly, several dozens of species of anopheline mosquitoes are responsible for the transmission of the 4 *Plasmodium* parasite species which cause human malaria. According to data from the World Health Organization (WHO), half the world's human population lives in regions where malaria is endemic [7]. On the other hand, non-biting or non-blood-sucking coprophagic (feces-eating), saprophagic (living on decaying or decomposing materials), and other arthropods, such as dung beetles, common houseflies, and cockroaches can be mechanical vectors of amebiasis [1, 2], cholera [7, 8], cryptosporidiosis [1, 7], giardia [1, 10, 11], isosporiasis [12], sarcocystosis [1], toxoplasmosis [1, 16], and

**Table 1** Human tropical diseases, etiological agents, and arthropod vectors.

Tropical disease			Arthropod vector		
Name	Etiologic agent	Occurrence	Common name	Scientific name	Source
Amebiasis	<i>Entamoeba histolytica</i> , <i>E. dispar</i>	China, Mexico, S. America, S.-E. & W. Africa, S.-E. Asia	German cockroach	<i>Blatella germanica</i> Linnaeus	[1, 2]
			American cockroach	<i>Periplaneta americana</i> Linnaeus	
			Common housefly	<i>Musca domestica</i> Linnaeus	
Chagas disease (American trypanosomiasis)	<i>Trypanosoma cruzi</i>	Americas, Europe	Kissing bug	<i>Triatoma</i> , <i>Rhodnius</i> Stål spp., <i>Panstrongylus megistus</i> Burmeister	[3–6]
Cholera	<i>Vibrio cholerae</i>	Worldwide	Common housefly	<i>Musca domestica</i> Linnaeus	[7, 8]
Cryptosporidiosis	<i>Cryptosporidium</i> sp.	Africa, Asia, Australia, Europe, Americas	Dung beetle	<i>Onthophagus</i> Latreille, <i>Anoplotrupes</i> Jekel & <i>Aphodius</i> Illiger spp.	[1, 7]
			Common housefly	<i>Musca domestica</i> Linnaeus	
Dengue (hemorrhagic fever)	<i>Flavivirus</i> sp.	Africa, Americas, E. Mediterranean, S.-E. Asia & W. Pacific	Mosquito	<i>Ae. Aegypti</i> L., <i>Ae. Albopictus</i> Skuse	[7]
Epidemic typhus (Brill-Zinsser disease)	<i>Rickettsia prowazekii</i>	Africa, C. & S. America	Louse	<i>Pediculus humanus corporis</i>	[9]
Filariasis or Elephantiasis	<i>Wuchereria bancrofti</i>	Africa, Asia, S. America	Mosquito	<i>Aedes</i> Meigen, <i>Anopheles</i> Meigen, <i>Culex</i> L. spp.	[7]
Giardia, giardiasis	<i>Giardia lamblia</i>	Worldwide	German cockroach	<i>Blatella germanica</i> Linnaeus	[1, 10, 11]
			Common housefly	<i>Musca domestica</i> Linnaeus	
Human African trypanosomiasis (sleeping sickness)	<i>Trypanosoma brucei</i>	Africa	Tsetse fly	<i>Glossina</i> Wiedemann spp.	[7]
Isosporiasis	<i>Isospora belli</i>	Africa, Australia, Caribbean Islands, Latin America, S.-E. Asia	Dung beetle	<i>Onthophagus</i> Latreille spp.	[12]
			Common housefly	<i>Musca domestica</i> Linnaeus	
Leishmaniasis	<i>Leishmania</i> sp.	Asia, C. & S. America, E. Africa, Europe, India	Sandfly	<i>Lutzomyia</i> França, <i>Phlebotomus</i> Loew & <i>Sergentomyia</i> França & Parrot spp.	[7, 13]
Lyme disease (Lyme borreliosis)	<i>Borrelia</i> spp.	Asia, Europe, Americas	Tick	<i>Ixodes</i> Latreille, <i>Amblyomma</i> Koch spp.	[14, 15]
Malaria	<i>Plasmodium</i> spp.	Africa, Mexico, C. & S. America, Asia	Mosquito	<i>Anopheles</i> Meigen spp.	[7]
Onchocerciasis	<i>Onchocerca volvulus</i>	Africa, Mexico, C. & S. America	Blackfly	<i>Simulium</i> Latreille spp.	[7]
Plague	<i>Yersinia pestis</i>	Africa, Asia, Brazil, Bolivia, Peru, Ecuador, USA	Oriental rat flea	<i>Xenopsylla cheopis</i> Rothschild	[10]
Recurrent fever	<i>Borrelia recurrentis</i>	Africa, Asia, Europe, N. America	Louse	<i>Pediculus humanus</i> spp.	[14]
			Tick	<i>Ornithodoros</i> C. L. Koch spp.	
Sarcocystosis	<i>Sarcocystis</i> spp.	Worldwide	German cockroach	<i>Blatella germanica</i> Linnaeus	[1]
			American cockroach	<i>Periplaneta Americana</i> Linnaeus	
			Common housefly	<i>Musca domestica</i> Linnaeus	
Scabies	–	Worldwide	(Itch) Mite	<i>Sarcoptes scabiei</i> var <i>hominis</i>	[17]
Spotted fever	<i>Rickettsia rickettsii</i>	Brazil, Colombia, Mexico, Panama, Canada, USA	Tick	<i>Dermacentor</i> Koch, <i>Rhipicephalus</i> Koch, <i>Amblyomma</i> Koch spp.	[7, 18]
Toxoplasmosis	<i>Toxoplasma gondii</i>	Worldwide	Dung beetle	<i>Onthophagus</i> Latreille spp.	[1, 16]
			German cockroach	<i>Blatella germanica</i> Linnaeus	
			American cockroach	<i>Periplaneta americana</i> Linnaeus	
			Common housefly	<i>Musca domestica</i> Linnaeus	
			Oriental latrine or blowfly	<i>Chrysomya megacephala</i> Fabricius	
West Nile fever	<i>Flavivirus</i> sp.	Worldwide	Mosquito	<i>Culex</i> L. spp., <i>Ochlerotatus</i> Lynch Arribálzaga spp.	[7]
Yellow fever	<i>Flavivirus</i> sp.	Africa, Latin America	Mosquito	<i>Ae. aegypti</i> L., <i>Ae. albopictus</i> Skuse	[7]

other infectious diseases and so must also be controlled for reasons of disease prevention and public health (Table 1).

### Literature on Plants for Arthropod Vector Control

In recent years, numerous scientific reports have been published on plants which are useful (or potentially useful) for the control of arthropod vectors of tropical diseases. Major emphasis has been on the most important arthropod-vector – the mosquito. Reviews

of the scientific literature have been published recently on mosquito larvicidal plant extracts and fractions [19], plant EOs exhibiting arthropod-killing (mosquitocidal, larvicidal) among other biological activities [20], and mosquito repellent and insecticidal plant EOs and chemical components [21–23]. The patent literature on plant EO-containing mosquito repellent inventions is also reviewed in this special issue of *Planta Medica* [24]. In the present review, emphasis is on literature published in the period 2007–2010 describing plant extracts and their active chemical components which cause the death of or are noxious to one or more of

the developmental stages (eggs, nymphs/larvae, pupae, adults) of a broad range of arthropod vectors of human tropical diseases such as blowflies, common houseflies, cockroaches, fleas, lice, mosquitoes, ticks, etc.

The plant sources of extracts, EOs, fractions, and isolated compounds which exhibit toxicity to, are noxious to, or are otherwise useful in the control of arthropods covered herein are terrestrial plants which generally have medicinal and other useful (economic) properties. Edible green and blue-green algae (cyanobacterium) having toxic properties towards insect species including larvae of *Aedes* Meigen, *Anopheles* Meigen, and *Culex* L. spp. were recently reviewed [25]. Thus, algae and aquatic plants are not covered herein nor are arthropod/insect control derivatives from bacteria and fungi (such as *Bacillus thuringiensis israelensis*, *B. sphaericus*, and *Saccharopolyspora spinosa*/Spinosa<sup>®</sup>) cultures.

### Blackflies

The water extracts of the leaves of *Chromolaena odorata* (L.) R. M. King & H. Rob. exhibited lethality to larvae of *Simulium damnosum* Theobald (LC<sub>50</sub> = 1 µg/mL) which was not statistically different from that of the synthetic organophosphorous larvicide chlorpyrifos [26].

Hydrogenated catnip (*Nepeta cataria* L. leaf, stem) EO containing 15 weight-percent (wt%) of stereoisomeric dihydronepetalactones (1.67 g/m<sup>2</sup>) as active repellent ingredients was formulated into a liquid and a lotion both of which provided > 7.5 h mean complete protection against adult *Simulium decorum* Walker in the field [27].

Remarkably, in a field study in Thailand, the blackfly *Simulium nigrogilvum* Summers was effectively repelled by lotions containing 10% w/w EO in absolute ethanol (60%) and additives vanillin (10%), propylene glycol (10%), and polyethylene glycol (10%). Thus finger root (*Boesenbergia rotunda* [L.] Mansf.) EO, guava (*Psidium guajava* L.) leaf EO, and turmeric (*Curcuma longa* L.) EO were separately formulated into lotions which were tested together with the proprietary product Repel Care<sup>®</sup> (active ingredients: 5% w/w turmeric EO and 4.5% w/w *Eucalyptus citriodora* Hook. EO) and DEET (10% w/w lotion formulated as for EOs). All five formulations provided 100% protection for 9 h and > 82% for 10 and 11 h against *S. nigrogilvum* [28].

### Blowflies

Topical application of eucalyptol (1,8-cineole) caused the death of *Chrysomya megacephala* Fabricius adult males and females (LD<sub>50</sub> = 197 and 221 µg/fly, respectively) [29]. Eucalyptol exhibited low activity against *C. megacephala* third instars (LD<sub>50</sub> = 642 µg/µL) using the dipping method. Also, *Azadirachta indica* A. Juss. seed extracts (containing 0.24% azadirachtin A) caused swelling of the protocuticle of third instars and first stage pupae of *C. megacephala* by the dipping method [30]. Larvae of *C. megacephala* were effectively killed by betel (*Piper betle* L.) EO. At concentrations of 3–4% betel EO, 100% larvae mortality was observed [31]. Nine plant EOs were screened for ovicidal activity against *C. megacephala* and *Eugenia caryophyllata* Thunb., *Illicium verum* Hook. f. and *Cinnamomum cassia* (L.) C. Presl EOs exhibited the most activity (LC<sub>50</sub> = 1.61, 2.49, and 0.43 mg/mL, respectively). Also, synthetic cinnamaldehyde showed ovicidal activity (LC<sub>50</sub> = 0.28 mg/mL) [32].

### Cockroaches

Pure components from EOs were screened for lethality against different developmental stages of *Blattella germanica* Linnaeus [34]. Topically applied pure components caused the death of adult males, females, gravid females, and nymphs at different stages of development of *B. germanica*. The most active substances against *B. germanica* adult males were thymol, *E*-cinnamaldehyde, carvacrol, and eugenol (LD<sub>50</sub> = 0.070, 0.078, 0.101, 0.109 mg/cockroach, respectively). Against *B. germanica* adult females the most active substances were carvacrol, *E*-cinnamaldehyde, and thymol (LD<sub>50</sub> = 0.186, 0.188, and 0.195 mg/cockroach, respectively), and these same substances were also the most active against gravid *B. germanica*: thymol > *E*-cinnamaldehyde > carvacrol (LD<sub>50</sub> = 0.122, 0.133, and 0.146 mg/cockroach, respectively).

Large *B. germanica* nymphs were also susceptible to topical treatment with EO components, and for large nymphs the most active substances were *E*-cinnamaldehyde, carvacrol, and thymol (LD<sub>50</sub> = 0.117, 0.129, and 0.220 mg/cockroach, respectively) while for medium nymphs the most active substances were thymol, carvacrol, *E*-cinnamaldehyde, and eugenol (LD<sub>50</sub> = 0.060, 0.061, 0.082, and 0.109 mg/cockroach, respectively) [34]. Small *B. germanica* nymphs were the most sensitive to natural components of EOs, and substances exhibited activity in the following order: *E*-cinnamaldehyde > thymol ~ geraniol > carvacrol ~ *S*-(-)-limonene > (-)-menthone (LD<sub>50</sub> = 0.036–0.060 mg/cockroach). Diminished nymph hatching was observed for *B. germanica* eggs treated with (-)-menthone. Based on these and other data, the group of substances thymol, *E*-cinnamaldehyde, carvacrol, geraniol, and eugenol exhibited interesting lethality to different stages of development of *B. germanica* [34].

Using a T-tube olfactometer, 17 EOs were screened, and 5 EOs exhibited significant repellency against *B. germanica* and other cockroach species. Thus, adult female *Periplaneta americana* Linnaeus (American cockroach) and *B. germanica* (German cockroach) were repelled by grapefruit (*Citrus × paradisi* Macfad.), lemon (*Citrus × limonum* Risso), lime [*Citrus × aurantiifolia* (Christm.) Swingle], orange [*Citrus × sinensis* (L.) Osbeck] EOs by 90.3, 85.7, 83.3, and 70.0%, respectively, and 96.7, 92.9, 86.7, and 71.4%, respectively [33]. Adult female *B. germanica* were also repelled by clove leaf (*Eugenia caryophyllata* Thunb.) EO (repellency = 70.0%). Another cockroach, *Periplaneta fuliginosa* Serville, was less repelled by these *Citrus* L. spp. EOs: 82.4, 72.0, 70.6, and 62.5%, respectively. Yoon et al. [33] identified two different “types” of *Citrus* oils based on the relative quantity of the components limonene (> 90% = type I = grapefruit and orange EOs; 48–61% = type II = lemon and lime EOs) and  $\alpha$ -pinene,  $\beta$ -pinene,  $\gamma$ -terpinene,  $\beta$ -myrcene, and benzene. Generally, type I EOs exhibited repellencies which could be reproduced by placing the volume of pure limonene present in 10 µL of EO in the olfactometer for both *B. germanica* and *P. americana*. To study the effects of type II EOs, limonene (6.4 µL) combined with  $\beta$ -pinene (1.4 µL) was found to provide repellency (90.3%) comparable to that of 10 µL of lemon and lime (type II) EOs against *B. germanica*. Adult female *B. germanica* were repelled about equally by limonene (ca. 6.1 µL) +  $\beta$ -pinene (1.6 µL) or 10 µL of type II (lemon and lime) EOs. In a ternary mixture of limonene +  $\beta$ -myrcene +  $\gamma$ -terpinene (6.1 + 0.1 + 0.4 µL), a repellency of 83.3% was attained which was comparable to the type II EO repellency against adult female *B. germanica*. When the individual components were tested for repellency against *P. americana* adult females in the volumes present in 10 µL of type II EOs,  $\beta$ -pinene (1.17 µL) fully repro-

duced the repellency of 10  $\mu\text{L}$  of lime EO. For type II EOs, to reproduce the repellent effect of 10  $\mu\text{L}$  of lime and lemon EOs to adult *P. americana*, only ternary combinations of limonene (6.1  $\mu\text{L}$ ) +  $\gamma$ -terpinene (0.4  $\mu\text{L}$ ) + [either  $\beta$ -myrcene (0.1  $\mu\text{L}$ ),  $\beta$ -pinene (1.4  $\mu\text{L}$ ), or  $\alpha$ -pinene (0.2  $\mu\text{L}$ )] produced repellencies (80–83%) comparable to those of both type II EOs [33]. This work demonstrates the very complex synergistic and suppressive interactions among the monoterpenes in these *Citrus* EOs and the species specific nature of repellency in these cockroach species.

### Common houseflies

Solvent extracts of plants and isolated chemical components have been tested for important control effects mainly against mature *Musca domestica* Linnaeus. Thus, seed and root petroleum ether extracts of *Griffonia simplicifolia* (M. Vahl ex DC.) Baill. and root and stem petroleum ether extracts of *Zanthoxylum xanthoxyloides* (Lam.) Waterman repelled (median repellent doses,  $\text{RD}_{50}$  = 1.0–1.7  $\mu\text{g}/\text{cm}^2$ ) and killed mature *M. domestica* (topical  $\text{LC}_{50}$  = 0.3–0.5  $\mu\text{g}/\text{fly}$ ) [35]. In other work, EtOH extracts of *A. indica* which contain the active ingredient azadirachtin A were found to be highly lethal to adult flies (94% mortality) at a concentration of 0.025%, but were only moderately lethal to earlier stages (larvae, pupae) even at higher concentrations [30]. Coumarin was isolated from the hexane extract of the leaves of *Ageratum conyzoides* L. and shown to be highly toxic ( $\text{LD}_{50}$  = 1.2,  $\text{LD}_{90}$  = 3.7 mg/g) to mature *M. domestica* [36]. Also, friedelin was isolated from *Cacalia tangutica* (Maxim.) Hand.-Mazz. extract and (as a coating on sugar) was found to be as lethal ( $\text{LC}_{50}$  = 0.130 mg/g sugar) to mature *M. domestica* as rotenone ( $\text{LC}_{50}$  = 0.091 mg/g sugar) which was used as control substance [37]. These examples demonstrate the potential of plant extracts and their active principles as sources of control agents for mature *M. domestica*.

A number of plant EOs and volatile chemical components exhibiting adulticidal activity against *M. domestica* have been identified in recent publications. *Eucalyptus* L'Hér. spp., *Citrus*  $\times$  *sinensis* (L.) Osbeck, *Lavandula angustifolia* Mill., *Mentha* L. spp., and *Pelargonium graveolens* L'Hér. EOs were found to effectively knock down and kill *M. domestica* ( $\text{KT}_{50}$  = 3–18 min;  $\text{LD}_{50}$  = 0.07–0.16  $\mu\text{g}/\text{insect}$ ) [38]. Several of the isolated monoterpenes of these EOs, namely 1,8-cineole ( $\text{KT}_{50}$  = 2.3 min,  $\text{LD}_{50}$  = 0.13  $\mu\text{g}/\text{insect}$ ), limonene ( $\text{KT}_{50}$  = 7.5 min,  $\text{LD}_{50}$  = 0.10  $\mu\text{g}/\text{insect}$ ), linalool ( $\text{KT}_{50}$  = 7.6 min,  $\text{LD}_{50}$  = 0.04  $\mu\text{g}/\text{insect}$ ), menthone ( $\text{KT}_{50}$  = 19.0 min,  $\text{LD}_{50}$  = 0.11  $\mu\text{g}/\text{insect}$ ), and menthyl acetate ( $\text{KT}_{50}$  = 22.6 min,  $\text{LD}_{50}$  = 0.09  $\mu\text{g}/\text{insect}$ ) were found to be toxic to *M. domestica* [38]. However, an independent study on 1,8-cineole found much lower lethal doses against *M. domestica* mature males ( $\text{LD}_{50}$  = 118  $\mu\text{g}/\text{fly}$ ) and females ( $\text{LD}_{50}$  = 177  $\mu\text{g}/\text{fly}$ ) [29]. In another screening for fumigant activity, *C. sinensis*, *Citrus*  $\times$  *aurantium* L., *Citrus*  $\times$  *limonum* Risso, *Citrus*  $\times$  *paradisi* Macfad., *Citrus*  $\times$  *reticulata* Blanco, *Coriandrum sativum* L., *Eucalyptus cinerea* F. Muell. ex Benth., *Laurus nobilis* L., and *Myristica fragrans* Houtt. EOs were found to be active ( $\text{LC}_{50}$  = 3.9–8.8 mg/cm<sup>3</sup>) against mature *M. domestica*, and several of the monoterpene components of these EOs were evaluated for activity against *M. domestica*. Besides 1,8-cineole ( $\text{LC}_{50}$  = 3.3 mg/dm<sup>3</sup>), (–) and (+)-limonene ( $\text{LC}_{50}$  = 5.0 and 6.2 mg/dm<sup>3</sup>, respectively), chemical components such as  $\gamma$ -terpinene ( $\text{LC}_{50}$  = 4.0 mg/dm<sup>3</sup>),  $\alpha$ -terpinene ( $\text{LC}_{50}$  = 6.2 mg/dm<sup>3</sup>), citronellal ( $\text{LC}_{50}$  = 8.1 mg/dm<sup>3</sup>), (–)- $\beta$ -pinene ( $\text{LC}_{50}$  = 6.4 mg/dm<sup>3</sup>), and (–)- $\alpha$ -pinene ( $\text{LC}_{50}$  = 8.9 mg/dm<sup>3</sup>) were also found to be toxic to mature *M. domestica* [39]. Phenylpropanoids are another class of volatile compounds having important lethality to *M. domestica*. In a

recent study on the lethality of the EO and chemical components from the leaves of *Piper betle* L. to *M. domestica*, EO ( $\text{LC}_{50}$  = 10.3 mg/dm<sup>3</sup>) and individual phenylpropanoid EO components safrole ( $\text{LC}_{50}$  = 4.8 mg/dm<sup>3</sup>), dihydrosafrole ( $\text{LC}_{50}$  = 4.7 mg/dm<sup>3</sup>), isosafrole ( $\text{LC}_{50}$  = 2.3 mg/dm<sup>3</sup>), and eugenol ( $\text{LC}_{50}$  = 7.3 mg/dm<sup>3</sup>) all proved to have significant toxicity to mature *M. domestica* [40].

In the Supporting Information, data from literature sources on plant extracts, EOs, monoterpenes, and phenylpropanoids exhibiting fumigant and other activities against *M. domestica* are presented.

### Dung beetles

No recent studies on plant extracts, oils, or chemical components for the control of *Anoplotrupes* Jekel & *Aphodius* Illiger spp. of dung beetles were found. This could be an interesting group for future study given that these filth-associated organisms exist in many countries around the world where basic hygiene and coexistence with these species may be a source of contamination and infections (● Table 1).

### Fleas

Incense cedar (*Calocedrus decurrens* [Torr.] Florin) heartwood EO ( $\text{LC}_{50}$  = 0.24,  $\text{LC}_{90}$  = 0.31 mg/mL), Port-Orford [*Chamaecyparis lawsoniana* (A. Murray) Parl.] cedarwood EO ( $\text{LC}_{50}$  = 1.21,  $\text{LC}_{90}$  = 1.85 mg/mL), and western juniper (*Juniperus occidentalis* Hook.) heartwood EO ( $\text{LC}_{50}$  = 0.31,  $\text{LC}_{90}$  = 0.93 mg/mL) exhibited adulticidal effects on *Xenopsylla cheopis* Rothschild fleas [41].

Isolated components from Alaska yellow cedar (*Chamaecyparis nootkatensis* [D. Don] Spach) heartwood EO, derivatives, and commercially acquired products were tested against *X. cheopis*. Carvacrol, valencene, nootkatene, crystalline nootkatone, nootkatone grapefruit extract, isolated nootkatone, valencene-13-ol, nootkatol, valencene-13-aldehyde, nootkatone 1,10 epoxide, nootkatone diepoxide all were active against *X. cheopis* exhibiting  $\text{LC}_{50}$  = 0.0029–0.064% w/v and  $\text{LC}_{90}$  = 0.0049–0.10% w/v 24 h after exposure. Nootkatone (grapefruit EO) was the most active sample tested ( $\text{LC}_{50}$  = 0.0029,  $\text{LC}_{90}$  = 0.008% w/v) [42].

### Kissing bugs (assassin bugs, bloodsucking conenoses, “barbeiros”)

24 plant extracts were screened for insecticidal activity against fourth stage blood-replete nymphs of *Rhodnius milesi* Carcallo, Rocha, Galvão & Jurberg by applying 50  $\mu\text{g}$  of each extract to the abdomen of the nymphs. Hexane and ethanol extracts of *Simarouba versicolor* A.St.-Hil., *Guarea kunthiana* A. Juss., *G. guidonia* (L.) Sleumer, and *Talauma ovata* A.St.-Hil. caused 20–95% mortality among nymphs, and the ethanol extract of the root bark of *S. versicolor* and hexane extract of the roots of *G. guidonia* were responsible for 95 and 75% nymph mortalities, respectively [43].

Topically applied *Pilocarpus spicatus* A. St.-Hil. leaf EO was toxic to and paralyzed *Rhodnius prolixus* Stål fifth stage male nymphs (0.5 and 1.0  $\mu\text{L}$  EO/insect, 90.5 and 91.1% mortality, respectively, after 24 h; 89 and 92% paralysis of surviving nymphs after 15 days) as well as retarded moulting and had partial antifeedant effects [44]. In other work, EOs and monoterpenes were screened for fumigant activity (exposure to vapors emitted by 100  $\mu\text{L}$  of EO or monoterpene in a closed vessel) against *R. prolixus* first instars. Eucalyptus EO was the most active fumigant ( $\text{KT}_{50}$  = 216 min) and eucalyptol (1,8-cineole) was the most active fumigant monoterpene ( $\text{KT}_{50}$  = 117 min) [45].



In the above work, EOs and monoterpenes were also screened for repellency against *R. prolixus* first instars. Thus, mint and lavender EOs produced slight repellent effects at 400 µg/cm<sup>2</sup>; geraniol and menthyl acetate produced slight repellent effects, respectively, at 40 and 400 µg/cm<sup>2</sup>; and menthone produced a slight repellent effect at 400 µg/cm<sup>2</sup> [45].

*Schinus molle* L. leaf and root hexane extracts exhibited greater repellency than DEET against first instars of blood-sucking cone-noses *Triatoma infestans* Klug. Also, 3% w/v (maximum concentration tested) hexane extract of the fruit of *S. molle* caused 80% inhibition of hatching of *T. infestans* eggs [46].

## Lice

*Hedychium spicatum* Buch.-Ham. ex Sm. rhizome EO at 1–5% concentrations exhibited pediculicidal activity against human body lice (*Pediculus humanus humanus*) which was greater than a 1% permethrin based product which was tested for comparison [47]. In other work, head lice (*P. humanus capitis*) were killed by the ethyl acetate extracts of the seeds of custard apple (*Annona squamosa* L.), and the hexane extract of seeds contains oleic acid (13.88 wt%) and a triglyceride with one oleate ester (7.70 wt%). Ethyl acetate extract, oleic acid, and triglyceride with one oleate ester were diluted (1 : 1) in inactive coconut oil and found to kill clinically obtained *P. humanus capitis* in 31.7, 47.3, and 10.0 min, respectively [48]. In another study involving clinically-obtained third instars and adult head lice (*P. humanus capitis*), 1,8-cineole was found to inhibit acetylcholine-esterase in a homogenate of *P. humanus capitis* and was found also to cause intoxication (knockdown) after 20 min of exposure to 1,8-cineole vapor. This result was better than contact with the standard lice control compound DDVP (dichlorvos) which after 60 min of exposure had only knocked down 50% of head lice [49]. This same group of researchers tested 23 monoterpenoid compounds for ovicidal and fumigant activity (adulticide activity) in permethrin-resistant *P. humanus capitis*. Of 6 monoterpenes screened, only (+)- $\alpha$ -pinene (KT<sub>50</sub> = 34.5 min) and (-)- $\alpha$ -pinene (KT<sub>50</sub> = 28.5 min) had fumigant activity against adult *P. humanus capitis* [50]. These authors compared this result to previous work by the group in which 1,8-cineole (KT<sub>50</sub> = 11.1 min), anisole (KT<sub>50</sub> = 12.7 min), limonene (KT<sub>50</sub> = 27.2 min),  $\beta$ -pinene (KT<sub>50</sub> = 33.9 min), linalool (KT<sub>50</sub> = 37.7 min), menthone (KT<sub>50</sub> = 39.7 min),  $\alpha$ -pinene (KT<sub>50</sub> = 42.7 min), and benzyl alcohol (KT<sub>50</sub> = 59.7 min) were shown to be active fumigants against adult *P. humanus capitis*.

Louse egg mortalities > 80% relative to negative controls were obtained for anisole (100%),  $\alpha$ -pinene (97%),  $\beta$ -pinene (96%), (+)- $\alpha$ -pinene (96%), 1S(-)- $\alpha$ -pinene (94%), anethole (93%), carvone (92%), limonene (90%), linalool (88%) and 1,8-cineole (84%) [50]. These results demonstrate the potential use of monoterpenes and other substances in the control of permethrin-resistant head lice.

## Mosquitoes

Several recent reviews have been published on mosquito control agents from plants and related topics. In 2010, Nerio et al. [23] published a review on mosquito-repellent EOs and their repellent components. Both Bakkali et al.'s 2008 review [20] on the biological activity and toxicity of EOs and components and Burfield & Reekie's (2005) review [21] on EOs and components for mosquito control included data from publications on larvicidal, adulticidal, and other biological activities related to mosquito control. In the present work, the literature on plant extracts, fractions, and isolated chemical components having useful biological activ-

ities for mosquito control were reviewed. Given the large number of publications, emphasis was given to literature describing the lethal activity of isolated phytochemicals against mosquitoes. As Supporting Information, adulticidal, larvicidal, and ovicidal activities and other effects of plant extracts, essential oils, and fractions are summarized for publications which failed to report biological activity for isolated component chemicals (active principles) and which were not covered in this review. Also, Supporting Information includes data on extracts, EOs, and fractions as well as most nonvolatile active components discussed in this review. Publications on active extracts and essential oils having at least one (isolated) active principle which is potentially relevant for the purpose of mosquito control are discussed below.

## Mosquitocides

EOs and their isolated components which exhibit important fumigant activity in knockdown and adulticide assays against mosquito species have been the subject of recent publications. Kiran and Devi [88] described fumigant activity of *Chloroxylon swietenia* DC. EO to adult *An. gambiae*, *Cx. quinquefasciatus*, and *Ae. aegypti* (LD<sub>50</sub> = 1.0, 1.2, and 1.7 µg/cm<sup>3</sup>, respectively; KT<sub>50</sub> = 0.32, 0.50, and 0.72 h, respectively). This fumigant activity was attributed to sesquiterpene components present in the EO such as germacrene D (LD<sub>50</sub> = 1.8, 2.1, and 2.8 µg/cm<sup>3</sup>, respectively) which was purportedly acting synergistically with other components in the EO (EO was more active than these individually tested components) (Table 4). In other work, *Eucalyptus* spp. EOs exhibited general efficiency in knocking down adult *Ae. aegypti* (KT<sub>50</sub> = 4–12 min), which was associated with the presence of 1,8-cineole and other components in the EOs (Table 4) [89]. These authors also confirmed the relationship between the vapor pressure of *Eucalyptus* spp. EOs and highly active individual components such as 1,8-cineole,  $\alpha$ -pinene, and *p*-cymene and median knockdown time (KT<sub>50</sub>) in *Ae. aegypti* adult mosquitoes. *Mentha × piperita* L. was also found to be highly active against mature *Ae. aegypti* [90]. *Chloroxylon swietenia*, *Eucalyptus* spp., and *Mentha × piperita* EOs contain a number of volatile mosquitocidal chemical components whose fumigant activity was investigated in the above studies and is summarized in Table 4. Knockdown times of just 4–6 min were observed for 1,8-cineole, *p*-cymene, and  $\alpha$ -pinene (*Eucalyptus* spp.) against mature *Ae. aegypti* while mugetanol,  $\alpha$ -terpineol, and thymol (found in *Mentha* L. spp. EO) were highly effective at knocking down and killing mature *Ae. aegypti*, *An. tessellates* Theobald, and *Cx. quinquefasciatus*. Interestingly, L-menthol was found not to be toxic to adult *Ae. aegypti* [90]. This is a reminder that molecular and taxonomic specificities may be important characteristics of the toxic effects of certain mosquitocidal substances.

## Mosquito larvicides

*A. indica* or neem oil formulations are important mosquito control agents, and a formula containing 0.15% of the limonoid azadirachtin was tested and found to be highly effective at killing *Ae. aegypti* (LC<sub>50</sub> = 1.7 ppm) and *An. stephensi* Liston (LC<sub>50</sub> = 1.6 ppm) in the lab and *Aedes* spp. (95–100% reduction of larvae over 7 days), *Anopheles* spp. (80–100% reduction of larvae over 3 weeks), and *Culex* spp. (≥ 80% reduction of larvae over 3 weeks) in the field [51]. For another formulation comprised of wood and bark chips of neem tree (*A. indica*) in water, effective larvicidal activity was observed against first thru fourth instars of *An. gambiae* (IE<sub>90</sub> = 0.12–0.6 g wood chips/L H<sub>2</sub>O) [52]. Interestingly, no azadirachtin was found in these aqueous solutions; however,

the limonoids nimbin and salannin were detected by HPLC analysis by comparison with authentic samples. As these and earlier studies have demonstrated, *A. indica* (neem) derivatives, especially neem oil and the active ingredient azadirachtin, show great promise as general plant-based mosquitocides and larvicides and are the basis for a number of commercially available products.

The limonoid calodendrolide was isolated from the root bark of *Calodendrum capense* Thunb. and was reduced to pyroangolenolide. These two limonoids exhibited important larvicidal activity against *Ae. aegypti* ( $LC_{50}$  = 13.2 and 16.6  $\mu$ M, respectively). In this same work, the authors also isolated the limonoids harrisonin and pedonin from the methanol extract of the root bark of *Harrisonia abyssinica* Oliv. and demonstrated that these two compounds had activity against *Ae. aegypti* larvae ( $LC_{50}$  = 28.1 and 59.2  $\mu$ M, respectively) [53]. Two quassinoids, neosergeolide and isobrucein B, were isolated from the roots and stems of the Amazonian medicinal plant *Picrolemma sprucei* Hook. f. and exhibited good larvicidal activity against *Ae. aegypti* ( $LC_{50}$  = 8.7 and 6.7  $\mu$ M, respectively) [54].

The sesquiterpenes cubebol ( $LC_{50}$  = 68.6 and 50.0  $\mu$ g/mL against *Ae. aegypti* and *Ae. albopictus* Skuse, respectively) and *epi*-cubebol ( $LC_{50}$  = 100 and 63.8  $\mu$ g/mL against *Ae. aegypti* and *Ae. albopictus*, respectively) together with ferruginol ( $LC_{50}$  = 64.1  $\mu$ g/mL against *Ae. aegypti*) were isolated from the wood extract of *Cryptomeria japonica* (Thunb. ex L.f.) D. Don [55]. Sesquiterpenoid metabolites 9-oxoneoprocumeneol ( $LC_{50}$  = 5.81 ppm) and neoprocumeneol ( $LC_{50}$  = 13.7 ppm) exhibited larvicidal activity against *Cx. quinquefasciatus* Say and were isolated from the petroleum ether extract ( $LC_{50}$  = 11.4 ppm) of wild turmeric (*Curcuma aromatic* Salisb.) root [56].

A group from Brazil described the larvicidal activity ( $LC_{50}$  = 8.9 ppm) of the oil-resin from the trunk of *Copaifera reticulata* Ducke against *Ae. aegypti* and also the successful fractionation of the oil-resin to obtain active sesquiterpene ( $LC_{50}$  = 0.2 ppm) and labdane-enriched ( $LC_{50}$  = 0.8 ppm) fractions [57]. Later, this group reported the isolation of mosquito larvicidal diterpenes 3- $\beta$ -acetoxylabdan-8(17)-13-dien-15-oic acid ( $LC_{50}$  = 0.8 ppm) and alepterolic acid ( $LC_{50}$  = 87.3 ppm) [58].

Rahuman et al. [59] isolated the tetracyclic triterpene derivative gluanol acetate from the acetone extract of the bark of *Ficus racemosa* L. and found that it had important larvicidal activity against *Ae. aegypti*, *An. stephensi*, and *Cx. quinquefasciatus* fourth instars ( $LD_{50}$  = 14.6, 28.5, and 41.4 ppm, respectively).

The dichloromethane extract of the root bark of the traditionally used mosquito repellent plant *Lantana viburnoides* subsp. *viburnoides* var. *kisi* had potent larvicidal activity against *An. gambiae* Giles s.s. (72 h  $LC_{50}$  = 7.70 ppm). Active fractions of this extract contained the lantadene-type triterpene camaric acid (72 h  $LC_{50}$  = 6.19 ppm) and the lupine triterpene betulinic acid (72 h  $LC_{50}$  < 10 ppm) [60].

The steroid  $\beta$ -sitosterol was isolated from the petroleum ether extracts of the leaves of *Abutilon indicum* (Linn.) Sweet [61] and a saponin of unknown identity was isolated from *Achyranthes aspera* L. [62], and both exhibited significant toxicity to *Ae. aegypti* larvae ( $LC_{50}$  = 11.5 ppm and 18.2 ppm, respectively) and *Cx. quinquefasciatus* larvae ( $LC_{50}$  = 26.7 ppm and 27.2 ppm, respectively).  $\beta$ -Sitosterol was further found to be a larvicide against *An. stephensi* ( $LC_{50}$  = 3.58 ppm) [61]. A phytosteroid of unknown identity was isolated from the juice of crushed fresh leaves, and 50 ppm of this substance killed 83, 93, and 100% of *Cx. quinquefasciatus* larvae after 24, 48, and 72 h, respectively [63].

Anthraquinone and naphthoquinone larvicides have been described in recent literature. Thus, using bioguided fractionation in the search for larvicidal compounds against *Ae. aegypti* and *Ae. albopictus*, Cheng et al. isolated the anthraquinone tectoquinone ( $LC_{50}$  = 3.3 and 5.4  $\mu$ g/mL, respectively) from the highly active hexane fraction ( $LC_{50}$  = 2.4 and 3.3  $\mu$ g/mL, respectively) of the methanol extract of the sapwood of *Cyptomeria japonica* (Forssk.) Vahl [64]. From the ethyl acetate extracts of the leaves of *Cassia nigricans* Vahl, the anthraquinones emodin ( $LC_{50}$  = 2.4  $\mu$ g/mL after 48 h), citreoresin ( $LC_{50}$  = 7.67  $\mu$ g/mL after 48 h), and emodic acid ( $LC_{50}$  = 2.88  $\mu$ g/mL after 48 h) were isolated and found to be highly active larvicides against *An. gambiae* [65].

From the chloroform extracts of the rhizomes of *Plumbago capensis* Thunb. naphthoquinones and naphthoquinone dimers isoshinanolone ( $LC_{50}$  = 1.26  $\mu$ g/mL), plumbagin ( $LC_{50}$  = 5.43  $\mu$ g/mL), 3-O-methylhydroserone ( $LC_{50}$  = 31.5  $\mu$ g/mL), 6-hydroxyplumbagin ( $LC_{50}$  = 13.6  $\mu$ g/mL), maritinone ( $LC_{50}$  = 40.7  $\mu$ g/mL), and chitranane ( $LC_{50}$  = 31.2  $\mu$ g/mL) were isolated and found to be larvicidal to *Ae. aegypti* [66]. In other work, *Plumbago dawei* Rolfe ethyl acetate, *P. stenophylla* Wilmot-Dear chloroform, and *P. zeylanica* L. hexane and chloroform extracts of root barks exhibited strong larvicidal activity against *An. gambiae* ( $LC_{50}$  = 4.1–6.7  $\mu$ g/mL), and plumbin was isolated and confirmed as a potent *An. gambiae* larvicide ( $LC_{50}$  = 1.9  $\mu$ g/mL) [67]. Also, from the dichloromethane extract of the root bark of *Lantana viburnoides* subsp. *viburnoides* var. *kisi*, which was shown above to contain mosquito larvicidal triterpenes, one of the active fractions contained furanoquinone regioisomers (72 h  $LC_{50}$  = 5.48–5.70 ppm) [60].

Plants from the Brazilian cerrado biome were screened for larvicidal activity against *Ae. aegypti*, and the most active species found was *Ocotea velloziana* (Meisn.) Mez. The ethanol extract of the trunk bark ( $LC_{50}$  = 214  $\mu$ g/mL) was fractionated using biomonitoring to yield the larvicidal aromatic alkaloid (+)-dicentrine ( $LC_{50}$  = 30.2  $\mu$ g/mL) [68].

Fatty acids can have important larvicidal effects against different mosquito species. Thus, oleic and linoleic acids ( $LC_{50}/LC_{90}$  = 8.8/35.4 and 18.2/96.3 ppm, respectively) were isolated from the petroleum ether extracts of *Citrullus colocynthis* (L.) Schrad. and were shown to be the active larvicidal principles against *Cx. quinquefasciatus* [69]. These fatty acids are readily available from (saponified) fats and glyceridic oils, and they deserve more attention and study to establish their potential as mosquito larvicides. Plant proteins also can be important mosquito larvicides. A water soluble lectin was purified from the water extracts of the seeds of *Moringa oleifera* Lam. and shown to retard development and kill *Ae. aegypti* ( $LC_{50}$  = 0.197 mg/mL) larvae [70]. In other work, from the 0.15 M NaCl extracts of *Myracrodruon urundeuva* Allemão bark and heartwood, lectins were isolated and found to have high larvicidal activity against *Ae. aegypti* ( $LC_{50}$  = 0.125 and 0.040 mg/mL, respectively) [71]. Lastly, from the water extract of the fresh leaves of *Solanum villosum* Mill., a protein was isolated and purified and shown to have larvicidal activity against *Ae. aegypti*, *An. stephensi*, and *Cx. quinquefasciatus* ( $LC_{50}$  = 747, 645, and 646 ppm, respectively) [72].

Mosquito larvicidal compounds representing other chemical classes have been identified recently. For example, ethanol extracts of the roots of *Tephrosia toxicaria* (Sw.) Pers. exhibited larvicidal activity against *Ae. aegypti* ( $LC_{50}$  = 47.9 ppm) as did the hexane, chloroform, and ethyl acetate fractions ( $LC_{50}$  = 13.8–95.5 ppm).  $\alpha$ -Toxicarol was isolated and shown to have larvicidal activity against *Ae. aegypti* ( $LC_{50}$  = 24.6 ppm) [73]. In other work, galgotannin was isolated from the ethyl acetate extracts of *Quer-*

*cus infectoria* G. Olivier galls and was found to have larvicidal activity ( $LC_{50} = 125$  ppm) against *An. stephensi* [74]. Also, *Piper peltatum* L. root ethanol extract exhibited only slight lethality to *Ae. aegypti* third instars after 24 h [75]; however, based on isolated yields, this dry root can contain at least 5.7% w/w 4-nerolidylcatechol which is lethal ( $LC_{50} = 26.0$   $\mu\text{g/mL}$ ) to *Ae. aegypti* larvae [76].

In recent work, chemical components of plant EOs which are toxic to mosquito larvae have been described. For example, from the leaf and twig EO of *Piper aduncum* L., the volatile phenylpropanoid compound dillapiol was isolated and shown to be toxic to *Ae. aegypti* larvae ( $LC_{50} = 36.2$   $\mu\text{g/mL}$ ). Furthermore, isodillapiol ( $LC_{50} = 21.7$   $\mu\text{g/mL}$ ) and *n*-propyloxy and *n*-butyloxy propan-2-yl ether derivatives ( $LC_{50} = 28.0$  and  $19.7$   $\mu\text{g/mL}$ , respectively) prepared from dillapiol are more active larvicides than dillapiol [76]. Through compositional studies on the larvicidal EO of the roots of *Asarum heterotropoides* F. Schmidt (against *Ae. aegypti*, *Cx. pipiens pallens* Coquillett, and *Ochlerotatus togoi* Theobald,  $LC_{50} = 23.8$ ,  $21.1$ , and  $27.6$  ppm, respectively), the major components were identified and found to be terpenoid and phenylpropanoid compounds. These components were individually tested for activity against mosquito larvae. Thus, 3-carene, (+)-limonene,  $\alpha$ -phellandrene, safrole,  $\gamma$ -terpinene, and terpinolene were found to be highly toxic to *Ae. aegypti*, *Cx. pipiens pallens*, and *Oc. togoi* larvae (Table 2) [77]. Independently, R-limonene ( $LC_{50} = 37$  ppm) was shown to be the most important larvicidal active principle of the leaf EOs of *Lippia gracilis* Schauer against *Ae. aegypti* [78], but several other active substances were also found (Table 2). In studies on the composition and larvicidal activity of *Clausena excavata* Burm. f. leaf and twig EOs ( $LC_{50} = 37.1$  and  $40.1$   $\mu\text{g/mL}$ , respectively, against *Ae. aegypti*;  $LC_{50} = 41.2$  and  $41.1$   $\mu\text{g/mL}$ , respectively, against *Ae. albopictus*) [79], *Cryptomeria japonica* leaf EO ( $LC_{50} = 28.4$  and  $51.2$   $\mu\text{g/mL}$  against *Ae. aegypti* and *Ae. albopictus*, respectively) [80], and *Eucalyptus camaldulensis* Dehnh. and *E. urophylla* S.T. Blake leaf EOs ( $LC_{50} = 31.0$  and  $95.5$   $\mu\text{g/mL}$ , respectively, against *Ae. aegypti*) [81], 3-carene (against *Ae. aegypti*) [79], (+)-limonene (against *Ae. aegypti*) [79, 81], *p*-cymene (against *Ae. aegypti* and *Ae. albopictus*) [80, 81],  $\alpha$ -phellandrene (against *Ae. aegypti*) [81], myrcene (against *Ae. aegypti* and *Ae. albopictus*) [79, 80], and  $\alpha$ -terpinene (against *Ae. aegypti*) [81] were also found to be highly active larvicides ( $LC_{50} < 20$  ppm) (Table 2).

The leaf EOs of 6 chemotypes of *Cinnamomum osmophloeum* Kaneh. were studied, and two chemotypes with expressive larvicidal activity against *Ae. albopictus* ( $LC_{50} = 40.8$ – $46.5$   $\mu\text{g/mL}$ ) and one with expressive activity against the larvae of *Cx. quinquefasciatus* ( $LC_{50} = 31.3$   $\mu\text{g/mL}$ ) were described. Also, components of *C. osmophloeum* EOs such as *E*-cinnamaldehyde, benzaldehyde, and cinnamyl acetate were individually tested and found to exhibit significant larvicidal activity [82]. Diallyl disulfide (from *Allium sativum* L.) [83] and linalyl acetate (from *Citrus × aurantium*) [84] also exhibit important larvicidal effects (Table 2).

Thyme (*Thymus vulgaris* L.), parsley (*Petroselinum crispum* [Mill.] Fuss), anise (*Pimpinella anisum* L.), and coriander (*Coriandrum sativum* L.) EOs exhibited larvicidal activities ( $LC_{50} = 15.0$ ,  $34.3$ ,  $65.1$ , and  $156$   $\mu\text{g/mL}$ , respectively) against the seaside mosquito species *Ochlerotatus caspius* Pallas. Eugenol, thymol, carvacrol, *trans*-anethole, and linalool, which are components of these EOs, exhibited larvicidal activity (Table 2) [85].

Interestingly, the larvicidal activity of each of the enantiomers of  $\alpha$  and  $\beta$ -pinene was studied. In general (–)- $\beta$ -pinene was found to be the most toxic ( $LC_{50} = 13$ – $37$  ppm) to the larvae of *Ae. aegypti*,

*Cx. pipiens pallens* and *Oc. togoi* [77], and *Cx. pipiens* biotype *molestus* [86]. Enantiomeric composition, especially in components of EOs, is an important and often neglected aspect of the study of insecticide interactions. Substances such as linalool, limonene, etc., occur in EOs in one or both enantiomeric forms and the enantiomeric composition of the commercially supplied substance (often used in biological testing) is not necessarily the same as that of the enantiomeric composition of this component in the EO of a given plant under study. The identification of these larvicidal active components in *in vitro* studies is significant and requires further work to evaluate the potential of these components in formulations, their acceptability to humans, and their toxicity to humans and to mosquitoes in field settings.

From the data presented in Table 2 on recent studies on the larvicidal activity of components of EOs against mosquito species, the monoterpenes  $\beta$ -asarone, *p*-cymene, (+)-limonene, linalyl acetate, myrcene,  $\alpha$ -phellandrene, (+)- $\beta$ -pinene, (–)- $\beta$ -pinene,  $\alpha$ -terpinene,  $\gamma$ -terpinene and terpinolene (and perhaps thymol), phenylpropenes safrole and eugenol, and the sulfur-containing compound diallyl disulfide have potent larvicidal action on one or more species of mosquitoes. Further studies are needed to show the robustness or limitations of these substances as single component mosquito larvicides. Also, further studies on combinations of these components are needed to explore possible synergism, especially in those cases where the EOs are more active than the individual, isolated components.

In what is apparently the first general demonstration that piperonyl butoxide (PBO) has synergistic effects on larvicidal monoterpenes and other classes of substances found in plant essential oils, Waliwitiya et al. [87] showed that synergism ratios of up to 250 (2.5 orders of magnitude) could be obtained for substances such as borneol, linalool, camphor, and 1,8-cineole, as summarized in Table 3. Importantly, 10 mg/L of piperonyl butoxide (PBO) was established as the minimum sublethal concentration in 1st through 4th instars of *Ae. aegypti*, and this (low) concentration was used in that study to guarantee that the observed enlarged effectiveness by PBO was synergistic in nature [87].

PBO is a synthetic synergist which is used in commercial applications together with pyrethroids and other contact insecticides. It is an insect monooxygenase inhibitor which reduces or eliminates the insect's capacity to detoxify larvicidal and adulticidal compounds [87].

### Sandflies

*Lutzomyia longipalpis* Lutz & Neiva adults (sandfly vector of American visceral leishmaniasis) were killed by water extracts of the leaves of *Antonia ovata* Pohl ( $LD_{50} = 233$  mg/mL) and water extracts of the roots of *Derris amazonica* Killip ( $LD_{50} = 212$  mg/mL) [92]. Also, *Eucalyptus* spp. EOs exhibit toxic effects in contact with *L. longipalpis* adults. Thus, adulticidal effects were observed for lemon ironbark (*E. staigeriana* F. Muell. ex F.M. Bailey) EO (major components: limonene, *Z*-citral,  $\alpha$ -citral), lemon eucalyptus (*E. citriodora* Hook.) EO (major component:  $\beta$ -citronellal), and eucalyptus (*E. globulus* Labill.) EO (major component: 1,8-cineole) ( $EC_{50} = 0.59$ ,  $5.04$ , and  $7.78$  mg EO/mL, respectively). The superior toxicity of lemon ironbark is evident from these and other data and is due presumably to the activity of (major) components of its EO, which were not individually evaluated for biological activity [93].

In the above study, *Eucalyptus* spp. EOs were also shown to exhibit toxic effects by spraying directly onto *L. longipalpis* larvae. Thus, larvicidal effects were observed for *E. staigeriana*,



**Table 2** Median lethal concentrations (LC<sub>50</sub>) for volatile components of essential oils against larvae of mosquito species. Unless specified otherwise, all LC<sub>50</sub> values refer to a 24 h period of exposure. All LC<sub>50</sub> values are in ppm.

Volatile component	<i>Ae. aegypti</i>				<i>Ae. albopictus</i>					<i>Cx. pipiens</i> biotype <i>molestus</i>	<i>Cx. pipiens pallens</i>	<i>Oc. caspius</i>	<i>Oc. togoi</i>	Misc.
	[77]	[79]	[80]	[81]	[78]	[81]	[80]	[79]	[82]	[86]	[77]	[85]	[77]	
<i>Trans</i> -anethole									24/48 h	48 h		74.0/76.8		
$\beta$ -Asarone	27.0										22.4		26.4	
Benzaldehyde									47.0/45.4					
Borneol	94.9										91.6		97.3	
$\gamma$ -Cadinol									92.7/74.5					
Camphene	67.0										70.5		68.7	
3-Carene	19.2	27.9	25.3			24.1	22.9				13.8		16.2	
Carvacrol					70							35.5/34.6		
$\beta$ -Caryophyllene	88.3										93.7		97.9	
Caryophyllene oxide					125				65.6/58.3					
1,8-cineole	74.9										79.0		83.2	
<i>E</i> -Cinnamaldehyde									48.1/47.4					37.5/ 18.3 <sup>f</sup>
Cinnamyl acetate									52.7/45.0					
Citral									70.7/65.4					
<i>p</i> -Cymene		43.3	37.1	19.2		46.7	25.9	34.9						
Diallyl disulfide														6.61 <sup>d</sup>
3,5-Dimethoxytoluene	64.1										55.1		67.0	
Estragole	46.4										54.0		58.5	
Eugenol									67.4/54.8			7.53/5.57		
Fenchene	69.3										72.2		95.2	
16-Kaurene			57.0				56.5							
(+)-Limonene	24.5	19.4 <sup>b</sup>		18.1 <sup>b</sup>	37	32.7 <sup>b</sup>		15.0 <sup>b</sup>			13.3		19.2	
Linalool	96.6										94.8		99.0	
Linalyl acetate														23.1 <sup>e</sup>
Methyleugenol	57.7										53.3		58.5	
Myrcene	66.4	27.9 <sup>a</sup>	35.8 <sup>a</sup>			27.0 <sup>a</sup>	23.5 <sup>a</sup>				66.3		64.8	
Myristicin	73.0										77.0		90.7	
Pentadecane	96.7										97.6		99.2	
$\alpha$ -Phellandrene	23.1			16.6		39.9					13.8		16.1	
(+)- $\alpha$ -Pinene	50.9									61.5	54.0		47.3	
(-)- $\alpha$ -Pinene	64.8		79.1 <sup>c</sup>			74.0 <sup>c</sup>				58.4	70.4		57.9	
(+)- $\beta$ -Pinene	22.4									66.5	21.1		25.6	
(-)- $\beta$ -Pinene	15.4									36.5	12.9		18.0	
Safrole	9.88										8.22		16.1	
Terpinen-4-ol	64.8										58.3		58.5	
$\alpha$ -Terpinene			28.1	14.7		25.2	22.4							
$\gamma$ -Terpinene	17.1	26.8	26.8	30.7	95	29.8	22.8	22.8			12.6		14.4	
Terpinolene	15.3		32.1	28.4		35.6	22.0	21.3			11.9		14.2	
Thymol					79							33.7/33.3		
3,4,5-Trimethoxytoluene	67.1										74.8		91.4	
Verbenone	93.2										96.0		90.6	

<sup>a</sup>  $\beta$ -myrcene was the compound mentioned; <sup>b</sup> limonene was the compound mentioned; <sup>c</sup>  $\alpha$ -pinene was the compound mentioned. <sup>d</sup> *Cx. pipiens* third and fourth instars, 48 h LC<sub>50</sub> mg/L from *Allium sativum* [83]. <sup>e</sup> *Cx. pipiens* biotype *molestus* larvae, LC<sub>50</sub> mg/L from *Citrus × aurantium* [84]. <sup>f</sup> *Cx. quinquefasciatus* fourth instars, 24/48 h LC<sub>50</sub> mg/L from *Cinnamomum osmophloeum* [82]

*E. citriodora*, and *E. globulus* EOs (EC<sub>50</sub> = 2.63, 1.78, and 25.3 mg EO/mL, respectively) [93].

Dry, powdered, and otherwise unprocessed fruit and leaves of the broad-spectrum insectidal plants *A. indica* and *Melia azedarach* L. were tested in a no-choice feeding experiment against *L. longipalpis* first instars which were allowed to develop over a period of 30 days. All 4 types of extract had significant larvicidal effects as compared to controls fed an untreated, normal diet.

*A. indica* fruit extracts totally prevented third instars from moulting, thus resulting in no fourth instars. Feeding *M. azedarach* fruit totally prevented fourth instars of *L. longipalpis* from moulting (100% mortality) while feeding leaves of *M. azedarach* totally prevented moulting of second instars (100% mortality) [91]. Also, direct spraying of *Eucalyptus* spp. EOs onto *L. longipalpis* eggs produced ovicidal effects. Thus, *E. staigeriana*, *E. citriodora*,



**Table 3** Larvicidal activity of selected monoterpenoids and *trans*-anethole with and without piperonyl butoxide (PBO) to first- through fourth-instars of *Aedes aegypti* exposed for 24 h. All values are means of n = 3 experiments [87].

Chemical	Range of lethality to 1st–4th instars		
	without PBO	with PBO	Range of Synergism Ratios
	LC <sub>50</sub> mg/L	LC <sub>50</sub> mg/L	SR <sub>1–4</sub>
<i>Trans</i> -anethole	13–88.5	2.6–25.3	4–17
Borneol	183–>500	2.0–24.6	20–250
Borneol acetate	>500	3.0–39.3	13–167
Camphor	>500	2.1–71.8	7–238
1,8-Cineole	>500	2.1–96.2	5–238
Citronellal	40.7–263	2.9–15.6	14–37
<i>p</i> -Cymene	226–>500	5.2–23.2	22–44
Eugenol	24.5–143	2.3–52.3	3–11
Linalool	>500	2.0–99.5	5–250
$\alpha$ -Pinene	82.3–>500	2.8–20.7	24–151
$\beta$ -Pinene	96.2–>500	2.5–30.7	16–167
Pulegone	10.3–48.7	3.2–15.1	3–8
Terpineol	83.9–>500	3.2–8.5	26–128
Thymol	17.3–53.5	2.7–19.8	3–8

and *E. globulus* EOs are active mosquito ovicides (EC<sub>50</sub> = 3.60, 9.44, and 9.23 mg EO/mL, respectively) [93].

In a study performed in Ethiopia, *Phlebotomus bergeroti* Parrot adults (visceral leishmaniasis vector) were repelled by *A. indica* (neem) oil as 2 and 5% solutions in coconut oil providing 96.3% protection (for up to a mean time of 440 min) and 98.3% protection (for up to 9 h), respectively, under lab conditions. Also, *P. bergeroti* adults were repelled by *M. azedarach* (chinaberry) oil in 2 and 5% formulations in coconut oil which provided 95.1% protection (for 440 min) and 96.2% protection (for 500 min), respectively, under lab conditions. In tests against field populations of *Phlebotomus orientalis* Parrot and *P. bergeroti*, 2 and 5% neem oil in coconut oil mixtures and DEET provided more than 95% protection against *P. orientalis* for mean times of 8 h 24–48 min. Inter-

estingly, pure coconut oil also provided good protection (86%) against *P. orientalis* (no significant difference compared to test oil solutions and DEET) and providing a protection time of 504 min. Two concentrations of neem oil solution and pure coconut oil provided >92% protection for 480–576 min against *P. bergeroti*. DEET provided a lower mean protection of 76% against *P. bergeroti* for about 528 min (though this was not significantly different than test solution and coconut oil results). These results provide evidence for the value of both neem and coconut oils in the field as effective repellents to *Phlebotomus* Rondani & Berté spp. [94]. In other work, *P. papatasi* Scopoli mature females were repelled by garlic clove (*Allium sativum*) oil (1 and 0.005% oil exhibited 97 and 49% repellency, respectively) [95].

### Scabies mites

“Itch mite”, *Sarcoptes scabiei* var *hominis* Hering, is the arthropod agent of human scabies, a debilitating skin disease. The median lethal time (LT<sub>50</sub>) for *Sarcoptes scabiei* Linnaeus [96] treated with a 10% neem (*A. indica*) oil microemulsion was 81.7 min; with a 10% neem oil aqueous emulsion, 95.6 min; and with a microemulsion without neem oil, 89.1 min. The effectiveness of the microemulsion without neem oil is due to sodium dodecyl benzene sulfonate (SDBS) in the formulation which is known to have acaricidal activity in other studies [96]. Also, from the chloroform extracts of *A. indica* (neem) oil the acaricidal substance octadecanoic acid-tetrahydrofuran-3,4-diyl ester (24 h LC<sub>50</sub> = 0.1 mg/mL; LT<sub>50</sub> = 15.3 h at a concentration of 7.5 mg/mL) was isolated [97]. In other work, EOs and their components were screened for activity against permethrin-resistant (*S. scabiei* var *canis* Gerlach mites harvested from rabbits) and permethrin-sensitive (*S. scabiei* var *suis* Gerlach mites harvested from pigs) scabies mites, and active samples which were: clove EO (EC<sub>100</sub> ≥ 6.25% EO and 1.56% EO, respectively; mean survival time = 0.25 h), eugenol (EC<sub>50</sub> = 13.0 and 40.7 mM, respectively), isoeugenol (EC<sub>50</sub> = 24.6 and 32.1 mM, respectively), acetyleugenol (EC<sub>50</sub> = 19.4 and 30.8 mM, respectively), benzyl benzoate (positive control, EC<sub>50</sub> = 24.5 and 27.2 mM, respectively). Derivatives of eugenol and eugenol-rich clove EO show potential for use in the treatment of resistant and sensitive scabies mites [98].

**Table 4** Median lethal and knockdown data (fumigant activity) for chemical components of EOs against adult mosquitoes.

Chemical component	<i>Ae. aegypti</i>		Female [90]		<i>An. gambiae</i>		<i>An. tessellatus</i>		<i>Cx. quinquefasciatus</i>		
	Male & female				Male & female		Male & female		Male & female	Female	
	LD <sub>50</sub> [88]	KT <sub>50</sub> [89]	KD <sub>50</sub>	LC <sub>50</sub>	LD <sub>50</sub>		KD <sub>50</sub>	LC <sub>50</sub>	LD <sub>50</sub>	KD <sub>50</sub>	LC <sub>50</sub>
$\beta$ -Caryophyllene							1.1	0.80		2.2	5.3
1,8-Cineole		3.90									
<i>p</i> -Cymene		5.82									
Geijerene	6.8				4.2				5.4		
Germacrene D	2.8				1.8				2.1		
L-Menthol							0.54	0.36		0.50	0.50
Mugetanol			0.36	0.80			0.31	0.55		0.17	0.17
$\alpha$ -Pinene		5.36									
Pregeijerene	5.1				3.0				3.9		
Pulegone			3.3	5.3			0.84	1.33		1.62	3.31
$\gamma$ -Terpinene		9.31									
4-Terpineol		9.27									
$\alpha$ -Terpineol			0.75	0.62			0.59	0.45		0.59	0.56
Thymol			0.49	0.66			0.38	0.51		0.60	0.71

LD<sub>50</sub> – median lethal dose; KD<sub>50</sub> – median knockdown dose; KT<sub>50</sub> – median knockdown time; LC<sub>50</sub> – median lethal concentration. LD<sub>50</sub>, KD<sub>50</sub>, and LC<sub>50</sub> in  $\mu$ g/mL, KT<sub>50</sub> in min

## Ticks

Plant EOs and components exhibit significant adulticidal activities against ticks. Java citronella (*Cymbopogon winterianus* Jowitt ex Bor) EO exhibited lethal effects against *Rhipicephalus (Boophilus) microplus* Canestrini engorged females ( $DL_{50} = 6.1\%$  EO) [109]. Furthermore, *C. winterianus* EO can be sprayed or poured onto feeding ticks and causes a 45–75% reduction 22–28 days after treatment began [110]. The major components of *C. winterianus* were each tested against *R. microplus* and exhibited  $LC_{50}$  of 21.0% for citronellal and 17.8% for geraniol against blood-engorged females, and citronellol was much less toxic ( $LC_{50} = 78.9\%$ ) [110]. In other work, the EO (contains carvacrol) of the aerial parts of *Origanum minutiflorum* O. Schwarz & P.H. Davis is lethal to *R. turanicus* Pomerantsev adults, and 10  $\mu$ L EO/L causes 100% death in 120 min [111]. In another study, the EO of the aerial parts of *Lavandula angustifolia* exhibited toxicity to blood-engorged female *R. annulatus* ( $LC_{50}/LC_{99} = 2.76/8.84\%$  EO) [112].

The bioactive compound azadirachtin present in *A. indica* (neem) fruit and kernel extracts affects tick embryo development and molting stages. Interestingly, sheep (*Ovis aries* L.) which consumed a feed containing *A. indica* fruit and kernel extracts exhibited no noticeable signs of toxicity. This diet negatively affected the ability of the American dog tick *Dermacentor variabilis* Say to feed on sheep blood which exhibited plasma levels of azadirachtin of 4.35–4.81  $\mu$ g/mL over 14 days [113]. Sheep which consumed feed containing 0.6% azadirachtin had ticks which weighed ca. half as much as the ones from sheep which did not consume the *A. indica* extract. Azadirachtin in blood plasma impaired blood feeding by ticks, and so *A. indica* extracts as food additives may have applications in tick control for use in public health and veterinary applications [113].

Plant EOs and component chemicals exhibit significant larvicidal activities against ticks. In the larvae immersion test, *Hesperozygis ringens* (Benth.) Epling EO (86% pulegone) was lethal to *R. microplus* ( $LC_{99.9} = 0.541 \mu$ L/mL,  $LC_{50} = 0.260 \mu$ L/mL) as was pulegone ( $LC_{99.9} = 0.602 \mu$ L/mL,  $LC_{50} = 0.321 \mu$ L/mL) [106]. *C. winterianus* EO had lethal effects on *R. microplus* larvae ( $DL_{50} = 4.1\%$  EO) [109]. Plant EOs and components deter oviposition by ticks. For example, 5% geranium (*Pelargonium roseum* Willd.) leaf and stem EO and 100% *Cymbopogon nardus* (L.) Rendle leaf EO exhibited 88 and 100% oviposition deterrent effects on *R. annulatus* Say blood-engorged females [107, 108]. Also, *C. winterianus* EO exhibited total oviposition deterrence ( $EC_{100} = 10\%$  EO) [109], and at a concentration of 20%, *C. winterianus* EO exhibited total oviposition deterrence in *R. microplus* blood-engorged females [110].

Plant EOs and component chemicals exhibit anti-hatching activity against ticks. *Hesperozygis ringens* (Benth.) Epling EO (86% pulegone) was tested at concentrations of 25 and 50  $\mu$ L/mL in the adult immersion test and exhibited inhibitions of 48 and 76%, respectively, in egg production by *R. microplus* blood-engorged females as compared to controls. The eggs which were produced by these treated females exhibited infertility (30–95% anti-hatching activity). Pure pulegone gave the same results as EO in the above tests [106]. Furthermore, *C. nardus* EO had 100% anti-hatching activity against *R. annulatus* [108]. Similarly, *C. winterianus* EO exhibited total anti-hatching activity on *R. annulatus* eggs ( $EC_{100} = 7.1\%$  EO) [109]. The major chemical components of *C. winterianus* were each tested against *R. microplus*, and 50% citronellal and 25% geraniol caused 100% sterility of eggs (no hatching) [110]. In another study, the EO of the aerial parts of *L. angustifolia* was tested against *R. annulatus* blood-engorged females. At

4.0% EO, female *R. annulatus* suffered a 100% failure to produce eggs and  $\geq 6\%$  EO provided 100% egg laying failure [112].

Several recent applications of plant extracts and isolated natural compounds as tick repellents have been described in the scientific literature. In the wrapped finger assay, the ethanol extracts of the aerial parts of *C.nardus* ( $EC_{50} = 0.089 \text{ mg/cm}^2$ ), the leaves of *Ageratum conyzoides* ( $EC_{50} = 0.205 \text{ mg/cm}^2$ ), and the isolated natural compound callicarpinal ( $EC_{50} = 0.084 \text{ mg/cm}^2$ ) (from the leaves of *Callicarpa americana* L.) all exhibited significant repellent effects against the dog tick *Amblyomma cajennense* Fabricius [99]. Also, the gum resin (composed of germacrene D,  $\delta$ -elemene,  $\beta$ -bourbonene) of *Commiphora holtziana* Engl. provided over 5 h of repellent protection against *R. microplus* adults [100]. In other work, *Ixodes ricinus* L. nymphs were repelled by the toluene extracts of the twigs and leaves of *Artemisia abrotanum* L. (57% repellency after 8 h) and by the active chemical components thujyl alcohol, eugenol, and coumarin (83, 98, and 98% repellency, respectively, after 8 h) [101].

Chemical components of plant EOs are a very interesting group of substances having confirmed efficacy in tick control. For example, *Ixodes ricinus* nymphs are repelled by carnation (*Dianthus caryophyllus* L.) EO (92% repellency after 8 h). The active components present in *D. caryophyllus* flower EO were 2-phenylethanol,  $\beta$ -citronellol, cinnamyl alcohol, geraniol, and  $\alpha$ -pinene which when tested individually exhibited 96, 84, 80, 79, and 75% repellency, respectively, against *I. ricinus* after 8 h [101].

Undecanone is a valuable tick repellent which originally derived from the wild tomato plant, *Lycopersicon hirsutum* fo. *glabratum* C.H. Mull. In assays involving treated vs. untreated cotton cheesecloth and direct comparison, a 7.75% 2-undecanone commercial formulation exhibited mean percentage repellency against the lone star tick *Amblyomma americanum* L. and *D. variabilis* which was greater than or comparable in efficacy to arthropod repellent products containing 98.1% N,N-diethyl-m-toluamide (DEET), 19.6% IR3535, and 30% lemon eucalyptus (*Eucalyptus citriodora*) EO. Products containing 5% and 15% picaridin and 0.5% permethrin were less repellent than the 7.75% 2-undecanone formulation [102]. The high repellency of a commercial formulation containing 7.75% 2-undecanone as the active ingredient against *D. variabilis* and blacklegged tick *Ixodes scapularis* Say was confirmed in other work [103].

Brown ear tick *Rhipicephalus appendiculatus* Neumann adult climbing is repelled by the EO of *Commiphora swynnertonii* Burt, and a 0.1% solution of  $\alpha$ -copaene provided 86% repellency while a 1% solution had repellency comparable to DEET. Isocaryophyllene also exhibited repellent activity against adult *R. appendiculatus* (10% solution, 55% repellency) [100].

In other work, *Amyris balsamifera* L. EO ( $EC_{50} = 9.0$  and  $22.9 \mu$ g/cm<sup>2</sup>, respectively) and elemol ( $EC_{50} = 10.9$  and  $14.8 \text{ nmol/cm}^2$ , respectively), a major chemical constituent of the EO of osage orange [*Maclura pomifera* (Raf.) C.K. Schneid.], effectively repelled *A. americanum* host-seeking nymphs in climbing filter paper and wrapped fingertip assays, and elemol exhibited comparable repellency to DEET. Elemol and amyris EO also effectively repelled *I. scapularis* ( $EC_{50} = 5.16$  and  $4.20 \mu$ g/cm<sup>2</sup>, respectively, in the wrapped fingertip assay) [104]. A common component of plant EOs, geraniol, in mixtures containing cinnamon (*Cinnamomum* Schaeff. sp.), lemongrass (*Cymbopogon citratus* (DC.) Stapf), rosemary (*Rosmarinus officinalis* L.), wintergreen (*Gaultheria procumbens* L.), and canola oils as topical repellents exhibits longevity and repellency comparable to or better than DEET against *D. variabilis* and *I. scapularis* [105].

## Conclusion

Much information has been made available in the past few years on active plant extracts, fractions, EOs, and their isolated components which are responsible for lethal effects against arthropods. Plant extracts or chemicals which owe their origins to plants such as *A. indica* (neem) oils and other derivatives, azadirachtin, undecanone, among others already have important commercial applications in a variety of commercial products which are useful for arthropod control. It is of major importance for future work that more importance be given to two general areas of research. One area involves the probable differences in enantiomeric purity of EO components which are purchased and used in bioassays to confirm the active anti-arthropod activity in many of the references cited and the enantiomeric purity of said component in the active EO. In future work involving chiral monoterpenes, the enantiomeric purity of the active component in the EO and in the purchased reference sample should be taken into account. Another area of importance is the deciphering of the synergistic, suppressive, and other interactions of the components of EOs (and extracts) as was done in the complex and stimulating work by Yoon et al. [33] for the components of *Citrus* L. EOs against several cockroach species. In many of the scientific papers reviewed herein, the isolated active components are recognizably less active than the plant extracts and EOs from which they were isolated, and so doubt remains whether the component is the most important active component or whether synergism or perhaps a "cocktail of components" is in fact the active agent responsible for the effect of the plant derivative. For many arthropod vectors, besides phytochemicals which are already found in commercial products, there are at least a few promising botanical candidates for control or development of control products in the future.

## Supporting information

In Supporting Information, URLs of documents on human diseases borne by arthropod vectors, data from literature sources on plant extracts, EOs, monoterpenes, and phenylpropanoids exhibiting fumigant and other activities against *M. domestica* and extracts, EOs and fractions exhibiting lethal effects to mosquitoes are presented.

## References

- 1 Graczyk TK, Knight R, Tamang L. Mechanical transmission of human protozoan parasites by insects. *Clin Microbiol Rev* 2005; 18: 128–132
- 2 Sharma P, Sharma JD. A review of plant species assessed *in vitro* for antiamebic activity or both antiamebic and antiplasmodial properties. *Phytother Res* 2001; 15: 1–17
- 3 Rassi Jr A, Marin-Neto JA. Chagas disease. *Lancet* 2010; 375: 449
- 4 Carod-Artal FJ, Gascon J. Chagas disease and stroke. *Lancet Neurol* 2010; 9: 533
- 5 Costa J. The synanthropic process of Chagas disease vectors in Brazil, with special attention to *Triatoma brasiliensis* Neiva, 1911 (Hemiptera, Reduviidae, Triatominae) population, genetical, ecological, and epidemiological aspects. *Mem Inst Oswaldo Cruz* 1999; 94: 239–241
- 6 Fernandes AJ, Chiari E, Casanova C, Dias JCP, Romanha AJ. The threat of reintroduction of natural transmission of Chagas' disease in Bambuí, Minas Gerais state, Brazil, due to *Panstrongylus megistus*. *Mem Inst Oswaldo Cruz* 1992; 87: 285–289
- 7 World Health Organization (WHO). Specific document URLs are provided in the supplementary data. Available at <http://www.who.int/>. Accessed September 16, 2010
- 8 Fotedar R. Vector potential of houseflies (*Musca domestica*) in the transmission of *Vibrio cholerae* in India. *Acta Tropica* 2001; 78: 31–34
- 9 Bechah Y, Capo C, Mege JL, Raoult D. Epidemic typhus. *Lancet Infect Dis* 2008; 8: 417–426
- 10 Center for Disease Control and Prevention (CDC). Specific document URLs are provided in the supplementary data (available online). Available at <http://www.cdc.gov/>. Accessed September 16, 2010
- 11 Doiz O, Clavel A, Morales S, Varea M, Castillo FJ, Rubio C, Gomez-Lus R. House fly (*Musca domestica*) as a transport vector of *Giardia lamblia*. *Folia Parasitol* 2000; 47: 330–331
- 12 Minnaganti VR. Isosporiasis. Medscape: eMedicine. Available at <http://emedicine.medscape.com/article/219776-overview>. Accessed September 16, 2010
- 13 Leishmaniasis. NIH. Available at <http://www.nlm.nih.gov/medlineplus/leishmaniasis.html#cat59>. Accessed September 16, 2010
- 14 Soares CO, Ishikawa MM, Fonseca AH, Yoshinari NH. Borreliose, agentes e vetores. *Pesq Vet Bras* 2000; 20: 1–19
- 15 Alvim NC, Bento MAF, Martins LA. Borreliose de Lyme – a doença da década. *Rev Cient Elet Med Vet* 2005; 4 – ISSN 1679–7353 (online). Available at <http://www.revista.inf.br/veterinaria04/revissao/revissao04.pdf>. Accessed September 16, 2010
- 16 The Center for Food Security and Public Health. Zoonotic diseases by routes of transmission: companion animals. Iowa State University (ISU). Available at <http://www.cfsph.iastate.edu/>. Accessed September 16, 2010
- 17 Tzanetou K. Scabies: reappearance of a forgotten parasitic disease. *Deltio Ilenikes Mikrobiologikes Etaireias* 2006; 51: 346–352 (Chemical Abstracts)
- 18 Angerami RN, Nunes EM, Nascimento EMM, Freitas AR, Kemp B, Feltrin AFC, Pacola MR, Perecin GE, Sinkov V, Resende MR, Katz G, Jacintho da Silva L. Clusters of Brazilian spotted fever in São Paulo State, southeastern Brazil. A review of official reports and the scientific literature. *Eur Soc Clin Microbiol Infect Dis* 2009; 15: 202–204
- 19 Shaalan EAS, Canyon D, Younes MWF, Abdel-Wahab H, Mansour AH. A review of botanical phytochemicals with mosquitoicidal potential. *Environ Int* 2005; 31: 1149–1166
- 20 Bakkali F, Averbeck S, Averbeck D, Idaomar M. Biological effects of essential oils – a review. *Food Chem Toxicol* 2008; 46: 446–475
- 21 Burfield T, Reekie SL. Mosquitoes, malaria and essential oils. *Int J Aromather* 2005; 15: 30–41
- 22 Isman MB, Machial CM. Pesticides based on plant essential oils: from traditional practice to commercialization. In: Rai M, Carpinella MC, editors. Naturally occurring bioactive compounds. *Adv Phytomed* 2006; 3: 29–44
- 23 Nerio LS, Olivero-Verbel J, Stashenko E. Repellent activity of essential oils: a review. *Bioresource Technol* 2010; 101: 372–378
- 24 Pohlit AM, Lopes NP, Gama RA, Tadei WP, Andrade-Neto VF. Patent literature on mosquito repellent inventions which contain plant essential oils – a review. *Planta Med*, advance online publication 15 February 2011; doi: 10.1055/s-0030-1270723
- 25 Marten GG. Larvicidal algae. In: Floore TG, editor. Biorational Control of Mosquitoes. *Am Mosq Ctrl Assoc Bull* 2007; 23 (Suppl. 2): 177–183
- 26 Matur BM, Davou BJ. Comparative larvicidal property of leaf extract of *Chromolaena odorata* L. (Compositae) and chlopyrifos (organophosphorus compound) on *Simulium* larvae. *Biomed Environ Sci* 2007; 20: 313–316
- 27 Spero NC, Gonzalez YI, Scialdone MA, Hallahan DL. Repellency of hydrogenated catmint oil formulations to black flies and mosquitoes in the field. *J Med Entomol* 2008; 45: 1080–1086
- 28 Twatsin A, Thavara U, Chansang U, Chavalittumrong P, Boonruad T, Wongsinkongman P, Bansidhi J, Mulla MS. Field evaluation of deet, Repel Care®, and three plant-based essential oil repellents against mosquitoes, black flies (Diptera: Simuliidae), and land leeches (Arhynchobdellida: Haemadipsidae) in Thailand. *J Am Mosq Ctrl Assoc* 2006; 22: 306–313
- 29 Sukontason KL, Boonchu N, Sukontason K, Choochote W. Effects of eucalyptol on house fly (Diptera: Muscidae) and blow fly (Diptera: Calliphoridae). *Rev Inst Med Trop S Paulo* 2004; 46: 97–101
- 30 Siri wattanarungsee S, Sukontason KL, Olson JK, Chailapakul O, Sukontason K. Efficacy of neem extract against the blowfly and housefly. *Parasitol Res* 2008; 103: 535–544
- 31 Kumarasinghe SPW, Karunaweera ND, Ihalamulla RL, Arambewela LSR, Dissanayake RDSCT. Larvicidal effects of mineral turpentine, low aromatic white spirits, aqueous extracts of *Cassia alata*, and aqueous extracts, ethanolic extracts and essential oil of betel leaf (*Piper betle*) on *Chrysomya megacephala*. *International J Dermatol* 2002; 41: 877–880
- 32 Shen LR, Li HY, Zhou YG, Gu S, Lou YG. Ovicidal activity of nine essential oils against *Chrysomya megacephala* in bacon and kipper. *Chin J Appl Ecol* 2007; 18: 2343–2346



- 33 Yoon C, Kang SH, Yang JO, Noh DJ, Indiragandhi P, Kim GH. Repellent activity of citrus oils against the cockroaches *Blattella germanica*, *Periplaneta americana* and *P. fuliginosa*. J Pestic Sci 2009; 34: 77–88
- 34 Phillips AK, Appel AG, Sims SR. Topical toxicity of essential oils to the German cockroach (Dictyoptera: Blattellidae). J Econ Entomol 2010; 103: 448–459
- 35 Bisseleua HBD, Gbewonyo SWK, Obeng-Ofori D. Toxicity, growth regulatory and repellent activities of medicinal plant extracts on *Musca domestica* L. (Diptera: Muscidae). African J Biotechnol 2008; 7: 4635–4642
- 36 Moreira MD, Picanço MC, Barbosa LCA, Guedes RNC, Barros EC, Campos MR. Compounds from *Ageratum conyzoides*: isolation, structural elucidation and insecticidal activity. Pest Manag Sci 2007; 63: 615–621
- 37 Huang JG, Zhou LJ, Xu HH, Li WO. Insecticidal and cytotoxic activities of extracts of *Cacalia tangutica* and its two active ingredients against *Musca domestica* and *Aedes albopictus*. J Econ Entomol 2009; 102: 1444–1447
- 38 Tarelli G, Zerba EN, Alzogaray RA. Toxicity to vapor exposure and topical application of essential oils and monoterpenes on *Musca domestica* (Diptera: Muscidae). J Econ Entomol 2009; 102: 1383–1388
- 39 Palacios SM, Bertoni A, Rossi Y, Santander R, Urzúa A. Efficacy of essential oils from edible plants as insecticides against the house fly, *Musca domestica* L. Molecules 2009; 14: 1938–1947
- 40 Mohottalage S, Tabacchi R, Guerin PM. Components from Sri Lankan *Piper betle* L. leaf oil and their analogues showing toxicity against the housefly, *Musca domestica*. Flavour Fragr J 2007; 22: 130–138
- 41 Dolan MC, Dietrich G, Panella NA, Monteneri JA, Karchesy JJ. Bicidal activity of three wood essential oils against *Ixodes scapularis* (Acari: Ixodidae), *Xenopsylla cheopis* (Siphonaptera: Pulicidae), and *Aedes aegypti* (Diptera: Culicidae). J Econ Entomol 2007; 100: 622–625
- 42 Panella NA, Dolan MC, Karchesy JJ, Xiong Y, Peralta-Cruz J, Khasawneh M, Monteneri JA, Maupin GO. Use of novel compounds for pest control: insecticidal and acaricidal activity of essential oil components from heartwood of Alaska yellow cedar. J Med Entomol 2005; 42: 352–358
- 43 Coelho AAM, Paula JE, Espíndola LS. Insecticidal activity of cerrado plant extracts on *Rhodnius milesi* Carcavallo, Rocha, Galvão & Jurberg (Hemiptera: Reduviidae), under laboratory conditions. Neotrop Entomol 2006; 35: 133–138
- 44 Mello CB, Uzeda CD, Bernardino MV, Mendonça-Lopes D, Kelecom A, Fevereiro PCA, Guerra MS, Oliveira AP, Rocha LM, Gonzalez MS. Effects of the essential oil obtained from *Pilocarpus spicatus* Saint-Hilaire (Rutaceae) on the development of *Rhodnius prolixus* nymphae. Rev Bras Farmacogn [Braz J Pharmacogn] 2007; 17: 514–520
- 45 Sfara V, Zerba EN, Alzogaray RA. Fumigant insecticidal activity and repellent effect of five essential oils and seven monoterpenes on first-instar nymphs of *Rhodnius prolixus*. J Med Entomol 2009; 46: 511–515
- 46 Ferrero AA, González JOW, Chopa CS. Biological activity of *Schinus molle* on *Tritoma infestans*. Fitoterapia 2006; 77: 381–383
- 47 Jadhav V, Kore A, Kadam VJ. In-vitro pediculicidal activity of *Hedychium spicatum* essential oil. Fitoterapia 2007; 78: 470–473
- 48 Ashok VA, Srinivas A, Ravitej K, Gopinadh V. Ethyl acetate extract of *Annona squamosa* seeds containing anti-head lice activity. Pharmacogn J 2009; 1: 207–209
- 49 Picollo MI, Toloza AC, Cueto GM, Zygodlo J, Zerba E. Anticholinesterase and pediculicidal activities of monoterpenoids. Fitoterapia 2008; 79: 271–278
- 50 Toloza AC, Vassena C, Picollo MI. Ovicidal and adulticidal effects of monoterpenoids against permethrin-resistant human head lice, *Pediculus humanus capitis*. Med Vet Entomol 2008; 22: 335–339
- 51 Dua VK, Pandey AC, Raghavendra K, Gupta A, Sharma T, Dash AP. Larvicidal activity of neem oil (*Azadirachta indica*) formulation against mosquitoes. Malaria J 2009; 8: 124 [online, 6 pp]
- 52 Howard AFV, Adongo EA, Hassanali A, Omlin FX, Wanjoya A, Zhou G, Vulule J. Laboratory evaluation of the aqueous extract of *Azadirachta indica* (Neem) wood chippings on *Anopheles gambiae* s.s. (Diptera: Culicidae) mosquitoes. J Med Entomol 2009; 46: 107–114
- 53 Kiprok AK, Kiprono PC, Rajab MS, Kosgei MK. Limonoids as larvicidal components against mosquito larvae (*Aedes aegypti* Linn.). Z Naturforsch C 2007; 62: 826–828
- 54 Silva ECC, Cavalcanti BC, Amorim RCN, Lucena JF, Quadros DS, Tadei WP, Montenegro RC, Costa-Lotufu LV, Pessoa C, Moraes MO, Nunomura RCS, Nunomura SM, Melo MRS, Andrade-Neto VF, Silva LFR, Vieira PPR, Pohlit AM. Biological activity of neosergeolide and isobrucein B (and two semi-synthetic derivatives) isolated from the Amazonian medicinal plant *Picrolemma sprucei* (Simaroubaceae). Mem Inst Oswaldo Cruz 2009; 104: 48–55
- 55 Gu HJ, Cheng SS, Huang CG, Chen WJ, Chang ST. Mosquito larvicidal activities of extracts from black heartwood-type *Cryptomeria japonica*. Parasitol Res 2009; 105: 1455–1458
- 56 Madhu SK, Shaikath AK, Vijayan VA. Efficacy of bioactive compounds from *Curcuma aromatica* against mosquito larvae. Acta Trop 2010; 113: 7–11
- 57 Silva HHG, Geris R, Rodrigues-Filho E, Rocha C, Silva IG. Larvicidal activity of oil-resin fractions from the Brazilian medicinal plant *Copaifera reticulata* Ducke (Leguminosae-Caesalpinioideae) against *Aedes aegypti* (Diptera, Culicidae). Rev Soc Bras Med Trop 2007; 40: 264–267
- 58 Geris R, Silva IG, Silva HHG, Barison A, Rodrigues-Filho E, Ferreira AG. Di-terpenoids from *Copaifera reticulata* Ducke with larvicidal activity against *Aedes aegypti* (L.) (Diptera, Culicidae). Rev Inst Med Trop S Paulo 2008; 50: 25–28
- 59 Rahuman AA, Venkatesan P, Geetha K, Gopalakrishnan G, Bagavan A, Kamaraj C. Mosquito larvicidal activity of gluanol acetate, a tetracyclic triterpene derived from *Ficus racemosa* Linn. Parasitol Res 2008; 103: 333–339
- 60 Innocent E, Joseph CC, Gikonyo NK, Moshi MJ, Nkunya MHH, Hassanali A. Mosquito larvicidal constituents from *Lantana viburnoides* sp *viburnoides* var. *livi* (A. Rich) Verdc (Verbenaceae). J Vector Borne Dis 2008; 45: 240–244
- 61 Rahuman AA, Gopalakrishnan G, Venkatesan P, Geetha K. Isolation and identification of mosquito larvicidal compound from *Abutilon indicum* (Linn.) Sweet. Parasitol Res 2008; 102: 981–988
- 62 Bagavan A, Rahuman AA, Kamaraj C, Geetha K. Larvicidal activity of saponin from *Achyranthes aspera* against *Aedes aegypti* and *Culex quinquefasciatus* (Diptera: Culicidae). Parasitol Res 2008; 103: 223–229
- 63 Ghosh A, Chowdhury N, Chandra G. Laboratory evaluation of a phyto-steroid compound of mature leaves of day jasmine (Solanaceae: Solanales) against larvae of *Culex quinquefasciatus* (Diptera: Culicidae) and nontarget organisms. Parasitol Res 2008; 103: 271–277
- 64 Cheng SS, Huang CG, Chen WJ, Kuo YH, Chang ST. Larvicidal activity of tectoquinone isolated from red heartwood-type *Cryptomeria japonica* against two mosquito species. Bioresour Technol 2008; 99: 3617–3622
- 65 Georges K, Jayaprakasam B, Dalavoy SS, Nair MG. Pest-managing activities of plant extracts and anthraquinones from *Cassia nigricans* from Burkina Faso. Bioresour Technol 2008; 99: 2037–2045
- 66 Sreelatha T, Hymavathi A, Murthy JM, Rani PU, Rao JM, Babu KS. Bioactivity-guided isolation of mosquitoicidal constituents from the rhizomes of *Plumbago capensis* Thunb. Bioorg Med Chem Lett 2010; 20: 2974–2977
- 67 Maniava BM, Wilber L, Ndiege IO, Wanjala CC, Akenga TA. Larvicidal activity of extracts from three *Plumbago* spp against *Anopheles gambiae*. Mem Inst Oswaldo Cruz 2009; 104: 813–817
- 68 Garcez WS, Garcez FR, da Silva LMGE, Hamerski L. Larvicidal activity against *Aedes aegypti* of some plants native to the West-Central region of Brazil. Bioresour Technol 2009; 100: 6647–6650
- 69 Rahuman AA, Venkatesan P, Gopalakrishnan G. Mosquito larvicidal activity of oleic and linoleic acids isolated from *Citrullus colocynthis* (Linn.) Schrad. Parasitol Res 2008; 103: 1383–1390
- 70 Coelho JS, Santos NDL, Napoleão TH, Gomes FS, Ferreira RS, Zingali RB, Coelho LCB, Leite SP, Navarro DMAF, Paiva PMG. Effect of *Moringa oleifera* lectin on development and mortality of *Aedes aegypti* larvae. Chemosphere 2009; 77: 934–938
- 71 Sá RA, Santos NDL, Silva CSB, Napoleão TH, Gomes FS, Cavada BS, Coelho LC, Navarro DM, Bieber LW, Paiva PM. Larvicidal activity of lectins from *Myracrodruon urundeuva* on *Aedes aegypti*. Comp Biochem Physiol C Pharmacol Toxicol Endocrinol 2009; 149: 300–306
- 72 Chowdhury N, Laskar S, Chandra G. Mosquito larvicidal and antimicrobial activity of protein of *Solanum villosum* leaves. BMC Complement Alternat Med 2008; 8: 62 [online, 6 pp]
- 73 Vasconcelos JN, Lima JQ, Lemos TLG, Oliveira MCF, Almeida MMB, Andrade-Neto M, Mafezoli J, Arriaga AMC, Santiago GMP, Braz-Filho R. Estudo químico e biológico de *Tephrosia toxicaria* Pers. Quim Nova 2009; 32: 382–386
- 74 Aivazi AA, Vijayan VA. Larvicidal activity of oak *Quercus infectoria* Oliv. (Fagaceae) gall extracts against *Anopheles stephensi* Liston. Parasitol Res 2009; 104: 1289–1293
- 75 Pohlit AM, Quignard ELJ, Nunomura SM, Tadei WP, Hidalgo AF, Pinto ACS, Santos EVM, Morais SKR, Saraiva RCG, Ming LC, Alecrim AM, Ferraz AB, Pedrosa ACS, Diniz EV, Finney EK, Gomes EO, Dias HB, Souza KS, Oliveira LCP, Don LC, Queiroz MMA, Henrique MC, Santos M, Lacerda-Junior OS, Pinto OS, Silva SG, Graça YR. Screening of plants found in the State of



- Amazonas, Brazil for larvicidal activity against *Aedes aegypti* larvae. *Acta Amazonica* 2004; 34: 97–105
- 76 Pohlit AM, Pinto ACS, Cavalcanti BC, Pessoa CO, Silva ECCS, Chaves FCM, Nogueira KL, Lotufo L, Silva LFR, Melo MRS, Vieira PPR, Mause R, Amorim RCN, Tadei WP, Andrade-Neto V. Biopharmaceutical synthesis. In: Carioca JOB, organizer. Brazilian network on green chemistry-awareness, responsibility and action. Fortaleza, Brasil: Edições Universidade do Ceará (UFC); 2008: 347–372
  - 77 Perumalsamy H, Kim N-J, Ahn Y-J. Larvicidal activity of compounds isolated from *Asarum heterotropoides* against *Culex pipiens pallens*, *Aedes aegypti*, and *Ochlerotatus togoi* (Diptera: Culicidae). *J Med Entomol* 2009; 46: 1420–1423
  - 78 Silva WJ, Dória GAA, Maia RT, Nunes RS, Carvalho GA, Blank AF, Alves PB, Marçal RM, Cavalcanti SCH. Effects of essential oils on *Aedes aegypti* larvae: alternatives to environmentally safe insecticides. *Bioresour Technol* 2008; 99: 3251–3255
  - 79 Cheng SS, Chang HT, Lin CY, Chen PS, Huang CG, Chen WJ, Chang ST. Insecticidal activities of leaf and twig essential oils from *Clausena excavata* against *Aedes aegypti* and *Aedes albopictus* larvae. *Pest Manag Sci* 2009; 65: 339–343
  - 80 Cheng SS, Chua MT, Chang EH, Huang CG, Chen WJ, Chang ST. Variations in insecticidal activity and chemical compositions of leaf essential oils from *Cryptomeria japonica* at different ages. *Bioresour Technol* 2009; 100: 465–470
  - 81 Cheng SS, Huang CG, Chen YJ, Yu JJ, Chen WJ, Chang ST. Chemical compositions and larvicidal activities of leaf essential oils from two eucalyptus species. *Bioresour Technol* 2009; 100: 452–456
  - 82 Cheng SS, Liu JY, Huang CG, Hsui YR, Chen WJ, Chang ST. Insecticidal activities of leaf essential oils from *Cinnamomum osmophloeum* against three mosquito species. *Bioresour Technol* 2009; 100: 457–464
  - 83 Kimbaris AC, Kioulos E, Koliopoulos G, Polissiou MG, Michaelakis A. Coactivity of sulfide ingredients: a new perspective of the larvicidal activity of garlic essential oil against mosquitoes. *Pest Manag Sci* 2009; 65: 249–254
  - 84 Melliou E, Michaelakis A, Koliopoulos G, Skaltsounis AL, Magiatis P. High quality bergamot oil from Greece: chemical analysis using chiral gas chromatography and larvicidal activity against the West Nile virus vector. *Molecules* 2009; 14: 839–849
  - 85 Knio KM, Usta J, Dagher S, Jounajian H, Kreydiyyeh S. Larvicidal activity of essential oils extracted from commonly used herbs in Lebanon against the seaside mosquito, *Ochlerotatus caspius*. *Bioresour Technol* 2008; 99: 763–768
  - 86 Michaelakis A, Papachristos D, Kimbaris A, Koliopoulos G, Giatropoulos A, Polissiou MG. Citrus essential oils and four enantiomeric pinenes against *Culex pipiens* (Diptera: Culicidae). *Parasitol Res* 2009; 105: 769–773
  - 87 Waliwitiya R, Kennedy CJ, Lowenberger CA. Larvicidal and oviposition-altering activity of monoterpenoids, *trans*-anethole and rosemary oil to the yellow fever mosquito *Aedes aegypti* (Diptera: Culicidae). *Pest Manag Sci* 2009; 65: 241–248
  - 88 Kiran SR, Devi PS. Evaluation of mosquitocidal activity of essential oil and sesquiterpenes from leaves of *Chloroxylon swietenia* DC. *Parasitol Res* 2007; 101: 413–418
  - 89 Lucia A, Licastro S, Zerba E, Audino PG, Masuh H. Sensitivity of *Aedes aegypti* adults (Diptera: Culicidae) to the vapors of *Eucalyptus* essential oils. *Bioresour Technol* 2009; 100: 6083–6087
  - 90 Samarasekera R, Weerasinghe IS, Hemalal KDP. Insecticidal activity of menthol derivatives against mosquitoes. *Pest Management Science* 2008; 64: 290–295
  - 91 Andrade-Coelho CA, Souza NA, Gouveia C, Silva VC, Gonzalez MS, Rangel EF. Effect of fruit and leaves of Meliaceae plants (*Azadirachta indica* and *Melia azedarach*) on the development of *Lutzomyia longipalpis* larvae (Diptera: Psychodidae: Phlebotominae) under experimental conditions. *J Med Entomol* 2009; 46: 1125–1130
  - 92 Luitgards-Moura JF, Bermudez EGC, Rocha AFI, Tsouris P, Rosa-Freitas MG. Preliminary assays indicate that *Antonia ovata* (Loganiaceae) and *Derris amazonica* (Papilionaceae), ichthyotoxic plants used for fishing in Roraima, Brazil, have an insecticide effect on *Lutzomyia longipalpis* (Diptera: Psychodidae: Phlebotominae). *Mem Inst Oswaldo Cruz* 2002; 97: 737–742
  - 93 Maciel MV, Morais SM, Bevilaqua CML, Silva RA, Barros RS, Sousa RN, Sousa LC, Brito ES, Souza-Neto MA. Chemical composition of *Eucalyptus* spp. essential oils and their insecticidal effects on *Lutzomyia longipalpis*. *Vet Parasitol* 2010; 167: 1–7
  - 94 Kebede Y, Gebre-Michael T, Balkew M. Laboratory and field evaluation of neem (*Azadirachta indica* A. Juss) and Chinaberry (*Melia azedarach* L.) oils as repellents against *Phlebotomus orientalis* and *P. bergerochi* (Diptera: Psychodidae) in Ethiopia. *Acta Trop* 2010; 113: 145–150
  - 95 Valerio L, Maroli M. Evaluation of repellent and anti-feeding effect of garlic oil (*Allium sativum*) against the bite of phlebotomine sandflies (Diptera: Psychodidae). *Annali dell'Istituto Superiore di Sanita* 2005; 41: 253–256
  - 96 Xu J, Fan QJ, Yin ZQ, Li XT, Du YH, Jia RY, Wang KY, Lv C, Ye G, Geng Y, Su G, Zhao L, Hua TX, Shi F, Zhang L, Wu CL, Tao C, Zhang YX, Shi DX. The preparation of neem oil microemulsion (*Azadirachta indica*) and the comparison of acaricidal time between neem oil microemulsion and other formulations *in vitro*. *Vet Parasitol* 2010; 169: 399–403
  - 97 Du YYH, Li JL, Jia RY, Yin ZQ, Li XT, Lv C, Ye G, Zhang L, Zhang YQ. Acaricidal activity of four fractions and octadecanoic acid-tetrahydrofuran-3,4-diyl ester isolated from chloroform extracts of neem (*Azadirachta indica*) oil against *Sarcoptes scabiei* var. *cuniculi* larvae *in vitro*. *Vet Parasitol* 2009; 163: 175–178
  - 98 Pasay C, Mounsey K, Stevenson G, Davis R, Arlian L, Morgan M, Vyszenski-Moher D, Andrews K, McCarthy J. Acaricidal activity of eugenol based compounds against scabies mites. *PLoS One* 2010; 5: 1–9
  - 99 Soares SF, Borge LMF, Braga RS, Ferreira LL, Louly CCB, Tresvenzol MF, Paula JR, Ferri PH. Repellent activity of plant-derived compounds against *Amblyomma cajennense* nymphs. *Vet Parasitol* 2010; 167: 67–73
  - 100 Kaoneka B, Mollel M, Lyatuu F. Leaf essential oil composition and tick repellency activity of *Commiphora swynnertonii* Burt. *J Biol Res Thessalon* 2007; 8: 213–216
  - 101 Tunón H, Thorsell W, Mikiver A, Malander I. Arthropod repellency, especially tick (*Ixodes ricinus*), exerted by extract from *Artemisia abrotanum* and essential oil from flowers of *Dianthus caryophyllum*. *Fito-terapia* 2006; 77: 257–261
  - 102 Bissinger BW, Zhu J, Apperson CS, Sonenshine DE, Watson DW, Roe RM. Comparative efficacy of BioUD to other commercially available arthropod repellents against the ticks *Amblyomma americanum* and *Dermacentor variabilis* on cotton cloth. *Am J Trop Med Hyg* 2009; 81: 685–690
  - 103 Witting-Bissinger BE, Stumpf CF, Donohue KV, Apperson CS, Roe RM. Novel arthropod repellent, BioUD, is an efficacious alternative to deet. *J Med Entomol* 2008; 45: 891–898
  - 104 Carrol JF, Paluch G, Coats J, Kramer M. Elemol and amyris oil repel the ticks *Ixodes scapularis* and *Amblyomma americanum* (Acari: Ixodidae) in laboratory bioassays. *Exp Appl Acarol* 2010; 51: 383–392
  - 105 Besette SM. Pest repellent compositions containing geraniol and essential oils and methods for use. US Patent 2009263515; 2009
  - 106 Ribeiro VLS, Santos JC, Bordignon SAL, Apel MA, Henriques AT, Poser GL. Acaricidal properties of the essential oil from *Hesperozygis ringens* on the cattle tick *Rhipicephalus microplus*. *Bioresour Technol* 2010; 101: 2506–2509
  - 107 Pirali-Kheirabadi K, Razzaghi-Abyaneh M, Halajian A. Acaricidal effect of *Pelargonium roseum* and *Eucalyptus globulus* essential oils against adult stage of *Rhipicephalus annulatus in vitro*. *Vet Parasitol* 2009; 162: 346–349
  - 108 Olivo CJ, Carvalho NM, Silva JHS, Vogel FF, Massariol P, Meinerz G, Agnolin C, Morel AF, Viau LV. Citronella oil on the control of cattle ticks. *Ciencia Rural* 2008; 38: 406–410
  - 109 Martins RM. *In vitro* study of the acaricidal activity of the essential oil from the citronella of Java (*Cymbopogon winterianus* Jowitt) to the tick *Boophilus microplus*. *Rev Bras Pl Med* 2006; 8: 71–78
  - 110 Martins RM, Gonzalez FHD. Use of Java (*Cymbopogon winterianus* Jowitt) (Panicoidideae) as an acaricide against the tick *Boophilus microplus* Canestrini. *Rev Bras Pl Med* 2007; 9: 1–8
  - 111 Cetin H, Cilek JE, Aydin L, Yanikoglu A. Acaricidal effects of the essential oil of *Origanum minutiflorum* against *Rhipicephalus turanicus*. *Vet Parasitol* 2009; 160: 359–361
  - 112 Pirali-Kheirabadi K, Silva JAT. *Lavandula angustifolia* essential oil as a novel and promising natural candidate for tick (*Rhipicephalus [Boophilus] annulatus*) control. *Exp Parasitol* 2010; 126: 184–186
  - 113 Landau SY, Provenza FD, Gardner DR, Pfister JA, Knoppel EL, Peterson C, Kababya D, Needham GR, Villalba JJ. Neem-tree (*Azadirachta indica* Juss.) extract as a feed additive against the American dog tick (*Dermacentor variabilis*) in sheep (*Ovis aries*). *Vet Parasitol* 2009; 165: 311–317