

Literatur zu

Postreanimationstherapie

Therapiekonzepte und Prognosestellung

Andreas Schneider • Erik Popp • Bernd W. Böttiger

1 Böttiger BW. et al.: Long term outcome after out-of-hospital cardiac arrest with physician staffed emergency medical services: the Utstein style applied to a midsized urban/suburban area. *Heart* 1999; 82: 674-679

2 Herlitz J. et al.: Resuscitation in Europe: a tale of five European regions. *Resuscitation* 1999; 41: 121-131

3 Laver Setal.: Mode of death after admission to an intensive care unit following cardiac arrest. *Intensive Care Med* 2004; 30: 2126-2128

4 European Resuscitation Council. European Resuscitation Council guidelines for resuscitation 2005. . *Resuscitation* 2005; 67: S1-S189. Deutsche Übersetzung: *Notfall Rettungsmed* 2006; 9: 6-170

5 Negovsky VA. Postresuscitation disease. *Crit Care Med* 1988; 16: 942-946

6 Padosch SA, Vogel P, Böttiger BW. Neuronale Apoptose nach zerebraler Ischämie. Grundlagen, Pathophysiologie und Interventionsmöglichkeiten. *Anaesthesist* 2001; 50: 905-920

7 Kern KBetal.: Myocardial dysfunction after resuscitation from cardiac arrest: an example of global myocardial stunning. *J Am Coll Cardiol* 1996; 28: 232-240

8 Laurent I et al.: Reversible myocardial dysfunction in survivors of out-of-hospital cardiac arrest. *J Am Coll Cardiol* 2002; 40: 2110-2116

9 Adrie C, Adib-Conquy M, Laurent I, Monchi M, Vinsonneau C, Fitting C, Fraisse F, Dinh-Xuan AT, Carli P, Spaulding CM, Dhainaut JF, Cavaillon JM. Successful cardiopulmonary resuscitation after cardiac arrest as a "sepsis-like syndrome". *Circulation* 2002; 106: 562-568

10 Böttiger BW, Motsch J, Braun V, Martin E, Kirschfink M. Marked activation of complement and leukocytes and an increase in the concentrations of soluble endothelial adhesion molecules during cardiopulmonary resuscitation and early reperfusion after cardiac arrest in humans. *Crit Care Med* 2002; 30: 2473-2480

11 Böttiger BW, Motsch J, Böhrer H, Böker T, Aulmann M, Nawroth PP, Martin E. Activation of blood coagulation after cardiac arrest is not balanced adequately by activation of endogenous fibrinolysis. *Circulation* 1995; 92: 2572-2578

12 Müllner M, Hirschl MM, Herkner H, Sterz F, Leitha T, Exner M, Binder M, Laggner AN. Creatine kinase-MB fraction and cardiac troponin T to diagnose acute myocardial infarction after cardiopulmonary resuscitation. *J Am Coll Cardiol* 1996; 28: 1220-1225

13 Spaulding CM et al.: Immediate coronary angiography in survivors of out-of-hospital cardiac arrest. *N Engl J Med* 1997; 336: 1629-1633

14 Vanbrabant P, Dhondt E, Billen P, Sabbe M. Aetiology of unsuccessful prehospital witnessed cardiac arrest of unclear origin. *Eur J Emerg Med* 2006; 13: 144-147

15 Oschatz E et al.: Cardiopulmonary resuscitation performed by bystanders does

not increase adverse effects as assessed by chest radiography. *Anesth Analg* 2001; 93: 128-133

16 Bernard SAetal.: Treatment of comatose survivors of out-of-hospital cardiac arrest with induced hypothermia. *N Engl J Med* 2002; 346: 557-563

17 Hypothermia after Cardiac Arrest Study Group.: Mild therapeutic hypothermia to improve the neurologic outcome after cardiac arrest. *N Engl J Med* 2002; 346: 549-556

18 Langhelle Aetal.: In-hospital factors associated with improved outcome after out-of-hospital cardiac arrest. A comparison between four regions in Norway. *Resuscitation* 2003; 56: 247-263

19 Sundgreen Cetal.: Autoregulation of cerebral blood flow in patients resuscitated from cardiac arrest. *Stroke* 2001; 32: 128-132

20 Sterz F. et al.: Hypertension with or without hemodilution after cardiac arrest in dogs. *Stroke* 1990; 21: 1178-1184

21 Bender Retal.: Hypertonic saline during CPR: Feasibility and safety of a new protocol of fluid management during resuscitation. *Resuscitation* 2007; 72: 74-81

22 Müllner M. et al.: Blood glucose concentration after cardiopulmonary resuscitation influences functional neurological recovery in human cardiac arrest survivors. *J Cereb Blood Flow Metab* 1997; 17: 430-436

23 Van den BergheG. et al.: Intensive insulin therapy in the critically ill patients. *N Engl J Med* 2001; 345: 1359-1367

24 Buunk G, van der Hoeven JG, Meinders AE. Cerebrovascular reactivity in comatose patients resuscitated from a cardiac arrest. *Stroke* 1997; 28: 1569-1573

25 Böttiger BW. The Thrombolysis In Cardiac Arrest (TROICA) trial. ESC Congress Reports 2006;. www.escardio.org/knowledge/congresses/CongressReports/hotlinesandctus/709011_Boettiger.htm

26 Spöhr F, Böttiger BW. Safety of thrombolysis during cardiopulmonary resuscitation. *Drug Saf* 2003; 26: 367-379

27 Ingvar M. Cerebral blood flow and metabolic rate during seizures. Relationship to epileptic brain damage. *Ann N Y Acad Sci* 1986; 462: 194-206

28 Bauer J. et al.. Status epilepticus im Erwachsenenalter. In: Diener HC, Putzki N, Berlit P (Hrsg.). Leitlinien für Diagnostik und Therapie in der Neurologie. 3. Aufl. Stuttgart: Thieme 2005, p. 20-27

29 Hachimi-Idrissi S. et al. Mild hypothermia induced by a helmet device: a clinical feasibility study. *Resuscitation* 2001; 51: 275-281

30 Coimbra C, Boris-Möller F, Drake M, Wieloch T. Diminished neuronal damage in the rat brain by late treatment with the antipyretic drug dipyron or cooling following cerebral ischemia. *Acta Neuropathol* 1996; 92: 447-453

31 Kuboyama K, Safar P, Radovsky A, Tisherman SA, Stezoski SW, Alexander H. Delay in cooling negates the beneficial effect of mild resuscitative cerebral hypothermia after cardiac arrest in dogs: a prospective, randomized study. *Crit Care Med* 1993; 21: 1348-1358

32 Bernard S, Buist M, Monteiro O, Smith K. Induced hypothermia using large volume, ice-cold intravenous fluid in comatose survivors of out-of-hospital cardiac arrest: a preliminary report. *Resuscitation* 2003; 56: 9-13

33 Virkkunen I, Yli-Hankala A, Silfvast T. Induction of therapeutic hypothermia after cardiac arrest in prehospital patients using ice-cold Ringer's solution: a pilot study. *Resuscitation* 2004; 62: 299-302

34 Schwab S. et al.. Moderate hypothermia and brain temperature in patients with severe middle cerebral artery infarction. *Acta Neurochir Suppl* 1998; 71: 131-134

35 Schwab S. et al.. Feasibility and safety of moderate hypothermia after massive hemispheric infarction. *Stroke* 2001; 32: 2033-2035

36 Sessler DI. Mild perioperative hypothermia. *N Engl J Med* 1997; 336: 1730-1737

37 Kranke P. et al. Pharmacological treatment of postoperative shivering: a quantitative systematic review of randomized controlled trials. *Anesth Analg* 2002; 94: 453-460

38 Booth CM. et al. Is this patient dead, vegetative, or severely neurologically impaired? Assessing outcome for comatose survivors of cardiac arrest. *JAMA* 2004; 291: 870-879

39 Zandbergen EGJ. et al.. Systematic review of early prediction of poor outcome in anoxic-ischaemic coma. *Lancet* 1998; 352: 1808-1812

40 Böttiger BW. et al. Astroglial protein S-100 is an early and sensitive marker of hypoxic brain damage and outcome after cardiac arrest in humans. *Circulation* 2001; 103: 2694-2698

41 Zandbergen EGJ, de HaanRJ, Hijdra A. Systematic review of prediction of poor outcome in anoxic-ischaemic coma with biochemical markers of brain damage. *Intensive Care Med* 2001; 27: 1661-1667

42 Torbey MT. et al. Quantitative analysis of the loss of distinction between gray and white matter in comatose patients after cardiac arrest. *Stroke* 2000; 31: 2163-2167

43 Dae MW, Gao DW, Sessler DI, Chair K, Stillson CA. Effect of endovascular cooling on myocardial temperature, infarct size, and cardiac output in human-sized pigs. *Am J Physiol Heart Circ Physiol* 2002; 282:

44 Popp E, Sterz F, Böttiger BW. Therapeutische milde Hypothermie nach Herz-Kreislauf-Stillstand. *Anaesthesist* 2005; 54: 96-106