

Literatur

- 1 <http://www.bundesaerztekammer.de/page.asp?his=3.71.11025.11634.11638>
- 2 <http://www.dso.de>
- 3 Bundesärztekammer, Dezernat VI. Richtlinien zur Organtransplantation gemäß § 16 TPG. Dtsch Arztebl2006; 103: A3282 – A3290
- 4 Schemmer P, Nickkholgh A, Hinz U et al. Extended donor criteria have no negative impact on early outcome after liver transplantation: a single-center multivariate analysis. *Transplant Proc*2007; 39: 529 – 534
- 5 Schemmer P, Nickkholgh A, Gerling T et al. Rescue allocation for liver transplantation within Eurotransplant: The Heidelberg experience. *Clin Transplant*2009; 23: 2142 – 48
- 6 Proneth A, Schnitzbauer AA, Zeman F et al. Extended pancreas donor program - the EXPAND study rationale and study protocol. *Transplant Res*2013; 2: 12
- 7 Gasser M, Waaga-Gasser AM, Grimm MW et al. Selectin blockade plus therapy with low-dose sirolimus and cyclosporin a prevent brain death-induced renal allograft dysfunction. *Am J Transplant*2005; 5: 662 – 670
- 8 Schemmer P, Schoonhoven R, Swenberg JA et al. Gentle in situ organ manipulation during harvest decreases survival after rat liver transplantation: The role of Kupffer cells. *Transplantation*1998; 65: 1015 – 1020
- 9 Schindler G, Kincius M, Liang R et al. Fundamental efforts towards the development of a therapeutic cocktail with a manifold ameliorative effect on hepatic ischemia/reperfusion injury. *Microcirculation*2009; 16: 593 – 602
- 10 Liu Q, Bruns H, Schultze D et al. HTK-N, a modified HTK solution, decreases preservation injury in a model of microsteatotic rat liver transplantation. *Langenbecks Arch Surg*2012; 397: 1323 – 1331
- 11 O'Callaghan JM, Morgan RD, Knight SR et al. Systematic review and meta-analysis of hypothermic machine perfusion versus static cold storage of kidney allografts on transplant outcomes. *Br J Surg*2013; 100: 991 – 1001
- 12 Minor T, Paul A. Hypothermic reconditioning in organ transplantation. *Curr Opin Organ Transplant*2013; 18: 161 – 167
- 13 Guan X, Dei-Anane G, Bruns H et al. Danshen protects kidney grafts from ischemia/reperfusion injury after experimental transplantation. *Transpl Int*2009; 22: 232 – 241
- 14 Schemmer P, Mehrabi A, Friess H et al. Living related liver transplantation: the ultimate technique to expand the donor pool? *Transplantation*2005; 80: S138 – S141
- 15 Weitz J, Koch M, Mehrabi A et al. Living donor kidney transplantation: risks of the donor - benefits of the recipient. *Clin Transplant*2006; 20: 1713 – 16
- 16 Eurotransplant Annual Report.2012: <http://www.eurotransplant.org/cms/mediaobject.php?file=AR2012.pdf>
- 17 Collaborative Transplant Study. [CTS-K-21181, K-TONS05E-0312] <http://www.ctstransplant.org/public/graphics.shtml>
- 18 Mange KC, Weir MR. Preemptive renal transplantation: why not? *Am J Transplant*2003; 3: 1336 – 1340
- 19 de Groot IB, Stiggelbout AM, van der Boog PJ et al. PARTNER-study group. Reduced quality of life in living kidney donors: association with fatigue, societal participation and pre-donation variables. *Transpl Int*2012; 25: 967 – 975
- 20 Weismüller TJ, Fikatas P, Schmidt J et al. Multicentric evaluation of model for end-stage liver disease-based allocation and survival after liver transplantation in Germany – limitations of the 'sickest first'-concept. *Transpl Int*2011; 24: 91 – 99
- 21 Weismüller TJ, Prokein J, Becker T et al. Prediction of survival after liver transplantation by pre-transplant parameters. *Scand J Gastroenterol*2008; 43: 736 – 746