

**Literatur**

- Dekker J, Van Henricus L, Hendriksen M. What Is the Best Sequential Treatment Strategy in the Treatment of Depression? Adding Pharmacotherapy to Psychotherapy or Vice Versa? *Psychother Psychosom* 2013; 82: 89–98
- Forand N, deRubeis R, Amsterdam J. Combining Medication and Psychotherapy in the Treatment of Major Mental Disorders. In: Lambert MJ, ed. *Bergin and Garfield's Handbook of Psychotherapy and Behavior Change*. 6th Ed. Hoboken, New Jersey: John Wiley; 2013: 735–774
- Grosse Holtforth M, Hayes AM, Sutter M et al. Fostering Cognitive-Emotional Processing in the Treatment of Depression: A Preliminary Investigation in Exposure-Based Cognitive Therapy. *Psychother Psychosom* 2012; 81: 259–260
- Härter M, Klesse C, Bermejo I et al. Unipolare Depression: Empfehlungen zur Diagnostik und Therapie aus der aktuellen S3- und Nationalen VersorgungsLeitlinie „Unipolare Depression“. *Deutsches Ärzteblatt International* 2010; 107: 700–708
- Hegerl U. Sollten leichte Depressionen ausschließlich psychotherapeutisch behandelt werden? *Contra. Nervenarzt* 2013; 84: 384–385
- Hofmann SG, Asnaani A, Vonk IJ et al. The Efficacy of Cognitive Behavioral Therapy: A Review of Meta-Analyses. *Cognit Ther Res* 2012; 36: 427–440
- Huber D, Zimmermann J, Henrich G et al. Comparison of Cognitive-Behaviour Therapy with Psychoanalytic and Psychodynamic Therapy for Depressed Patients – A Three-Year Follow-Up Study. *Z Psychosom Med Psychother* 2012; 58: 299–316
- Joutsenniemi K, Laaksonen MA, Knekt P et al. Prediction of the Outcome of Short- and Long-Term Psychotherapy Based on Socio-Demographic Factors. *J Affect Disord* 2012; 141: 331–342
- Kirsch I, Deacon BJ, Huedo-Medina TB et al. Initial Severity and Antidepressant Benefits: A Meta-Analysis of Data Submitted to the Food and Drug Administration. *PLoS Med* 2008; 5: e45
- Knekt P, Lindfors O, Sares-Jäske L et al. Randomized Trial on the Effectiveness of Long- and Short-Term Psychotherapy on Psychiatric Symptoms and Working Ability during a 5-Year Follow-Up. *Nord J Psychiatry* 2013; 69: 59–68
- Laaksonen MA, Knekt P, Lindfors O. Psychological Predictors of the Recovery from Mood or Anxiety Disorders in Short-term and Long-term Psychotherapy During a 3-Year Follow-up. *Psychiatry Res* 2012; DOI: 10.1016/j.psychres.2012.09.053
- Lambert MJ. The Efficacy and Effectiveness of Psychotherapy. In: Lambert MJ, ed. *Bergin and Garfield's Handbook of Psychotherapy and Behavior Change*. 6th Ed. Hoboken, New Jersey: John Wiley; 2013: 169–218
- Schauenburg H, Bschor T. Sollten leichte Depressionen ausschließlich psychotherapeutisch behandelt werden? *Pro. Nervenarzt* 2013; 84: 386–387
- Schramm E, Schweiger U, Hohagen F et al. Psychotherapie für chronische Depression. *Cognitive Behavioral Analysis System of Psychotherapy (CBASP)* von James P. McCullough. Deutsche Übersetzung und Bearbeitung. München: Elsevier; 2006
- Shedler J. The Efficacy of Psychodynamic Psychotherapy. *Am Psychol* 2010; 99: 98–109
- Shimokova K, Lambert MJ, Smart DW. Enhancing Outcome of Patients at Risk for Failure: Meta-Analytic and Mega-Analytic Review of a Psychotherapy Quality Assurance System. *J Consult Clin Psychol* 2010; 78: 298–311