

Supporting Information

Phillyrin, a Natural Lignan, Attenuates Tumor Necrosis Factor α -Mediated Insulin Resistance and Lipolytic Acceleration in 3T3-L1 Adipocytes

Poren Kong¹, Linlin Zhang², Yuyu Guo¹, Yingli Lu¹, Dongping Lin¹

Affiliations

¹Department of Endocrine and Metabolic Diseases, Laboratory of Endocrine and Metabolic Diseases, Ninth People's Hospital, Shanghai Jiaotong University, School of Medicine, Shanghai, China

²Yantai Traditional Chinese Medicine Hospital, Affiliated to Shandong University of Traditional Chinese Medicine, Yantai, China

Correspondence

Dr. Poren Kong (Bo-ren Jiang in Chinese); Dr. Dongping Lin

Department of Endocrine and Metabolic Diseases

Laboratory of Endocrine and Metabolic Diseases, Ninth People's Hospital

Shanghai Jiaotong University, School of Medicine

No. 639, Zhizhaoju Road

Shanghai 200211 Shanghai

China

Phone: +86 21 2327 1699 5139

Fax: +86 21 2327 1699

porenkong@gmail.com; lindongping2005@126.com

Fig. 1S Relative quantities of total Glut4 (A) and Glut4 in PM fraction (B) from **Fig. 2E**. The data are shown as mean \pm SE of three independent experiments.

