


Available online at www.sciencedirect.com

SciVerse ScienceDirect

journal homepage: www.elsevier.com/locate/ijnt

Letter to the Editor

Role of autoantibodies in the pathophysiology of hypopituitarism following traumatic brain injury

We read with interest the review 'Endocrine manifestations of traumatic brain injury' written by Agrawal et al.¹ The authors have broadly described the causes of post-traumatic brain injury (TBI) hypopituitarism to be ischemia-hypoxia, direct mechanical injury, compression injury and vascular injury. It is worthwhile to mention here that recent studies have shown the presence of anti-pituitary antibodies (APAs) and anti-hypothalamic antibodies (AHAs) following TBI as an important potential cause of hypopituitarism. Preliminary studies on animals have shown the presence of APAs and AHAs, similar to those seen in Sheehan's syndrome of postpartum pituitary necrosis.^{2,3} Up to 44% patients with TBI showed the presence of APAs in a study by Tanriverdi et al.³ These APAs may have distinct selective bias toward the growth hormone (GH) releasing hormone secreting cells as compared to the vasopressin secreting cells, as suggested by Tanriverdi's study, which showed that all AHA-positive cases were accompanied by at least some GH deficiency but no diabetes insipidus.⁴ Damage to the blood-brain barrier allowing escape of brain proteins into the circulation and thus evoking an immunological response⁵ or, inflammation triggering a vascular cascade causing histopathological changes may be responsible.³ Spontaneous disappearance of these APAs from the circulation, allowing functional recovery of the pituitary may be the cause of transient hypopituitarism seen in a number of TBI patients.⁴ De Bellis et al⁶ suggest that if detectable in high titer (>1.8) these antibodies may be used as markers of hypopituitarism. Research on the mechanism of formation of these autoantibodies after TBI may have promising rewards in preventing post-TBI hypopituitarism-induced morbidity and mortality in the future.

2. De Bellis A, Kelestimur F, Sinisi AA, et al. Anti-hypothalamus and anti-pituitary antibodies may contribute to perpetuate the hypopituitarism in patients with Sheehan's syndrome. *Eur J Endocrinol.* 2008;158(2):147–152.
3. Tanriverdi F, De BA, Bizzarro A, et al. Antipituitary antibodies after traumatic brain injury: is head trauma-induced pituitary dysfunction associated with autoimmunity? *Eur J Endocrinol.* 2008;159(1):7–13. <http://dx.doi.org/10.1530/EJE-08-0050>. pii: EJE-08-0050.
4. Tanriverdi F, De BA, Battaglia M, et al. Investigation of antihypothalamus and antipituitary antibodies in amateur boxers: is chronic repetitive head trauma-induced pituitary dysfunction associated with autoimmunity? *Eur J Endocrinol.* 2010;162(5):861–867. <http://dx.doi.org/10.1530/EJE-09-1024>. pii: EJE-09-1024.
5. Papa L, Lewis LM, Falk JL, et al. Elevated levels of serum glial fibrillary acidic protein breakdown products in mild and moderate traumatic brain injury are associated with intracranial lesions and neurosurgical intervention. *Ann Emerg Med.* 2012;59(6):471–483. <http://dx.doi.org/10.1016/j.annemergmed.2011.08.021>. pii: S0196-0644(11)01545-9.
6. De Bellis A, Bizzarro A, Conte M, et al. Antipituitary antibodies in adults with apparently idiopathic growth hormone deficiency and in adults with autoimmune endocrine diseases. *J Clin Endocrinol Metab.* 2003;88(2):650–654.

Viraat Harsh

Post-Doctoral Research Associate, Department of Neurosurgery,
Baylor College of Medicine, Houston, TX 77030, USA

Anil Kumar*

Professor and Head, Department of Neurosurgery,
Rajendra Institute of Medical Sciences, Ranchi,
Jharkhand 834009, India*Corresponding author. Tel.: +91 943 111 5510.
E-mail address: dr.anilkumarranchi@gmail.com

12 February 2013

Available online 9 April 2013

REFERENCES

1. Agrawal A, Reddy PA, Prasad NR. Endocrine manifestations of traumatic brain injury. *The Indian J Neurotrauma.* 2012;9(2):123–128. <http://dx.doi.org/10.1016/j.ijnt.2012.11.005>.

0973-0508/\$ – see front matter
Copyright © 2013, Neurotrauma Society of India. All rights reserved.
<http://dx.doi.org/10.1016/j.ijnt.2013.04.001>