


## Endoscopic ultrasound-guided hepaticoduodenostomy for an obstructed anterior hepatic branch in a patient with Billroth-II reconstruction

A 70-year-old man who had previously undergone distal gastrectomy with Billroth-II reconstruction visited our hospital complaining of a repetitive fever. Contrast-enhanced computed tomography (CT) revealed a 25-mm liver abscess in the right anterior segment, communicating with the anterior hepatic duct, which was dilated due to stenosis at the anomalous confluence with the middle of the cystic duct; this obstruction was believed to be the cause of the fever (► **Fig. 1**). Conservative treatment with antimicrobial agents was not sufficiently effective. After obtaining written informed consent from the patient, endoscopic ultrasound (EUS)-guided biliary drainage was performed to resolve the structural problem (► **Fig. 2**, ► **Video 1**) after considering other interventions, including transpapillary stenting through


the narrow, winding cystic duct and highly stenotic confluence. Using a forward-viewing echoendoscope (TGFUC260J; Olympus Co., Tokyo, Japan) inserted into the blind end of the duodenum, the dilated anterior hepatic duct was punctured using a 19-gauge needle (Expect; Boston Scientific Japan K. K., Tokyo, Japan) under EUS guidance. After dilation of the punctured tract using a 7-Fr bougie catheter, a 4-mm balloon catheter, and a 6-Fr cauterly dilator, a fully covered metallic stent (Bonastent, M-Intraductal, 10 mm × 7 cm; Medico's Hirata Inc., Tokyo, Japan) was deployed at the puncture site. No procedure-related adverse events were observed.

CT showed diminution of the drained duct with air influx 7 days after the procedure (► **Fig. 3**), and disappearance of the abscess 3 months later (► **Fig. 4**). No symptoms indicating recurrence were experienced in the subsequent 10 months.


Previous reports have described the use of the EUS-guided biliary drainage technique for hepatic abscess drainage performed using a forward-viewing scope in patients with surgically altered anatomies [1–3]. The present case was the first application of these techniques for a patient with both challenging conditions and with reasons that made other treatment options difficult.




► **Fig. 1** Computed tomography images. Axial (a) and coronal (b) views showing a 25-mm liver abscess in the right anterior segment (arrow).




► **Fig. 2** Endoscopic ultrasound (EUS)-guided biliary drainage. a The hepatic bile duct, which was filled with biliary sludge, was punctured using a 19-gauge needle from the duodenum under EUS guidance. b Injected contrast medium revealed the dilated hepatic duct, which was obstructed at the confluence with the middle of the cystic duct (arrow). c, d A fully covered self-expandable metallic stent was deployed at the puncture site.



► **Fig. 3** Computed tomography images. Axial (a) and coronal (b) views showing diminution of the drained duct with air influx 7 days after the procedure.



► **Fig. 4** Computed tomography images. Axial (a) and coronal (b) views showing the disappearance of the abscess 3 months after the procedure.



► **Video 1** Endoscopic ultrasound-guided biliary drainage procedures for the right anterior hepatic duct using a forward-viewing echoendoscope.



## Corresponding author

### Yujiro Kawakami, MD

Department of Gastroenterology, Sendai City Medical Center, Tsurugaya 5-22-1, Miyagino-ku, Sendai, Japan  
Fax: +81-22-2529431  
yujiro.kawakami@gmail.com

## References

- [1] Kanno Y, Koshita S, Ogawa T et al. EUS-guided biliary drainage for unresectable malignant biliary obstruction: 10-year experience of 99 cases at a single center. *J Gastrointest Cancer* 2018. doi:10.1007/s12029-018-0096-1
- [2] Yamamoto K, Itoi T, Tsuchiya T et al. EUS-guided drainage of hepatic abscess in the right side of the liver of a patient with Chlaiditi syndrome. *VideoGIE* 2017; 2: 299–300
- [3] Kida M, Yamauchi H, Okuwaki K et al. Endoscopic ultrasound-guided choledochojunostomy with a forward-viewing echoendoscope for severe benign bilioenteric stricture in a patient with Child's resection. *Endoscopy* 2015; 47: E303–E304

## Bibliography

DOI <https://doi.org/10.1055/a-0885-9348>  
Published online: 30.4.2019  
*Endoscopy* 2019; 51: E221–E222  
© Georg Thieme Verlag KG  
Stuttgart · New York  
ISSN 0013-726X

## ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



*Endoscopy E-Videos* is a free access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at

<https://mc.manuscriptcentral.com/e-videos>

Endoscopy\_UCTN\_Code\_TTT\_1AR\_2AG

## The authors

Yujiro Kawakami, Yoshihide Kanno, Shinsuke Koshita, Takahisa Ogawa, Hiroaki Kusunose, Toji Murabayashi, Kei Ito

Department of Gastroenterology, Sendai City Medical Center, Sendai, Japan

## Competing interests

None