


Concise Total Synthesis of Curvulone B

Shivalal Banoth^{a,b}Utkal Mani Choudhury^{a,b}Kanakaraju Marumudi^cAjit C. Kunwar^cDebendra K. Mohapatra^{*a,b}

^a Department of Organic Synthesis and Process Chemistry, CSIR-Indian Institute of Chemical Technology, Hyderabad 500 007, India

^b Academy of Scientific and Innovative Research (AcSIR), Ghaziabad 201 002, India

^c Centre for NMR and Structural Chemistry, CSIR-Indian Institute of Chemical Technology, Hyderabad 500 007, India
mohapatra@iict.res.in



Received: 08.10.2020

Accepted after revision: 26.10.2020

Published online: 26.10.2020

DOI: 10.1055/a-1297-6838; Art ID: St-2020-d0546-1



Abstract A concise and convergent stereoselective synthesis of curvulone B is described. The synthesis utilized a tandem isomerization followed by C–O and C–C bond-forming reactions following Mukaiyama-type aldol conditions for the construction of the *trans*-2,6-disubstituted dihydropyran ring system as the key steps. Other important features of this synthesis are a cross-metathesis, epimerization, and Friedel–Crafts acylation.

Key words curvulone B, natural products, Mukaiyama–aldol reaction, Friedel–Crafts acylation reaction

Marine fungi have been long recognized as a rich source of novel secondary metabolites with such biological properties as antitumor, phytotoxic, or antifungal activities, as well as cytotoxicity against human cancer cell lines.¹ In connection with the search for biologically active metabolites from fungi, Krohn and Kurtán and their co-workers isolated two new curvularin-type metabolites, curvulone A (**1**) and curvulone B (**2**; Figure 1) from a *Curvularia* sp. obtained from the marine alga *Gracilaria folifera*.¹ Curvulone B (**2**) features a 2,6-disubstituted *cis*-tetrahydropyran ring, and displays antitumor, antifungal, and cytotoxic activities.²

The structure of curvulone B was determined by 2D NMR spectroscopy, and its absolute configuration was de-

duced by comparison of the experimental electronic circular dichroism spectra in acetonitrile with the Boltzman-averaged spectrum.³ Total syntheses of curvulone B (**2**) have been reported by the groups of Takahashi,² Bates,⁴ and, more recently, He.⁵ None of these syntheses involved fewer than ten linear steps, and all employed an intramolecular oxa-Michael addition for the formation of THP ring.

We recently reported the synthesis of 2,6-*trans*-disubstituted tetrahydropyrans with a keto functionality by means of a Mukaiyama-type aldol reactions of 1-phenyl-1-(trimethylsiloxy)ethylene with six-membered cyclic hemiacetals in the presence of iodine.⁶ As a further application of this Mukaiyama-type aldol reaction, and as a part of our ongoing research on the total synthesis of biologically active natural products containing pyran rings,⁷ we report an efficient and convergent synthesis of curvulone B in seven steps.

Our retrosynthetic analysis of curvulone B is outlined in Scheme 1. It was envisaged that curvulone B might be prepared by a Friedel–Crafts acylation reaction of aromatic ester **3** and the acid fragment **4** (Scheme 1). We planned to obtain intermediate **4** from the *trans*-2,6-disubstituted dihydropyran **5** that, in turn, would be accessed from a δ -hydroxy α,β -unsaturated aldehyde through tandem isomeri-






Figure 1 Structures of curvulone A (**1**) and curvulone B (**2**)



Scheme 1 Retrosynthetic analysis of curvulone B (**2**)


zation followed by C–O and C–C bond-forming reactions of a silyl enol ether under Mukaiyama-type aldol reaction conditions. The δ -hydroxy α,β -unsaturated aldehyde would be obtained from the commercially available chiral homoallylic alcohol **6**.

The synthesis of the key intermediate **5** began with the commercially available homoallylic alcohol **6** and acrolein, which, on treatment with the Hoveyda–Grubbs catalyst (10 mol%) in CH_2Cl_2 gave the cross-metathesis⁸ product, the δ -hydroxy α,β -unsaturated aldehyde **7**, in 87% yield (Scheme 2). Tandem isomerization followed by a C–O and C–C bond-formation protocol under Mukaiyama-type conditions was performed by treating **7** with trimethyl(vinyloxy)silane in the presence of a catalytic amount of molecular iodine in anhydrous CH_2Cl_2 at room temperature to furnish the *trans*-2,6-disubstituted-3,4-dihydropyran **5** as the sole product in 81% yield.^{6,9,10}




Scheme 2 Synthesis of compound **5**

The next task was to reduce the internal double bond and then to perform the isomerization reaction. Accordingly, the double bond in compound **5** was reduced in the presence of a catalytic amount of Adams's catalyst under hydrogen in anhydrous ethyl acetate to furnish compound **8** in excellent yield (Scheme 3). Epimerization was performed by a retro-oxa-Michael/oxa-Michael reaction with potassium *tert*-butoxide in THF at 0 °C; this reaction was highly stereoselective, favoring the desired C- β -epimer **9**, which was obtained in 92% yield.¹¹ Acid **4**, a key fragment for the Friedel–Crafts acylation strategy, was then synthesized in 86% yield from *cis*-pyran aldehyde **9** by Pinnick oxidation¹² with NaClO_2 , NaH_2PO_4 , *t*-BuOH– H_2O (1:1), and 2-methylbut-2-ene.




Scheme 3 Synthesis of compound **4**

The aromatic coupling fragment **3** was synthesized by methylation of commercially available methyl 2-(3,5-dihydroxyphenyl)acetate (**10**) with potassium carbonate and dimethyl sulfate in acetone to afford methyl 2-(3,5-dimethoxyphenyl)acetate (**3**) in 95% yield (Scheme 4).¹³



Scheme 4 Synthesis of fragment **3**

With *cis*-pyran acid **4** and methyl 2-(3,5-dimethoxyphenyl)acetate (**3**) in hand, our next objective was to combine both fragments by using the key Friedel–Crafts acylation reaction. Accordingly, treatment of *cis*-pyran acid **4** with methyl 2-(3,5-dimethoxyphenyl)acetate (**3**) in TFA/TFAA, afforded the desired ketone **11** in 93% yield (Scheme 5).¹⁴



Scheme 5 Completion of the total synthesis of curvulone B (**2**)

The structure of compound **11** was confirmed by extensive NMR experiments, including DQF-COSY, TOCSY, NOESY, HSQC, and HMBC experiments. The distinctive AB spin system of double doublets at $\delta = 2.97$ and 3.04 ppm, due to 10-H and 10-H' displaying a HMBC correlation with the carbonyl carbon ($\delta = 204.3$ ppm), was used to initiate the assignments. The DQF-COSY experiment helped us to assign the protons from 11-H to 15-H and the 16- CH_3 protons. The 2- CH_2 protons appear as a broad singlet at $\delta = 3.60$ ppm. The nOe correlations 11-H/15-H, 11-H/13-H, 13-H/15-H, and 12-H'/14-H' strongly supported the *syn* orientation of the 11-H and 15-H protons, as well as the $^{14}\text{C}_{11}$ chair conformation of the six-membered ring. Furthermore, nOe correlations between 10-H/2- CH_2 , 2- CH_2 /4-H, and 7'- OCH_3 /10-H provided strong evidence that the pyran ring occupies a position *ortho* to methyl ester of the benzene ring, providing firm support for the proposed structure of **11** (Figure 2).

Finally, demethylation of the methoxy group of **11** was successfully achieved under Maier's conditions (AlI_3 , TBAI, phloroglucinol)¹⁵ in benzene at 0 °C to furnish curvulone B (**2**) in 91% yield.¹⁶ The spectroscopic and analytical data for synthetic compound **2** were in good agreement with those reported for the natural product.¹




Figure 2 Energy-minimized structure of **11**, along with key nOe correlations (double-headed arrows)

In summary, a concise and stereoselective synthesis of the curvulone B (**2**) was achieved in seven steps and a 46% overall yield by using iodine-catalyzed tandem isomerization followed by C–O and C–C bond-formation through a Mukaiyama-type aldol reaction for the construction of the *trans*-2,6-disubstituted dihydropyran ring system as the key step. The other important reactions involved in the current synthetic approach were cross-metathesis, epimerization, and Friedel–Crafts acylation reactions.

Funding Information

The authors gratefully acknowledge financial support received from the Science and Engineering Research Board (SERB), a statutory body of the Department of Science & Technology (DST), New Delhi, Government of India, through Grant No. EMR/2017/002298.

Acknowledgment

We are grateful to the Director, CSIR-IICT for his kind support and for providing research facilities (Manuscript communication number IICT/Pubs/2020/305). S.B., U.M.C., and K.M. thank UGC, New Delhi, India, for financial assistance in the form of fellowships.

Supporting Information

Supporting information for this article is available online at <https://doi.org/10.1055/a-1297-6838>.

References and Notes

- (1) Dai, J.; Krohn, K.; Flörke, U.; Pescitelli, G.; Kerti, G.; Papp, T.; Kövér, K. E.; Bényei, A. C.; Draeger, S.; Schulz, B.; Kurtán, T. *Eur. J. Org. Chem.* **2010**, 6928; and references therein.
- (2) Takahashi, S.; Akita, Y.; Nakamura, T.; Koshino, H. *Tetrahedron: Asymmetry* **2012**, *23*, 952; and references therein.
- (3) Guo, Y.-W.; Kurtán, T.; Krohn, K.; Pescitelli, G.; Zhang, W. *Chirality* **2009**, *21*, 561.
- (4) Bates, R. W.; Wang, K.; Zhou, G.; Kang, D. Z. *Synlett* **2015**, *26*, 751.
- (5) Allu, S. R.; Banne, S.; Jiang, J.; Qi, N.; Guo, J.; He, Y. J. *Org. Chem.* **2019**, *84*, 7227.
- (6) Bharath, Y.; Choudhury, U. M.; Sadhana, N.; Mohapatra, D. K. *Org. Biomol. Chem.* **2019**, *17*, 9169.

- (7) (a) Mallampudi, N. A.; Srinivas, B.; Reddy, J. G.; Mohapatra, D. K. *Org. Lett.* **2019**, *21*, 5952. (b) Srinivas, B.; Reddy, D. S.; Mallampudi, N. A.; Mohapatra, D. K. *Org. Lett.* **2018**, *20*, 6910. (c) Reddy, G. S.; Padhi, B.; Bharath, Y.; Mohapatra, D. K. *Org. Lett.* **2017**, *19*, 6506. (d) Maity, S.; Kanikarapu, S.; Marumudi, K.; Kunwar, A. C.; Yadav, J. S.; Mohapatra, D. K. *J. Org. Chem.* **2017**, *82*, 4561. (e) Banoth, S.; Maity, S.; Kumar, S. R.; Yadav, J. S.; Mohapatra, D. K. *Eur. J. Org. Chem.* **2016**, 2300. (f) Thirupathi, B.; Mohapatra, D. K. *Org. Biomol. Chem.* **2016**, *14*, 6212. (g) Padhi, B.; Reddy, D. S.; Mohapatra, D. K. *Eur. J. Org. Chem.* **2015**, 542. (h) Reddy, D. S.; Padhi, B.; Mohapatra, D. K. *J. Org. Chem.* **2015**, *80*, 1365.
- (8) (a) Garber, S. B.; Kingsbury, J. S.; Gray, B. L.; Hoveyda, A. H. *J. Am. Chem. Soc.* **2000**, *122*, 8168. (b) Dinh, M.-T.; Bouzbouz, S.; Peglion, J.-L.; Cossy, J. *Tetrahedron* **2008**, *64*, 5703.
- (9) Mohapatra, D. K.; Das, P. P.; Pattanayak, M. R.; Yadav, J. S. *Chem. Eur. J.* **2010**, *16*, 2072.
- (10) [(**2R,6R**)-6-Methyl-5,6-dihydro-2H-pyran-2-yl]acetaldehyde (**5**)
I₂ (0.89 g, 3.51 mmol) was added to a stirred solution of aldehyde **7** (2.0 g, 17.54 mmol) and trimethyl(vinyloxy)silane (3.85 mL, 26.31 mmol) in anhyd CH₂Cl₂ (50 mL) at 0 °C, and the mixture was allowed to warm to rt. When the reaction was complete (TLC), it was quenched with sat. aq Na₂S₂O₃ (30 mL), and the mixture was extracted with CH₂Cl₂ (2 × 50 mL). The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated under reduced pressure. The crude product was purified by column chromatography [silica gel, EtOAc–hexane (1:5)] to give a pale-yellow liquid; yield: 1.99 g (81%); [α]_D²⁰ –68.0 (c = 0.85, CHCl₃).
IR (neat): 3033, 2971, 2928, 1721, 1636, 1373, 1187, 1137, 1090 cm⁻¹. ¹H NMR (300 MHz, CDCl₃): δ = 9.81 (s, 1 H), 5.87 (m, 1 H), 5.70 (d, J = 11.7 Hz, 1 H), 4.78 (m, 1 H), 3.83 (m, 1 H), 2.76 (ddd, J = 16.2, 8.8, 3.0 Hz, 1 H), 2.55 (dd, J = 16.2, 4.7 Hz, 1 H), 2.06–1.91 (m, 2 H), 1.21 (d, J = 6.2 Hz, 3 H). ¹³C NMR (75 MHz, CDCl₃): δ = 201.0, 127.5, 125.2, 67.6, 64.1, 47.9, 31.5, 20.6. HRMS (ESI): m/z [M + Na]⁺ calcd for C₈H₁₂NaO₂: 163.0728; found: 163.0724.
- (11) Yakambram, P.; Puranik, V. G.; Gurjar, M. K. *Tetrahedron Lett.* **2006**, *47*, 3781.
- (12) (a) Ballakrishna, S. B.; Childers, W. E. Jr.; Pinnick, H. W. *Tetrahedron* **1981**, *37*, 2091. (b) Dalcanale, E.; Montanari, F. *J. Org. Chem.* **1986**, *51*, 567.
- (13) Liang, Q.; Zhang, J.; Quan, W.; Sun, Y.; She, X.; Pan, X. *J. Org. Chem.* **2007**, *72*, 2694.
- (14) Tauber, J.; Rudolph, K.; Rohr, M.; Erkel, G.; Opatz, T. *Eur. J. Org. Chem.* **2015**, 3587.
- (15) Rink, C.; Sasse, F.; Zubriené, A.; Matulis, D.; Maier, M. E. *Chem. Eur. J.* **2010**, *16*, 14469.
- (16) **Curvulone B (2)**
A suspension of Al powder (192 mg, 7.37 mmol) in anhyd benzene (5 mL) was treated with I₂ (0.7 g, 2.74 mmol) under argon, and the violet mixture was stirred under reflux for 30 min until the mixture became colorless. The mixture was then cooled to 0 °C, and TBAI (12.7 mg, 0.034 mmol) and phloroglucinol (108 mg, 0.85 mmol) were added, followed by a solution of **11** (60 mg, 0.17 mmol) in anhyd benzene (2 mL) added in one portion. The resulting green–brown suspension was stirred for 30 min at 0 °C. When the reaction was complete (TLC), it was quenched with sat. aq aqueous Na₂S₂O₃ (10 mL), and the mixture was diluted with EtOAc (15 mL). The organic layer was separated, and the aqueous layer was extracted with EtOAc (3 × 15 mL). The combined organic extracts were washed with brine (25 mL), filtered, dried (Na₂SO₄), and concentrated under reduced pressure. The crude product was purified by column

chromatography [silica gel, EtOAc-hexane (1:1)] to give a colorless liquid; yield: 50 mg (91%); $[\alpha]_D^{20}$ -18.2 ($c = 0.2$, EtOH).

IR (neat): 3410, 2928, 1713, 1613, 1451, 1334, 1166 cm^{-1} . ^1H NMR (400 MHz, CDCl_3): $\delta = 9.79$ (s, 1 H), 6.28 (d, $J = 2.3$ Hz, 1 H), 6.22 (d, $J = 2.3$ Hz, 1 H), 6.08 (br s, 1 H), 4.13 (brtt, $J = 10.5, 2.3$ Hz, 1 H), 3.92 (d, $J = 16.5$ Hz, 1 H), 3.70 (s, 3 H), 3.57 (m, 1 H),

3.51 (d, $J = 16.6$ Hz, 1 H), 3.30 (dd, $J = 14.3, 10.1$ Hz, 1 H), 2.56 (dd, $J = 14.3, 3.1$ Hz, 1 H), 1.85 (m, 1 H), 1.65–1.50 (m, 3 H), 1.42 (m, 1 H), 1.25 (m, 1 H), 1.17 (d, $J = 6.2$ Hz, 3 H). ^{13}C NMR (100 MHz, CDCl_3): $\delta = 204.2, 172.4, 159.4, 135.7, 120.7, 111.7, 104.0, 77.8, 74.8, 52.0, 49.0, 39.7, 32.6, 30.7, 23.1, 21.5$. HRMS (ESI): m/z $[\text{M} + \text{H}]^+$ calcd for $\text{C}_{17}\text{H}_{23}\text{O}_6$: 323.1489; found: 323.1494.