



Forward-viewing echoendoscope plus gel immersion technique in a patient with Billroth II reconstruction

▶ **Video 1** The intrapancreatic pancreaticobiliary ducts and papilla were successfully observed in a patient with Billroth II reconstruction using forward-view echoendoscopy and the gel immersion technique.



▶ **Fig. 1** Magnetic resonance cholangiopancreatography demonstrated no apparent obstruction of the bile ducts.



▶ **Fig. 2** View during forward-view echoendoscopy showing that, although visualization of the intrapancreatic bile ducts and papilla was possible, detailed evaluation was difficult.

Endoscopic ultrasound (EUS) in patients with surgically altered anatomy is often difficult, and forward-view echoendoscopy (FV-EUS) has proven to be useful in such cases [1]. Recently, the gel immersion technique has been used for various endoscopic diagnostic and therapeutic procedures [2–4]. In the case reported here, the bile ducts and papilla were successfully observed in a patient with a Billroth II reconstruction using FV-EUS and the gel immersion technique (▶ **Video 1**).

An 82-year-old man who had undergone distal gastrectomy with Billroth II reconstruction for a gastric ulcer was referred to our center with elevated hepatobiliary enzymes and suspected cholangitis. Blood tests had not revealed an identifiable cause of hepatitis, while abdominal ultrasonography, computed tomography, and magnetic resonance cholangiopancreatography showed no bile duct dilatation and no obvious origin of the obstruction (▶ **Fig. 1**).

EUS was performed using a linear echoendoscope; however, scope insertion into

the afferent limb was difficult and the bile ducts could not be visualized in the stomach region. By switching to FV-EUS (TGF-UC260J; Olympus Medical Systems Corp., Tokyo, Japan), we safely guided the scope to the afferent limb with visualization of the anastomosis and successfully reached the ampulla of Vater. FV-EUS revealed mild dilatation of the common bile duct, with no obvious stones. The peripapillary delineation was unclear, making it difficult to detect the presence of small stones or tumors (▶ **Fig. 2**). For gel immersion, the duodenum was filled with gel (VISCOCLEAR; Otsuka Pharmaceutical Factory, Tokushima, Japan) via the scope working channel. The intrapancreatic pancreaticobiliary ducts and papilla could then be delineated clearly (▶ **Fig. 3**). Therefore, with the absence of stones/tumors confirmed, endoscopic retrograde cholangiopancreatography was deemed unnecessary. Thereafter, the patient's hepatobiliary enzyme levels slowly decreased. In the afferent limb, the gel tended to pool at the blind end and could be clearly observed.

FV-EUS plus gel immersion was considered to be the best combination for pancreaticobiliary examination in such patients with surgically altered anatomy.


Endoscopy_UCTN_Code_CCL_1AF_2AF

Acknowledgments

We would like to thank Editage (www.editage.com) for English language editing.

Competing interests

A. Katanuma has received speaker's fees from Olympus Co., Tokyo, Japan. The remaining authors declare that they have no conflict of interest.



► **Fig. 3** Gel immersion endoscopic ultrasound view clearly showing the papilla and intrapancreatic bile duct.

Bibliography

Endoscopy 2022; 54: E867–E868

DOI 10.1055/a-1841-6048

ISSN 0013-726X

published online 3.6.2022

© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14,
70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>

The authors

Tatsuya Ishii **Tsuyoshi Hayashi** **Kuniyuki Takahashi**, **Toshifumi Kin**, **Akio Katanuma**
Center for Gastroenterology, Teine Keijinkai Hospital, Sapporo, Hokkaido, Japan

Corresponding author

Tatsuya Ishii, MD
Center for Gastroenterology, Teine Keijinkai Hospital, 1-40, 12-chome, 1-jou, Maeda, Teine-ku, Sapporo 006-8555, Hokkaido, Japan
ttyisi141@outlook.com

References

- [1] Fusaroli P, Serrani M, Lisotti A et al. Performance of the forward-view echoendoscope for pancreaticobiliary examination in patients with status post-upper gastrointestinal surgery. *Endosc Ultrasound* 2015; 4: 336–341
- [2] Yano T, Nemoto D, Ono K et al. Gel immersion endoscopy: a novel method to secure the visual field during endoscopy in bleeding patients (with videos). *Gastrointest Endosc* 2016; 83: 809–811
- [3] Hanaoka N, Ishihara R, Matsuura N et al. Esophageal EUS by filling water-soluble lubricating jelly for diagnosis of depth of invasion in superficial esophageal cancer. *Gastrointest Endosc* 2015; 82: 164–165
- [4] Toyonaga H, Takahashi K, Kin T et al. Gel immersion technique for the examination and treatment of an ampullary tumor. *Endoscopy* 2022; 54: E115–E116