


Double-balloon enteroscopy-assisted polypectomy for a solitary jejunal Peutz–Jeghers polyp causing anemia



► **Fig. 1** Capsule enteroscopic view of a reddish tumor in the proximal small intestine.

A 53-year-old man with no family history of hereditary disease was referred to our hospital for evaluation of anemia. He had a 2-year history of postprandial nausea and discomfort in the upper abdomen. There was no hyperpigmentation of the oral mucosa or skin on physical examination. Esophagogastroduodenoscopy and colonoscopy performed at his previous hospital had revealed no specific findings. Laboratory data showed iron deficiency anemia, with a hemoglobin level of 6.0 g/dL. Abdominal computed tomography was unremarkable.

Capsule enteroscopy detected a reddish tumor in the proximal small intestine ► (**Fig. 1**), but there were no other polyps. Peroral double-balloon enteroscopy (DBE) identified a 30-mm polyp in the jejunum ► (**Video 1**), which had a slightly lobulated head and stalk ► (**Fig. 2a, b**). After ligating the stalk with a detachable snare to prevent bleeding, we achieved en bloc resection with polypectomy ► (**Fig. 2c**). Histopathological examination of the resected specimen revealed radially branching muscularis mucosa and growth of the lamina



► **Video 1** Double-balloon enteroscopy-assisted polypectomy for a solitary Peutz–Jeghers polyp in the jejunum.

propria with no atypia, indicating a diagnosis of a solitary Peutz–Jeghers polyp ► (**Fig. 3**). After undergoing this endoscopic treatment, the patient's abdominal symptoms immediately disappeared and his anemia improved.

Solitary Peutz–Jeghers polyp is defined as a hamartomatous polyp, having similar histopathological findings to the polyps found in Peutz–Jeghers syndrome, in patients without a family history or mucocutaneous pigmentation [1]. Solitary Peutz–Jeghers polyp is rare, especially in the jejunum, and can cause gastrointestinal bleeding and intussusception [2, 3]. Most jejunal cases have been treated with surgery and only a few cases have been endoscopically resected [2, 4]. Earlier studies reported that solitary Peutz–Jeghers polyps tended to present with lobulated and pedunculated morphology [1, 5]. Therefore, if suspected, endoscopists should carefully look for a stalk and consider endoscopic resection, even when the polyp is large.

In our patient, the use of DBE and a detachable snare allowed safe resection of a symptomatic solitary Peutz–Jeghers polyp in the jejunum, thereby avoiding the need for surgery.

Endoscopy_UCTN_Code_TTT_1AP_2AD

Acknowledgments


We would like to thank Editage (<https://www.editage.jp>) for English language editing.

Competing interests

The authors declare that they have no conflict of interest.



► **Fig. 2** Images during peroral double-balloon enteroscopy (DBE) showing: **a** a 30-mm lobulated polyp in the jejunum; **b** a thick stalk visible on careful endoscopic observation of the jejunal polyp; **c** appearance after DBE-assisted polypectomy with a detachable snare.



► **Fig. 3** Histopathological image of a solitary Peutz-Jeghers polyp showing radially branching muscularis mucosa and growth of the lamina propria without atypia.

The authors

Kazunori Onuma¹, Gota Sudo^{1,2}, Hiroyuki Inoue¹, Takahiro Kuriyama³, Yumemi Takada¹, Atsushi Yawata¹, Hiroshi Nakase²

- 1 Department of Gastroenterology and Hepatology, Hakodate Goryoukaku Hospital, Hakodate, Japan
- 2 Department of Gastroenterology and Hepatology, Sapporo Medical University School of Medicine, Sapporo, Japan
- 3 Division of Clinical Engineering, Hakodate Goryoukaku Hospital, Hakodate, Japan

Corresponding author

Gota Sudo, MD

Department of Gastroenterology and Hepatology, Hakodate Goryoukaku Hospital, 3-38, Goryoukaku-cho, Hakodate 040-8611, Japan
gotasudo@gmail.com

References

- [1] Lantz H, Doos WG, Affi A. Peutz-Jeghers-type hamartomatous polyp in a patient without Peutz-Jeghers syndrome. *Gastrointest Endosc* 2004; 60: 316–317
- [2] Chang CH, Cheng KS, Chou JW. Small-bowel solitary Peutz-Jeghers-type hamartomatous polyp treated with endoscopic polypectomy. *Clin Gastroenterol Hepatol* 2019; 17: e44
- [3] Endo K, Kawamura K, Murakami K et al. A case of jejunal solitary Peutz-Jeghers polyp with intussusception identified by double-balloon enteroscopy. *Clin J Gastroenterol* 2020; 13: 1129–1135
- [4] Sone Y, Nakano S, Takeda I et al. Solitary hamartomatous polyp of Peutz-Jeghers type in the jejunum resected endoscopically. *Gastrointest Endosc* 2000; 51: 620–622
- [5] Iwamura M, Aoyama Y, Suzuki S et al. Long-term outcome in patients with a solitary Peutz-Jeghers polyp. *Gastroenterol Res Pract* 2019; 2019: 8159072

Bibliography

Endoscopy 2022; 54: E906–E907
DOI 10.1055/a-1866-3758
ISSN 0013-726X
published online 1.7.2022
© 2022. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS

<https://eref.thieme.de/e-videos>



Endoscopy E-Videos is an open access online section, reporting on interesting cases and new techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online. Processing charges apply (currently EUR 375), discounts and waivers acc. to HINARI are available.

This section has its own submission website at
<https://mc.manuscriptcentral.com/e-videos>