




Low α -Thrombin/GPIb α Interaction Is a Potential Contributor to Platelet Hyper-reactivity in COVID-19 Patients

Younes Zaid^{1,2,3}  Loubna Khalki⁴ Farid Jalali⁵ Youssef Tijani⁴ Nabil Zaid^{1,6} Abdallah Naya³
Mounia Oudghiri³ Khadija Akarid⁷ Ejaife O. Agbani⁸ Fadila Guessous^{4,9}

¹ Department of Biology, Faculty of Sciences, Mohammed V University, Rabat, Morocco

² Research Center of Abulcasis University of Health Sciences, Rabat, Morocco

³ Immunology and Biodiversity Laboratory, Department of Biology, Ain Chock Faculty of Sciences, Hassan II University, Casablanca, Morocco

⁴ Laboratory of Neurosciences and Oncogenetics, Neurosciences and Cellular Physiology Team, Mohammed VI Center for Research & Innovation, Higher Institute of Biosciences and Biotechnology and Faculty of Medicine, Mohammed VI University of Sciences and Health, Casablanca, Morocco

⁵ Department of Gastroenterology, Saddleback Medical Group, Laguna Hills, California, United States

Address for correspondence Younes Zaid, PhD, Faculty of Sciences, 4 Ibn Battouta Av, Agdal, Rabat, Morocco (e-mail: y.zaid@um5r.ac.ma).

⁶ Health, Care, and Environment Laboratory, Higher Institute of Nursing Professions and Health Technology of Rabat, Morocco

⁷ Health and Environment Laboratory, Ain Chock Faculty of Sciences, Hassan II University of Casablanca, Morocco

⁸ Department of Physiology & Pharmacology, University of Calgary, Calgary, Canada

⁹ Department of Microbiology, Immunology and Cancer Biology, School of Medicine, University of Virginia, Charlottesville, Virginia

Thromb Haemost 2023;123:804–807.

Several studies have demonstrated that platelets can interact with severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), and consequently undergo programmed cell death, extracellular vesicle release, and increased platelet reactivity in coronavirus disease 2019 (COVID-19) patients in response to low doses of α -thrombin.^{1–3} Such hyper-sensitivity could be partially due to increased mitogen-activated protein kinase (MAPK) signaling pathway activation, protein kinase C delta (PKC δ) phosphorylation, and thromboxane synthesis.^{1,2} These posttranslational modifications are often triggered following activation of one or more receptors on the platelet surface.

GPIb α (CD42b) in the GPIb-IX-V receptor complex is the major binding site for α -thrombin associated with platelets, and through this function may support procoagulant activities and contribute to platelet activation and aggregation.⁴ Importantly, blocking α -thrombin binding to the N-terminal region of GPIb α (His1-Glu282) can be achieved by a blocking antibody (SZ2) that selectively inhibits the α -thrombin binding site on GPIb α .⁴ According to recent findings, GPIb α is the receptor through which SARS-CoV-2 spike protein binds to platelets as well as activates their increased expression of ligands.⁵ However, the role of GPIb α

in the α -thrombin-induced platelet hyper-responsiveness observed during SARS-CoV-2 infection is not well understood.

At the low enzyme concentration, we identified α -thrombin/GPIb α interaction as a novel mechanism triggering the hyper-reactivity observed in COVID-19 patients. Our findings showed that during SARS-CoV-2 infection, pretreatment of platelets with human anti-GPIb α antibody (SZ2) prevents platelet hyper-aggregation and degranulation. Interestingly, we found that platelets derived from patients with Bernard-Soulier syndrome (BSS) and who are infected with SARS-CoV-2 are not hyper-reactive. We thus have identified low α -thrombin-GPIb α interaction as a novel prothrombotic pathway which triggers the formation of procoagulant platelets in COVID-19 patients.

COVID-19 patients ($n = 10$) and COVID-19 patients with BSS ($n = 3$) who were admitted to the Cheikh Zaid Hospital (Rabat, Morocco) were included in this prospective, observational study conducted from December 7, 2020 to January 12, 2022. Patients with BSS have a homozygous mutation on GP9 which prevents GPIb α expression on platelets. All patients with COVID-19 were studied (on average) at

received

November 8, 2022

accepted after revision

March 14, 2023

accepted manuscript online

April 12, 2023

article published online

May 2, 2023

DOI <https://doi.org/10.1055/a-2072-0366>.

ISSN 0340-6245.

© 2023. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

5.8 ± 1.2 hours after receiving a nasopharyngeal swab that showed positivity for SARS-CoV-2. Sex- and age-matched (age: 47.5, interquartile range: 38–52.4, 50% female) healthy blood donors ($n = 10$) were used as controls. None of the patients or healthy volunteers were treated with antithrombotic drugs (either pre-morbidly or for the treatment of their COVID-19 infection), and that could affect platelet functions or coagulation. All human blood studies were approved by the Local Ethics Committee of Cheikh Zaid Hospital, Rabat, Morocco; Project: CEFCZ/PR/2020-PR04. Informed consent was obtained from each subject and all experiments were

conducted according to the principles set out in the Declaration of Helsinki.

Washed platelets were prepared as previously described.¹ Markers of platelet activation, α -granule release (CD62P or P-selectin expression) (\blacktriangleright Fig. 1Ai, Aii) and dense granule secretion, as assessed by ATP release (\blacktriangleright Fig. 1Aiii) and loss of mepacrine fluorescence (\blacktriangleright Fig. 1Aiv), were significantly enhanced in the presence of sub-threshold concentrations of α -thrombin in platelets from COVID-19 patients. We sought to investigate the contribution of GPIIb/IIIa to the regulation of platelet hyper-activation. As shown in \blacktriangleright Fig. 1A(i–iv), this




Fig. 1 (A) GPIIb/IIIa positively regulates human platelet degranulation induced by low doses of α -thrombin in COVID-19 patients. (Ai) Platelet P-selectin expression was measured (percent of CD62P positive platelets) in washed human platelets by flow cytometry at baseline in 10 healthy donors and 10 COVID-19 patients. Platelets were pretreated with human anti-GPIIb/IIIa antibody (S22, 20 μ g/mL) or its isotype IgG control for 5 minutes at 37°C. Degranulation was then initiated by α -thrombin at 0, 0.05 (priming dose), and 0.2 U/mL (positive control). Histogram represents the mean of data \pm standard deviation (SD) of plots for P-selectin (CD62P) expression ($n = 10$); ** $p < 0.01$; **** $p < 0.0001$. Statistical significance was analyzed using one-way analysis of variance (ANOVA) with subsequent Dunnett's t -test for comparison against a single group. (Aii) Effect of anti-GPIIb/IIIa on P-selectin (CD62P) expression in COVID-19 patients, as assessed by flow cytometry. Left plots represent resting platelets. Right plots represent platelets in the presence of a priming concentration of α -thrombin (0.05 U/mL). (Aiii) ATP release was measured by a Lumi-Aggregometer (Luciferase assay; Chrono-Lume, Chrono-log). Results are expressed as a measure of increase in luminescence and (Aiv) dense granule secretion was evaluated by measuring the loss of mepacrine fluorescence following activation by α -thrombin (0.05 U/mL). Histogram represents the mean of data \pm SD of (Aiii) ATP release (fold increase) and (Aiv) remaining mepacrine fluorescence ($n = 10$); **** $p < 0.0001$. The reference range (resting platelets) is represented by the shaded gray region. (B) Pretreatment with a human anti-GPIIb/IIIa antibody prevents COVID-19-induced potentiation of platelet aggregation while Bernard–Soulier Syndrome (BSS) platelets are not hyper-reactive. (Bi) Platelets were pretreated with the human anti-GPIIb/IIIa antibody (S22, 20 μ g/mL) for 5 minutes at 37°C. Aggregation was then initiated by low concentrations of α -thrombin (0.05 U/mL). Histogram represents the mean of data \pm SD of aggregation traces ($n = 10$); **** $p < 0.0001$. (Bii) Representative traces of platelet aggregation induced by a priming dose of α -thrombin (0.05 U/mL). (Biii) Specific blockade of the thrombin-binding site on GPIIb/IIIa, by either S22 or VM16d blocking monoclonal antibodies, prevented the potentiation of thrombin-induced platelet aggregation in COVID-19 patients. (Biv) Platelets isolated from COVID-19 patients with (or without) BSS (deficiency in GPIIb-VIX complex) were stimulated by low doses of α -thrombin (0.05 U/mL). (Bv) Representative traces of platelet aggregation (shown in Biv) induced by a priming dose of α -thrombin (0.05 U/mL). (C) Scheme of the potential role the α -thrombin/GPIIb/IIIa axis plays in platelet hyper-reactivity in COVID-19 patients. During SARS-CoV-2 infection, α -thrombin stimulates its high-affinity receptor GPIIb/IIIa leading to α - and dense granule release and platelet hyper-aggregation. Receptor blockade or deficiency prevents the hyper-coagulable state occurring in COVID-19 patients. IgG, immunoglobulin G.

process was prevented following pretreatment of platelets with a selective human anti-GPIb α antibody (S22), whereas a control immunoglobulin G (IgG) antibody had no effect. This suggests that SARS-CoV-2 triggered platelet degranulation in an α -thrombin- and GPIb α -dependent manner. This is consistent with a report that GPIb α is the major α -thrombin binding site on the platelet surface with evidence for a high affinity site within its extracellular domain adjacent to the von Willebrand factor (VWF) binding site.⁶ Also, at a higher α -thrombin concentration, the protease-activated receptors (PAR1 and PAR4) likely become significantly engaged and further enhance degranulation and extent of platelet hyper-activation and aggregation.⁷

Similarly, we assessed whether the α -thrombin-GPIb α axis plays a role in the hyper-aggregation of platelets from COVID-19 patients. As anticipated, priming, but not high concentration of α -thrombin-induced aggregation, was significantly inhibited in platelets that were pretreated with a human anti-GPIb antibody (S22) (Beckman Coulter, sodium azide free), as compared with COVID-19 and to COVID-19 control IgG groups (**Fig. 1Bi, Bii**). Thus, specific inhibition of the high-affinity binding site of α -thrombin on GPIb α was found to inhibit potentiated washed human platelet secretion and aggregation response, indicating that the α -thrombin/GPIb α axis can potentiate platelet function in the presence of suboptimal α -thrombin concentrations.

To further assert our finding based on the mouse-anti-human antibody, clone S22, that targets the anionic/sulfated tyrosine sequence 269 to 282 of GPIb α , we performed a similar experiment using a mouse anti-human antibody, clone VM16d (Abcam, dialyzed against phosphate-buffered saline in our laboratory to remove sodium azide), that maps the C-terminal flanking sequence 226 to 268 of GPIb α .⁸ Our results show that specific blockade of the thrombin-binding site on GPIb α , by either S22 or VM16d (**Fig. 1Biii**) blocking monoclonal antibodies, prevented the potentiation of thrombin-induced platelet aggregation in COVID-19 patients.

To confirm our pharmacological-based approach, the key role for GPIb α was assessed using platelets isolated from COVID-19 patients with BSS (deficiency in GPIb-V-IX complex causing an absence of GPIb expression). Platelets from these patients were found to have markedly diminished aggregation in response to low concentrations of α -thrombin compared with COVID-19 patients without BSS (**Fig. 1Biii, Biv**). Indeed, SARS-CoV-2 infection failed to trigger platelet hyper-aggregation in patients with BSS suggesting that GPIb α positively regulates platelet aggregation downstream of α -thrombin in COVID-19 patients (**Fig. 1C**).

BSS is characterized by prolonged bleeding time, thrombocytopenia,^{9,10} and giant platelets lacking the surface membrane glycoprotein GPIb of the GPIb-IX-V complex.¹⁰⁻¹² The GPIb-IX-V complex is a platelet-specific adhesion-signaling complex, and it consists of GPIb α linked to GPIb β via a disulfide bond and to GPIX and GPV non-covalently. Here, GPIb α is the major ligand-binding subunit and binds thrombin and the adhesive ligand VWF.⁸ Given

that platelets of BSS patients hardly express GPIb, BSS platelets lack GPIb-specific response to α -thrombin. Similarly, S22, an anti-GPIb α monoclonal antibody, is known to inhibit VWF binding to platelet and platelet aggregation.¹³ Thus, both S22 antibody-treated platelets and platelets of BSS lack GPIb α function and GPIb α -driven response to α -thrombin. Other factors may contribute to the diminished response of BSS platelets to thrombin and SARS-CoV-2 infection, including the additional loss of GPV and GPIX. The loss of surface-membrane glycoprotein in BSS platelets has been closely linked to in vitro and in vivo functional impairment.^{9,10}

Our finding was supported by a recent study¹⁴ that worked on S100A8/A9, also known as “calprotectin” or “MRP8/14,” an alarmin primarily secreted by activated myeloid cells with antimicrobial, proinflammatory, and prothrombotic properties. This research group identified the S100A8/A9-GPIb α axis as a novel targetable prothrombotic pathway inducing procoagulant platelets and fibrin formation, in particular in diseases associated with high levels of S100A8/A9, such as COVID-19.

Taken together, GPIb α interaction with low concentration α -thrombin is a potential contributor to the formation of procoagulant platelets in COVID-19 patients. Such interaction is probably only one aspect of many that likely regulate platelet activity in COVID-19 patients. These observations provide a valuable foundation for understanding the disease pathophysiology and to identify a new target for treatment options.

Data Availability Statement

The original contributions presented in the study are included in the article, and further inquiries can be directed to the corresponding author.

Ethical Approval Statement

Written informed consent was obtained from the individuals for the publication of any data included in this article.

Authors' Contribution

Y.Z., F.J., and F.G. designed the research. Y.Z., L.K., Y.T., N.Z., A.N., and M.O. performed the research and statistical analyses. Y.Z., L.K., K.A., and F.G. analyzed the data. Y.Z., E.O.A., and F.G. contributed to discussion and revised the manuscript. Y.Z. and F.G. wrote the manuscript. All authors read and approved the paper.

Funding

This study was supported by Balvi Filantropic Fund (PR-BLV-20220527-B19).

Conflict of Interest

None declared.

Acknowledgment

The authors thank Dr. Eric Boilard (Université Laval, Québec, Canada) for helpful discussions, and are grateful to the patients and their families.

References

- 1 Zaid Y, Puhm F, Allaey I, et al. Platelets can associate with SARS-CoV-2 RNA and are hyperactivated in COVID-19. *Circ Res* 2020;127(11):1404–1418
- 2 Manne BK, Denorme F, Middleton EA, et al. Platelet gene expression and function in patients with COVID-19. *Blood* 2020;136(11):1317–1329
- 3 Koupenova M, Corkrey HA, Vitseva O, et al. SARS-CoV-2 initiates programmed cell death in platelets. *Circ Res* 2021;129(06):631–646
- 4 Dörmann D, Clemetson KJ, Kehrel BE. The GPIb thrombin-binding site is essential for thrombin-induced platelet procoagulant activity. *Blood* 2000;96(07):2469–2478
- 5 Li T, Yang Y, Li Y, et al. Platelets mediate inflammatory monocyte activation by SARS-CoV-2 spike protein. *J Clin Invest* 2022;132(04):e150101
- 6 Sixma JJ, Schiphorst ME, Verweij CL, Pannekoek H. Effect of deletion of the A1 domain of von Willebrand factor on its binding to heparin, collagen and platelets in the presence of ristocetin. *Eur J Biochem* 1991;196(02):369–375
- 7 Ware J. A binding relationship with thrombin. *Blood* 2016;127(05):525–526
- 8 Andrews RK, Berndt MC. Bernard-Soulier syndrome: an update. *Semin Thromb Hemost* 2013;39(06):656–662
- 9 López JA, Andrews RK, Afshar-Kharghan V, Berndt MC. Bernard-Soulier syndrome. *Blood* 1998;91(12):4397–4418
- 10 Berndt MC, Gregory C, Chong BH, Zola H, Castaldi PA. Additional glycoprotein defects in Bernard-Soulier's syndrome: confirmation of genetic basis by parental analysis. *Blood* 1983;62(04):800–807
- 11 Nurden AT, Caen JP. Specific roles for platelet surface glycoproteins in platelet function. *Nature* 1975;255(5511):720–722
- 12 Clemetson KJ, McGregor JL, James E, Dechavanne M, Lüscher EF. Characterization of the platelet membrane glycoprotein abnormalities in Bernard-Soulier syndrome and comparison with normal by surface-labeling techniques and high-resolution two-dimensional gel electrophoresis. *J Clin Invest* 1982;70(02):304–311
- 13 Cauwenberghs N, Vanhoorelbeke K, Vauterin S, et al. Epitope mapping of inhibitory antibodies against platelet glycoprotein Ibalph reveals interaction between the leucine-rich repeat N-terminal and C-terminal flanking domains of glycoprotein Ibalph. *Blood* 2001;98(03):652–660
- 14 Colicchia M, Schrottmaier WC, Perrella G, et al. S100A8/A9 drives the formation of procoagulant platelets through GPIb α . *Blood* 2022;140(24):2626–2643