

Breathing Motion Pattern in Cyclists: Role of Inferior against Superior Thorax Compartment



Authors

Ana Luiza Lopes¹, Karine Jacon Sarro¹, Isabella Martins Rodrigues¹, Richard Diego Leite², Carlo Massaroni³, Paulo Roberto dos Santos Amorim⁴, Pietro Cerveri^{5, 6}, Amanda P. Silvatti⁴

Affiliations

- 1 Faculdade de Educação Física, Universidade Estadual de Campinas, Campinas, Brazil
- 2 Centro de Educação Física e Desportos, Universidade Federal do Espírito Santo, Vitória, Brazil
- 3 Unit of Measurements and Biomedical Instrumentation, Università Campus Bio-Medico di Roma, Roma, Italy
- 4 Departamento de Educação Física, Universidade Federal de Viçosa, Viçosa, Brazil
- 5 Department of Electronics, Information and Bioengineering, Politecnico di Milano, Milano, Italy
- 6 Center for Intelligent Technologies in Sleep Medicine, Istituto Auxologico Italiano Istituto di Ricovero e Cura a Carattere Scientifico, Milano, Italy

Key words

respiratory, mechanics, cycling

accepted 15.11.2023

accepted manuscript online 15.11.2023

published online 2024

Bibliography

Int J Sports Med

DOI 10.1055/a-2211-9421

ISSN 0172-4622

© 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Georg Thieme Verlag, Rüdigerstraße 14,
70469 Stuttgart, Germany

Correspondence

Ana Luiza Lopes

Universidade Estadual de Campinas

Faculdade de Educação física

Av. Érico Veríssimo, 701 – Cidade Universitária Zeferino Vaz

13083-872 Campinas

Brazil

Tel.: 31971716143

analuiza.castrolopes@gmail.com

ABSTRACT

The thoracoabdominal breathing motion pattern is being considered in sports training because of its contribution, along with other physiological adaptations, to overall performance. We examined whether and how experience with cycling training modifies the thoracoabdominal motion patterns. We utilized optoelectronic plethysmography to monitor ten trained male cyclists and compared them to ten physically active male participants performing breathing maneuvers. Cyclists then participated in a self-paced time trial to explore the similarity between that observed during resting breathing. From the 3D coordinates of 32 markers positioned on each participant's trunk, we calculated the percentage of contribution of the superior thorax, inferior thorax, and abdomen and the correlation coefficient among these compartments. During the rest maneuvers, the cyclists showed a thoracoabdominal motion pattern characterized by an increased role of the inferior thorax relative to the superior thorax ($26.69 \pm 5.88\%$, $34.93 \pm 5.03\%$; $p = 0.002$, respectively), in contrast to the control group ($26.69 \pm 5.88\%$; $25.71 \pm 6.04\%$, $p = 0.4$, respectively). In addition, the inferior thorax showed higher coordination in phase with the abdomen. Furthermore, the results of the time trial test underscored the same pattern found in cyclists breathing at rest, suggesting that the development of a permanent modification in respiratory mechanics may be associated with cycling practice.

Introduction

Cycling is a demanding endurance sport that requires high physical fitness and stamina. Professional cyclists exhibit specific physiological responses such as considerable reliance on fat metabolism, even at high power outputs, and several neuromuscular adaptations [1–3]. Respiratory efficiency is also critical for cyclists [4–6]. Through training, they can improve oxygen uptake, which helps to improve endurance and delay the onset of fatigue during prolonged efforts [6–8]. In particular, they develop distinctive breathing patterns at high workloads that are distinguished by a lack of tachypnoeic shift, as they maintain increased pulmonary ventilation by increasing tidal volume rather than respiratory frequency of breathing [5, 9]. It is suggested that this breathing adaptation can enhance metabolic efficiency, partly accounting for the VO_2 kinetics [1].

While all such physiological adaptations are well documented, there is insufficient information regarding respiratory mechanics, described as thoracoabdominal (THA) motion patterns, i. e. displacement and coordination of the rib cage and abdomen [10]. Generally speaking, THA motion patterns modify according to the physical activity intensity, being dependent also on other factors such as gender, posture, and motion task [11–14]. Conversely, long-term sports training can permanently modify the THA motion patterns (evidenced by measurements taken during rest breathing maneuvers), to the advantage of better respiratory performance [15–17].

Especially, optimized lung and chest wall compliance, required to progressively suppress dyspnea and minimize the work of breathing, elicits respiratory mechanics adaptation [18, 19]. At rest and during exercise, there are distinct patterns of muscular recruitment. During rest, the rib cage muscles (as external intercostals, scalene, sternocleidomastoid) and the diaphragm act as pressure generators, and the abdominal muscle is not active. Instead, during exercise, the diaphragm behaves essentially as a flow generator because its shortening ability is greater than the inspiratory rib cage muscles, and the rib cage and abdominal muscles (such as transversus abdominis and internal and external oblique) are pressure generators [20, 21].

Among different non-invasive technologies to measure chest wall displacements, optoelectronic systems have been largely adopted, which record the 3D position of physical markers distributed homogeneously on the trunk surface [15, 16, 22–24]. Traditionally, the shape of the chest wall is partitioned into three functional compartments independent of each other [25]. This compartmental division provides information on respiratory muscles' action and assesses breathing movement patterns [18].

During pedaling, breathing muscles act as a crucial postural function, where optimal relation between the diaphragm, abdominal wall, parasternal, and pelvic floor provides production, transfer, and control of the active force [26]. In particular, the bike sitting posture may induce physical constraints to THA expansion and could affect respiratory mechanics. These differences in thoracoabdominal patterns become particularly prominent during moderate-intensity efforts, where there is an expected gradual increase in ventilation aligned with alveolar ventilation [27] and respiratory function [28, 29]. This phenomenon can be observed, for instance, during a time trial race that involves cyclists completing a pre-

etermined distance as swiftly as possible. Such races have been utilized in laboratory settings to simulate endurance events and predict aerobic performance indicators like VO_2 and lactate threshold [30].

Since 2008, our group has been investigating factors affecting THA motion patterns, such as physical activities, exercise, long-term training, and more in general anamnestic data [15, 16, 22, 31]. Experienced swim training was shown to increase coordination of the THA volumes and the ribs motion [22], and the contribution of the abdominal compartment [15]. Experienced ballet dancers, both men and women, showed the predominant respiratory contribution of the upper thorax and abdomen [16]. Interestingly, age was found to be a weak predictive factor for changes in breathing motion patterns, contrary to what is commonly believed in the field of respiratory physiology [31].

Capitalizing on such previous studies, this work aimed to investigate the THA breathing motion pattern in experienced cyclists compared to non-cyclists. To provide further clarity, a secondary objective was to examine potential disparities in the THA motion pattern of experienced cyclists during periods of rest and exercise, with the intention of evaluating the consistency and stability of such patterns. To achieve these objectives, our study involved subjecting the cyclists to a targeted simulated 20-km time trial.


Material and Methods

Participants and experimental protocol

Twenty healthy and non-smoker participants were enrolled in this study. They were divided into two sets, respectively control and cyclist groups. The control group comprised ten males physically active in different recreational sports, none of whom had ever been involved in intensive cycling practice. Every participant in the control group engaged in physical exercise for more than 3 hours per week. The cyclist group was composed of ten trained cyclist males with at least five years of competitive experience on speed bikes, who regularly participate in regional cycling events. The acquisition protocol involved a pulmonary function test performed by both groups. Specifically, three maneuvers of forced vital capacity were monitored first throughout spirometry. Then breathing maneuvers at rest, namely quiet breathing, and vital capacity trials, were monitored by optoelectronic plethysmography (► **Fig. 1a**). The cyclist group performed a simulated 20-km time-trial test (► **Fig. 1b**). The study was approved by the University Ethics Committee (Number 59773616.0.0000.5153) in compliance with the Declaration of Helsinki, and all participants provided written informed consent.

Pulmonary function test

To compare the pulmonary function, a spirometer (Micro-medical, Rochester, Kent, England) was used. The control and cyclist groups performed three maneuvers of the forced vital capacity according to a standard protocol in a sitting posture [32]. Measurements were obtained for forced vital capacity (FVC), which is the greatest total amount of air expired, and for the forced expiratory volume in one second (FEV_1), which is the volume delivered in the first second of a forced vital capacity maneuver according to the guidelines estab-



► **Fig. 1** General protocol and tests performed.

lished by the American Thoracic Society/European Respiratory Society [33]. These data were expressed as mean and percentages of predicted values for the Brazilian population, according to [34].

THA – movement patterns by three-dimensional kinematics of the trunk compartments

Thirty-two retro-reflective markers were placed on the trunk of the participants [25]. The markers were positioned as a grid, using anatomical references allowing trunk division into three compartments namely superior thorax (ST), from 2nd rib to the xiphoid process, inferior thorax (IT), from this line to the 10th rib, and abdomen (AB), from this line to transverse of the abdomen [16]. The 3D coordinates of the markers were acquired with eleven OptiTrack Prime 17 W cameras (NaturalPoint, Corvallis, OR, USA, 360 Hz), positioned around the participants. A low pass, cutoff frequency of 10 Hz, was used to filter the marker trajectories. From the filtered marker locations, the compartmental volumes were calculated frame by frame using a volumetric convex hull method in the software Visual 3D (C-Motion, USA) [23]. The variation of the compartmental volume was expressed as a function of the time, divided into breathing cycles, defined as the beginning of inspiration to the end of expiration. Considering n breathing cycles, the percentage of contribution of each compartment (%ST, %IT, %AB) was computed as:

$$\begin{aligned}
 \%ST &= \frac{1}{n} \sum_{i=1}^n \left(\% \frac{ST_i}{V_{tot_i}} \right) \\
 \%IT &= \frac{1}{n} \sum_{i=1}^n \left(\% \frac{IT_i}{V_{tot_i}} \right) \\
 \%AB &= \frac{1}{n} \sum_{i=1}^n \left(\% \frac{AB_i}{V_{tot_i}} \right)
 \end{aligned}
 \quad (1)$$

where V_{tot} is the total trunk volume in the i^{th} breathing cycle. We also calculated the relative ratio between percentage of contribution of superior thorax and abdomen, and inferior thorax and abdomen. In order to assess the coordination among the THA compartments, the

correlation coefficient was computed by means of the cross-correlation between the pairs of the compartmental volume during the inspiration phase. Positive and negative correlations indicated coordinated and paradoxical movements, respectively.

Breathing maneuvers at rest

Participants (control group and cyclist group) were seated on a chair without back support, with shoulder abduction, forearms leaned on a rigid support, 90° of knee flexion, and feet on the ground (► **Fig. 1a**). Following an adaptation period (at least 2 min) to the experimental setup, they performed two quiet breathing trials (30 s, each). Then, they performed two trials of five cycles of vital capacity, characterized by five periods of maximum inspiration followed by maximum forced expiration. The participants were directed to perform this maneuver using a verbal cue.

Simulation of the 20-km time trial

The cyclists performed a self-paced time trial in laboratory settings (► **Fig. 1b**) using their bicycle coupled to a cycle trainer (Compu-trainer ProLab 3D, Racermate, Seattle, WA, USA). The rear wheel of the personal bike was changed to Powermeter equipment (PowerTap, Saris, Madison, USA). The 20-km time trial protocol was configured in the Computrainer software as a flat course and consisted of a 10-min warm-up, and a 20-km time trial simulated test performed in the shortest time possible. The participants received verbal encouragement stimulus when they were nearly to achieve a distance of 5, 10, 15 km, and at the end of the test (20 km). They were instructed to maintain their hands at brake hoods enabling trunk markers' tracking. The THA movement pattern was evaluated one minute earlier each step in 5, 10, 15, and 20 kilometers. Heart rate (HR) and ratings of perceived exertion (RPE) were acquired at the following distances, namely 5, 10, 15, and 20 km. HR was measured using a Polar Fitness tracker chest strap and the data was acquired by smartphone app (Elite HRV). The RPE was measured using Borg's CR-10 scale (1 to 10) [35], where 1 corresponds to 'no exertion' and 10 corresponds to 'extreme/maximal exertion'.

The participants received standardized instructions for each measure using this scale before the exercise session.

Statistical analysis

The normality of the data was evaluated using the Shapiro-Wilk test. The percentage of contribution and correlation coefficient were not normally distributed and the arcsine transformation and Fisher's z-transformation were applied, respectively. Quiet breathing and vital capacity maneuvers were analyzed separately. To analyze breathing maneuvers at rest, the Mann-Whitney test was adopted to compare the descriptive participants' characteristics between groups. Repeated measure ANOVAs were then considered to evaluate the percentage of contribution and correlation coefficient both with one between factors (groups: control group and cyclist group) and one within factor (contribution of compartments: superior thorax, inferior thorax and abdomen or pairs compartments: superior thorax × inferior thorax, superior thorax × ab-

domen and inferior thorax × abdomen, respectively). The repeated measure ANOVA with one within factor (time) was applied to evaluate the power output and HR, and for the RPE the Friedman test was used. To analyze the breathing motion pattern during the 20-km time trial, the repeated measure ANOVA with two within factors (contribution of compartments: superior thorax, inferior thorax and abdomen or pairs compartments: superior thorax × inferior thorax, superior thorax × abdomen and inferior thorax × abdomen and time: quiet breathing, vital capacity, 5, 10, 15, 20 km) were employed, respectively. Mauchly's test of sphericity was performed and the Huynh-Feldt correction was exploited to correct the variability in experimental error. If significant F-ratios were detected, a Bonferroni post-hoc comparison was applied to determine where the differences occurred. Cohen's *d* standardized effect sizes and confidence intervals (set at 95%) were reported when appropriate. Statistical significance was set at ($\alpha = 5\%$) for all analyses and was performed using SPSS version 18.0.


► **Table 1** Description of the study sample (mean ± SD) for the subjects' characteristics, pulmonary function, and training.

	Cyclists	Control
Subject characteristics		
Age (years)	42.7 ± 6.3	41.9 ± 8.7
Height (m)	1.75 ± 0.07	1.7 ± 0.08
Weight (kg)	75.9 ± 7.4	81.4 ± 10.2
Pulmonary function		
FEV ₁ (l)	3.9 ± 0.6	3.6 ± 0.5
FEV ₁ (% predict)	109.7 ± 19.3	105.2 ± 15.8
FVC (l)	4.3 ± 0.8	4.4 ± 0.7
FVC (% predict)	100.3 ± 20.6	104.8 ± 14.8
Training description		
Time per week (h)	11.2 ± 2.1	5.9 ± 2.7
Experience (years)	19.2 ± 10.9	22.6 ± 14*

Results

The two groups did not differ significantly in terms of age, height, and weight (► **Table 1**). All the participants showed pulmonary function results equal to or higher than predicted, and no significant differences between groups were found.

Regarding the relative ratio between inferior thorax and abdomen, the cyclist group exhibited values of 0.81 for quiet breathing and 0.89 for vital capacity, whereas the control group showed values of 0.56 for quiet breathing and 0.60 for vital capacity. The ANOVA analyses of the percentage of contribution in quiet breathing (► **Fig. 2a**) showed that the effect of the factor group was not significant ($p = 0.458$). The effect of the factor compartment was significant ($p < 0.001$) and inferior thorax and superior thorax had a similar contribution, while abdomen had a significantly higher contribution than both inferior thorax (AB: 41.6 ± 7.9%, 95%CI [37.8, 45.3]; IT: 28.3 ± 7.7%, 95%CI [24.7, 31.9]; $p = 0.002$. $d = 1.68$)



► **Fig. 2** Boxplot of the percentage of contribution of superior thorax (ST), inferior thorax (IT), and abdomen (AB) during quiet Breathing (QB) (a) and vital Capacity (VC) (b) for the control and cyclist group; + : outlier data.

► **Table 2** Mean (\pm SD) values of the descriptive characteristics and thoracoabdominal movement pattern during TT20.

	5 km	10 km	15 km	20 km
RPE	3.6 \pm 0.5	5.1 \pm 0.67 [*]	6.87 \pm 0.76 [◊]	9.25 \pm 0.31 ^{◊†}
HR (BPM)	171.5 \pm 3.45	173.4 \pm 3.34	176.4 \pm 2.69	184 \pm 3.18 ^{◊†}
Power Output (W)	245.76 \pm 14.93	239.78 \pm 14.16	237.41 \pm 15.69	278.87 \pm 20.81 [†]
Percentage of contribution				
%ST	23.59 \pm 3.06	23.60 \pm 3.66	24.51 \pm 3.23	23.59 \pm 4.60
%IT	32.88 \pm 2.33	32.94 \pm 3.54	32.47 \pm 2.84	33.01 \pm 2.91
%AB	43.51 \pm 3.28	43.45 \pm 3.78	43.01 \pm 4.20	43.38 \pm 5.22
ST:AB	0.54	0.53	0.59	0.54
IT:AB	0.76	0.76	0.76	0.76
Correlation coefficient				
ST \times IT	0.93 \pm 0.03	0.92 \pm 0.04	0.88 \pm 0.07	0.89 \pm 0.03
ST \times AB	0.88 \pm 0.06	0.87 \pm 0.06	0.83 \pm 0.09	0.84 \pm 0.05
IT \times AB	0.94 \pm 0.04	0.94 \pm 0.04	0.93 \pm 0.03	0.92 \pm 0.03
*significantly higher than 5 km; ◊significantly higher than 10 km; † significantly higher than 15 km. Significance level $p < 0.05$.				

and superior thorax (30.1 \pm 5.1 %, 95 %CI [27.6, 32.4]; $p = 0.001$, $d = 1.72$). Additionally, there was no significant interaction effect between the groups.

Again, in the vital capacity maneuver (► **Fig. 2b**), the ANOVA analyses showed that the effect of the factor group was not significant ($p = 0.336$), and the factor compartment was significant ($p < 0.001$), the abdomen contributing significantly more than inferior thorax (AB: 42.0 \pm 8.4 %, 95 %CI [38.1, 45.9]; IT: 30.8 \pm 6.8 %, 95 %CI [27.6, 34.0], $p = 0.003$, $d = 1.46$) and superior thorax (27.2 \pm 5.9 %, 95 %CI [24.4, 29.9], $p < 0.001$, $d = 2.03$). In the cyclist group, the superior thorax had a lower contribution than the inferior thorax (ST: 25.7 \pm 6.0 %, 95 %CI [21.4, 30.0]; IT: 34.9 \pm 5.1 %, 95 %CI [22.5, 30.9], $p = 0.002$, $d = 1.65$) and abdomen (39.3 \pm 4.8 %, 95 %CI [35.9, 42.8], $p = 0.004$, $d = 2.49$). In the control group, the abdomen had a higher contribution than the inferior thorax (AB: 44.6 \pm 10.5 %, 95 %CI [37.1, 52.1]; IT: 26.7 \pm 5.8 %, 95 %CI [31.3, 38.5], $p = 0.001$, $d = 2.11$) and superior thorax (28.7 \pm 5.6 %, 95 %CI [24.6, 32.7], $p < 0.001$, $d = 1.88$). Additionally, there was a significant interaction effect between the factors and the inferior thorax contribution of the cyclist group, which was significantly higher than the control group ($p = 0.004$, $d = 1.46$).


Both groups presented strong positive correlation coefficient values for quiet breathing and vital capacity (0.9 to 0.97). The cyclist group had correlation coefficient values significantly higher than the control group in quiet breathing ($p = 0.01$) and in vital capacity ($p = 0.034$). The inferior thorax \times abdomen compartmental pair had significantly higher correlation coefficient values than superior thorax \times inferior thorax and superior thorax \times abdomen and superior thorax \times inferior thorax was significantly higher than superior thorax \times abdomen. In both maneuvers, Cohen's d values ranged from 0.22 to 0.99 ($p < 0.001$). Both groups presented this coordination pattern in quiet breathing. However, during vital capacity, this pattern changed. While the control group had similar coordination between abdomen \times inferior thorax and superior thorax \times inferior thorax, the abdomen \times inferior thorax of the cyclist group was higher than the superior thorax \times inferior thorax. Sum-

marized statistical results are available in the supplementary material.

Due to missing data of some retro-reflective markers placed on the trunk of two cycling athletes during time trial, the results were reported only for eight participants (► **Table 2**). The RPE was different from each step of the time trial ($\chi^2 = 22.41$, $p < 0.001$; $5 > 10 > 15 > 20$ km). In the last step of the time trial (20 km), HR was significantly higher than 5 km ($p = 0.001$), 10 km ($p = 0.011$), and 15 km ($p = 0.017$). Also, significant differences in power output were found ($p < 0.001$) and the 20 km was higher compared to 15 km ($p = 0.021$).

Related to the percentage of contribution, compartments were different from each other ($p < 0.001$, ► **Table 2**), the abdomen was significantly higher than inferior thorax (AB: 43.34 \pm 3.98, 95 %CI [41.9, 44.78]; IF: 32.82 \pm 2.8, 95 %CI [31.81, 33.83], $p = 0.004$, $d = 3.08$) and superior thorax (23.82 \pm 3.52, 95 %CI [22.55, 25.10]; $p < 0.001$, $d = 5.31$), and inferior thorax was significantly higher compared to the superior thorax ($p = 0.004$, $d = 2.87$). The percentage of contribution of each compartment was the same in all the steps of the 20-km time trial (5, 10, 15, 20 km, $p = 0.879$). While the ratio superior thorax and abdomen oscillated between the steps, inferior thorax and abdomen was stable during all of the 20-km time trial (► **Table 2**). All compartment pairs presented strong correlation coefficient values. The superior thorax \times abdomen was significantly lower than superior thorax \times inferior thorax and inferior thorax \times abdomen ($p = 0.004$, $p = 0.001$, respectively). In addition, the correlation coefficient was statistically different between steps of the 20-km time trial ($p < 0.001$). Step 5 km was higher than 20 km ($p = 0.039$, $d = 0.72$), and 10 km presented higher values compared to 15 km ($p = 0.012$, $d = 0.38$) and 20 km ($p = 0.036$, $d = 0.55$).

Since no significant differences were observed among the distances of 5 km, 10 km, 15 km, and 20 km, we employed the mean value for comparison with the rest maneuvers. No statistical difference was found when the percentage of contribution of cyclists at rest were compared with the 20-km time trial ($p = 0.19$, ► **Fig. 3a**). The relative ratios between superior thorax and abdomen, likewise inferior thorax and abdomen, were higher in the quiet breathing



► **Fig. 3** Boxplot of the percentage of contribution of each compartment superior thorax (ST), inferior thorax(IT), and abdomen (AB) (a). Boxplot of the correlation coefficients of superior thorax× inferior thorax(ST×IT), superior thorax×abdomen (ST×AB), and inferior thorax×abdomen (IT×AB) (b) during the quiet breathing (QB) and vital capacity (VC) and during the 20-km time trial (20-TT).

► **Table 3** Summarized statistical results of quiet breathing, vital capacity and 20-km time trial for both groups and variables.

Summarized Statistical Results – Rest Maneuvers		
	Control Group	Cyclist Group
Percentage of Contribution		
Quiet Breathing	AB>ST>IT	AB>IT=ST
Vital Capacity	AB>ST=IT	AB=IT>ST
20-km Time Trial	n. a.	AB>IT>ST
Correlation Coefficient		
Quiet Breathing	AB×IT=ST×IT=ST×AB	AB×IT>ST×IT=ST×AB
Vital Capacity	AB×IT=ST×IT>ST×AB	AB×IT=ST×IT>ST×AB
20-km Time Trial	n. a.	AB×IT=ST×IT>ST×AB
Legend: AB: abdomen; IT: inferior thorax; ST: superior thorax; n. a.: not applicable.		

and vital capacity than in the last step of the test. The 20-km time trial correlation coefficients of all compartmental pairs, superior thorax× inferior thorax, superior thorax× abdomen, and inferior thorax× abdomen, were significantly lower than quiet breathing and vital capacity ($p<0.001$, ► **Fig. 3b**), and the Cohen’s *d* values ranged from 2.32 to 2.78. Concise summary of the statistical outcomes was reported (► **Table 3**).

Discussion

Globally, as verified in rest breathing maneuvers, the abdomen contribution was greater than superior thorax and inferior thorax con-

tributions individually in both groups. Specifically, cyclists demonstrated a specific THA motion pattern characterized by an increased role of the inferior thorax compartment with respect to the superior thorax (► **Fig. 1**), in opposition to the control group. In addition, inferior thorax contribution showed higher coordination with abdomen. The 20-km time trial test results highlighted the same THA motion pattern, which was kept stable throughout the trial, thus confirming the development of a permanent modification of the breathing mechanics in the cyclist group (► **Table 2**). Since no significant differences in anthropometric and pulmonary function values (► **Table 1**) were found between control group and cyclist group groups, this suggests that the differences in THA motion pattern may be attributed to cycling training.

While the statistical analysis supported the conclusions, the small cohort reduced the generalization of the results. We did not calculate post hoc power analyses given current statistical recommendations, but we do report confidence intervals for our primary results, which may be a superior means of interpreting the null and positive effects. In addition, the trained cyclists involved were not professionals; however, differences in the THA motion pattern with non-cyclists were detected.

Early studies showed that intense engagement in sports may easily elicit long-lasting THA patterns, with the greatest contribution to ventilation provided by the abdomen compartment [18, 36, 37], and our results agree with these findings. It is well known that physical activity recruits the abdominal muscles, featuring a higher magnitude than rib cage muscles, leading to a rise in diaphragmatic oxidative capacity [38]. Nonetheless, it was clear-

ly documented that exercise training improves diaphragm antioxidant capacity and endurance [39]. More recent studies have shown that sport specificity is responsible for developing diverse changes in THA patterns [15–17], triggering specific improvements in breathing mechanics efficiency [40]. Nonetheless, it was argued that postural constraints may further contribute to the development of custom THA patterns [1]. In Silvatti and coauthors [15], the authors argued that long-term swimming training contributed to the development of very specific THA motion patterns. Likewise, professional ballet dancers were able to keep unchanged breathing mechanics in quiet breathing and vital capacity maneuvers [16]. The effect of training in Pilates practitioners was documented to modify the thoracoabdominal motion pattern by increasing the contribution of the abdomen [17]. A raise in the force, in both appositional and insertional of the diaphragm, related to the inferior thorax compartment, was already found in cyclists [12, 41].

The 20-km time trial simulation allowed us to use an ecological methodology to understand the THA pattern of cyclists, with a major contribution of inferior thorax than superior thorax, along with different coordination with abdomen. As a matter of fact, all the cyclists showed a U power-output strategy, with higher values at the beginning and end of the test. Nonetheless, as shown, the THA pattern was not different during the steps evaluated, highlighting their capacity to maintain the breathing motion pattern during the whole test. The reduced coordination between superior thorax and abdomen can be motivated by the limited mechanical linkage between them, which is intentional to minimize elastic work when moving the chest wall [42]. We may argue that another possible explanation is the difference in the inspiratory and expiratory reserve volumes, which are relatively greater in the rib cage and abdominal compartment [18], respectively, and prevent paradoxical motion of any chest wall compartment during exercise [21]. The greater percentage of contribution of the inferior thorax found in our results, and the long-lasting pattern due to training, was further supported by literature outcomes about untrained individuals performing cycling [12, 18, 43]. All studies pointed out the major contribution of superior thorax to the breathing pattern. In trained cyclists, it can be hypothesized that the inferior thorax mechanics have a greater role in intra-abdominal pressure regulation, contributing the most to spinal and pelvic floor stabilization [44, 45], respiratory function, and power output [41, 45, 46]. As a result, we argue that the planning of specific inspiratory muscle training is to be considered to further improve the performance of competitive cyclists, in agreement with [5] who suggested that inspiratory muscle training attenuates the perceptual response to maximal incremental exercise.

While our findings provide valuable insights into the differences between cyclists and non-cyclists, it is essential to acknowledge the inherent limitations of our approach. As our study employed a cross-sectional design, we lacked information about cyclists' breathing patterns from the inception of their training journey. Additionally, we did not track cyclists over the course of their progression from novice to experienced cyclists or monitor changes in their breathing patterns over time. To gain a more comprehensive understanding of the impact of cycling training on the thoracoabdominal motion pattern, future research should incorporate longitudinal designs. Longitudinal studies would allow for tracking the

trajectory of individuals' thoracoabdominal motion patterns from the beginning of cycling training, potentially capturing the magnitude of its influence and uncovering causal relationships. Additionally, differentiating patterns between novice and experienced cyclists warrants investigation. Such studies would significantly contribute to a more in-depth exploration of the dynamic nature of thoracoabdominal motion in the context of cycling training.

In summary, cyclists demonstrated a specific thoracoabdominal motion pattern characterized by an increased role of the inferior thorax compartment with respect to the superior thorax. In addition, the inferior thorax contribution showed higher coordination with the abdomen. During the simulated 20-km time trial, the significant participation of the abdomen and inferior thorax was retained during all the tests, which can be associated with the cyclists' capacity to maintain during the effort imposed. Finally, the breathing mechanics outlined in this study can assist in designing specific programs aimed at improving the performance of competitive cyclists by enhancing the functionality of their inspiratory muscles, mainly the diaphragm.

Fundings

FAPEMIG – Fundação de Amparo à Pesquisa do Estado de Minas Gerais – PEE-00596–14 and 12040/2017

CNPq – Conselho Nacional de Desenvolvimento Científico e Tecnológico – 481391/2013–4

CAPES – Coordenação de Aperfeiçoamento de Pessoal de Nível Superior – 2011/10–7 and 08/2014

Acknowledgement

The authors special thank the laboratory technician, Nara Luiza Soares for her collaboration and commitment to this study. The authors are grateful to the volunteers for their participation.

Conflict of Interest

The authors declare that they have no conflict of interest.

References

- [1] Faria EW, Parker DL, Faria IE. The science of cycling: Physiology and training – Part 1. *Sports Med* 2005; 35: 285–312
- [2] Atkinson G, Davison R, Jeukendrup A et al. Science and cycling: Current knowledge and future directions for research. *J Sports Sci* 2003; 21: 767–787
- [3] Röthlin P, Wyler M, Müller B et al. Body and mind? Exploring physiological and psychological factors to explain endurance performance in cycling. *Eur J Sport Sci* 2023; 23: 101–108
- [4] Lucía A, Carvajal A, Calderón FJ et al. Breathing pattern in highly competitive cyclists during incremental exercise. *Eur J Appl Physiol Occup Physiol* 1999; 79: 512–521
- [5] Romer LM, McConnell AK, Jones DA. Effects of inspiratory muscle training on time-trial performance in trained cyclists. *J Sports Sci* 2002; 20: 547–562

- [6] Dionne JF, Lajoie C, Gendron P et al. Physiological and psychological adaptations of trained cyclists to spring cycling camps. *J Hum Kinet* 2018; 64: 137–146
- [7] Jeukendrup AE, Craig NP, Hawley JA. The bioenergetics of world class cycling. *J Sci Med Sport* 2000; 3: 414–433
- [8] Lucía A, Hoyos J, Chicharro JL. Physiology of professional road cycling. *Sports Med* 2001; 31: 325–337
- [9] Lucía A, Hoyos J, Pardo J et al. Effects of endurance training on the breathing pattern of professional cyclists. *Jpn J Physiol* 2001; 51: 133–141
- [10] Aliverti A, Dellacà R, Pelosi P et al. Compartmental analysis of breathing in the supine and prone positions by optoelectronic plethysmography. *Ann Biomed Eng* 2001; 29: 60–70
- [11] Sanna A, Bertoli F, Misuri G et al. Chest wall kinematics and respiratory muscle action in walking healthy humans. *J Appl Physiol* 1999; 87: 938–946
- [12] Vogiatzis I, Aliverti A, Golemati S et al. Respiratory kinematics by optoelectronic plethysmography during exercise in men and women. *Eur J Appl Physiol* 2005; 93: 581–587
- [13] Romagnoli I, Gorini M, Gigliotti F et al. Chest wall kinematics, respiratory muscle action and dyspnoea during arm vs. leg exercise in humans. *Acta Physiol (Oxf)* 2006; 188: 63–73
- [14] Layton AM, Garber CE, Thomashow BM et al. Exercise ventilatory kinematics in endurance trained and untrained men and women. *Respir Physiol Neurobiol* 2011; 178: 223–229
- [15] Silvatti AP, Sarro KJ, Cerveri P et al. A 3d kinematic analysis of breathing patterns in competitive swimmers. *J Sports Sci* 2012; 30: 1551–1560
- [16] Rodrigues IM, Bernardina GRD, Sarro KJ et al. Thoracoabdominal breathing motion pattern and coordination of professional ballet dancers. *Sports Biomech* 2019; 18: 51–62
- [17] Campos JL, Vancini RL, Zanoni GR et al. Effects of mat Pilates training and habitual physical activity on thoracoabdominal expansion during quiet and vital capacity breathing in healthy women. *J Sports Med Phys Fitness* 2019; 59: 57–64
- [18] Aliverti A, Cala SJ, Duranti R et al. Human respiratory muscle actions and control during exercise. *J Appl Physiol* 1997; 83: 1256–1269
- [19] Babcock MA, Pegelow DF, Harms CA et al. Effects of respiratory muscle unloading on exercise-induced diaphragm fatigue. *J Appl Physiol* 2002; 93: 201–206
- [20] Shephard RJ, Astrand PO. (ed). *Endurance in Sport*. John Wiley & Sons, 2008
- [21] Aliverti A. The respiratory muscles during exercise. *Breathe (Sheff)* 2016; 12: 165–168
- [22] Sarro KJ, Silvatti AP, Barros RML. Coordination between ribs motion and thoracoabdominal volumes in swimmers during respiratory maneuvers. *J Sports Sci Med* 2008; 7: 195–200
- [23] Massaroni C, Silvatti AP, Levai IK et al. Comparison of marker models for the analysis of the volume variation and thoracoabdominal motion pattern in untrained and trained participants. *J Biomech* 2018; 76: 247–252
- [24] Cavassini CL, Gomes EL, Luiz JG et al. Respiratory and thoracoabdominal motion pattern at rest and after sub-maximum effort in children with asthma. *Curr Res Physiol* 2022; 5: 287–291
- [25] Ferrigno G, Carnevali P, Aliverti A et al. Three-dimensional optical analysis of chest wall motion. *J Appl Physiol* 1994; 77: 1224–1231
- [26] Kibler WB, Press J, Sciascia A. The role of core stability in athletic function. *Sports Med* 2006; 36: 189–198
- [27] Stegmann H, Kindermann W. Comparison of prolonged exercise tests at the individual anaerobic threshold and the fixed anaerobic threshold of 4 mmol.l⁻¹ lactate. *Int J Sports Med* 1982; 3: 105–110
- [28] Fernandez-Garcia B, Pérez-Landaluce J, Rodríguez-Alonso M et al. Intensity of exercise during road race pro-cycling competition. *Med Sci Sports Exerc* 2000; 32: 1002–1006
- [29] Earnest CP, Foster C, Hoyos J et al. Time trial exertion traits of cycling's Grand Tours. *Int J Sports Med* 2009; 30: 240–244
- [30] Zavorsky GS, Murias JM, Gow J et al. Laboratory 20-km cycle time trial reproducibility. *Int J Sports Med* 2007; 28: 743–748
- [31] Rodrigues IM, Pereira ET, Lopes AL et al. Is age rating enough to investigate changes in breathing motion pattern associated with aging of physically active women? *J Biomech* 2021; 125: 110582
- [32] Miller MR, Hankinson J, Brusasco V et al. Standardisation of spirometry. *ERJ Open Res* 2005; 26: 319–338
- [33] Society AT, Society ER. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med* 2002; 166: 518–624
- [34] Rufino R, Costa CH, Lopes AJ et al. Spirometry reference values in the Brazilian population. *Braz J Med Biol Res* 2017; 50: e5700
- [35] Borg GA. Psychophysical bases of perceived exertion. *Med Sci Sports Exerc* 1982; 14: 377–381
- [36] Grimby G, Goldman M, Mead J. Respiratory muscle action inferred from rib cage and abdominal v-p partitioning. *J Appl Physiol* 1976; 41: 739–751
- [37] Johnson BD, Babcock MA, Suman OE et al. Exercise-induced diaphragmatic fatigue in healthy humans. *J Physiol* 1993; 460: 385–405
- [38] Powers SK, Criswell D, Lieu FK et al. Diaphragmatic fiber type specific adaptation to endurance exercise. *Respir Physiol* 1992; 89: 195–207
- [39] Vincent HK, Powers SK, Stewart DJ et al. Short-term exercise training improves diaphragm antioxidant capacity and endurance. *Eur J Appl Physiol* 2000; 81: 67–74
- [40] Lucía A, Hoyos J, Pardo J et al. Metabolic and neuromuscular adaptations to endurance training in professional cyclists: A longitudinal study. *Jpn J Physiol* 2000; 50: 381–388
- [41] Hodges PW, Gandevia SC. Activation of the human diaphragm during a repetitive postural task. *J Physiol* 2000; 522: 165–175
- [42] Deschamps C, Rodarte JR, Wilson TA. Coupling between rib cage and abdominal compartments of the relaxed chest wall. *J Appl Physiol* 1988; 65: 2265–2269
- [43] Iandelli I, Aliverti A, Kayser B et al. Determinants of exercise performance in normal men with externally imposed expiratory flow limitation. *J Appl Physiol* 2002; 92: 1943–1952
- [44] Hodges PW, Sapsford R, Pengel LHM. Postural and respiratory functions of the pelvic floor muscles. *NeuroUrol Urodyn* 2007; 26: 362–371
- [45] Kocjan J, Adamek M, Gzik-Zroska B et al. Network of breathing. Multifunctional role of the diaphragm: A review. *Adv Respir Med* 2017; 85: 224–232
- [46] Frank C, Kobesova A, Kolar P. Dynamic neuromuscular stabilization & sports rehabilitation. *Int J Sports Phys Ther* 2013; 8: 62–73