Detective flow imaging endoscopic ultrasound for localizing pancreatic insulinomas that are undetectable with other imaging modalities





▶ Fig. 1 B-mode endoscopic ultrasound showed a 9-mm pale tumor (arrows) with slightly higher echogenicity than the surrounding area.

Among pancreatic neuroendocrine tumors (PNETs), pancreatic insulinomas are often particularly difficult to detect by imaging studies, making localization diagnosis difficult [1,2]. Recently, the new technique of detective flow imaging endoscopic ultrasound (DFI-EUS), which is capable of displaying minute blood flow in the entire pancreas without contrast medium, has been introduced. We report a pancreatic insulinoma that was difficult to detect with other imaging tests, and for which DFI-EUS was useful for tumor detection.

A 40-year-old woman presented with a chief complaint of dizziness. She had recurrent hypoglycemic attacks. Contrast-enhanced dynamic computed tomography, magnetic resonance imaging, fluorodeoxyglucose-positron emission tomography, and somatostatin receptor scintigraphy did not detect any lesions. EUS was performed using an ultrasound scanning system (ARIETTA 850; FUJIFILM Medical Co., Ltd., Tokyo, Japan) and convex-type endoscope (GF-UCT260; Olympus, Tokyo, Japan). B-mode EUS showed a 9-mm pale tumor with slightly higher



Fig. 2 Detective flow imaging endoscopic ultrasound (DFI-EUS) detected a distinct hypervascular tumor (arrows). Strong blood flow effect was observed on the tumor surface and small vascular blood flow was observed in the interior. **a** B-mode image. **b** DFI-EUS.

echogenicity than the surrounding area (> Fig. 1, > Video 1). DFI-EUS detected a distinct multivessel tumor in the body of the pancreas (> Fig. 2). Contrast-enhanced EUS by Sonazoid (Daiichi-Sankyo, Tokyo, Japan) revealed a hypervascularized tumor (> Fig. 3). EUS-guided fineneedle aspiration was not performed because of the high risk of complications due to the main pancreatic duct interposition in the puncture line; selective arterial secretagogue injection test was positive in the body of the pancreas. The clinical diagnosis was insulinoma and a distal pancreatic body resection was performed. The final pathological diagnosis was PNET G1, insulinoma.

DFI-EUS is a new method for imaging and detecting small vessels and low-velocity blood flow without the use of ultrasound contrast agents. A previous study reported the utility of DFI-EUS for pancreaticobiliary lesions [3]. In this case, DFI-EUS clearly delineated a tumor with more vascularity than contrast EUS and was useful in the diagnosis of insulinoma. DFI-EUS is useful in the evaluation of



Video 1 Imaging findings of B-mode endoscopic ultrasound (EUS), detective flow imaging endoscopic ultrasound (DFI-EUS), and contrast-enhanced EUS in pancreatic insulinoma. DFI-EUS clearly delineated a tumor with more vascularity than contrast EUS.

tumor blood flow in pancreatic insulinomas. While Sonazoid contrast can only adequately observe the field of view at the time of contrast injection, DFI-EUS can screen the entire pancreas. DFI-EUS



▶ Fig. 3 Contrast-enhanced endoscopic ultrasound depicted a mildly hypervascularized tumor (arrows), although the boundary with the surrounding area was somewhat difficult to discern. a B-mode image. b Contrast-enhanced endoscopic ultrasound.

may be useful for screening and localizing the entire pancreas for insulinomas and other PNETs.

Endoscopy_UCTN_Code_CCL_1AF_2AZ

Conflict of Interest

The authors declare that they have no conflict of interest.

The authors

Shinichi Nihei¹, Yusuke Kurita¹, Sho Hasegawa¹, Kunihiro Hosono¹, Noritoshi Kobayashi², Kensuke Kubota¹, Atsushi Nakajima¹

- 1 Department of Gastroenterology and Hepatology, Yokohama City University Hospital, Yokohama, Japan
- 2 Department of Oncology, Yokohama City University Hospital, Yokohama, Japan

Corresponding author

Yusuke Kurita, MD, PhD

Department of Gastroenterology and Hepatology, Yokohama City University Hospital, 3-9 Fukuura, Kanazawa-ku, Yokohama 236-0004, Japan kuritay@yokohama-cu.ac.jp

References

- Khashab MA, Yong E, Lennon AM et al. EUS is still superior to multidetector computerized tomography for detection of pancreatic neuroendocrine tumors. Gastrointest Endosc 2011; 73: 691–696
- [2] Kurita Y, Hara K, Kobayashi N et al. Detection rate of endoscopic ultrasound and computed tomography in diagnosing pancreatic neuroendocrine neoplasms including small lesions: a multicenter study. J Hepatobiliary Pancreat Sci 2022; 29: 950–959
- [3] Yamashita Y, Yoshikawa T, Yamazaki H et al. A novel endoscopic ultrasonography imaging technique for depicting microcirculation in pancreatobiliary lesions without the need for contrast-enhancement: a prospective exploratory study. Diagnostics (Basel) 2021; 11: 2018

Bibliography

Endoscopy 2024; 56: E342–E343 DOI 10.1055/a-2291-9116 ISSN 0013-726X © 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/) Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany



ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



E-Videos is an open access online section of the journal *Endoscopy*, reporting on interesting cases

and new techniques in gastroenterological endoscopy. All papers include a high-quality video and are published with a Creative Commons CC-BY license. Endoscopy E-Videos qualify for HINARI discounts and waivers and eligibility is automatically checked during the submission process. We grant 100% waivers to articles whose corresponding authors are based in Group A countries and 50% waivers to those who are based in Group B countries as classified by Research4Life (see: https:// www.research4life.org/access/eligibility/).

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos