

SYNLETT Spotlight 278

Iodoxybenzene (PhIO₂)

Compiled by Cai Zhang



This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Cai Zhang was born in Anhui Province, P. R. of China. He received his B.Sc. (2005) from Huaibei Coal Industry Teachers College, Anhui, P. R. of China. He is currently pursuing his M.Sc. under the tutelage of Professor Dacheng Yang at Southwest University, Chongqing, P. R. of China. His current research interests include the development of hypervalent iodine for application in organic synthesis.


College of Chemistry and Chemical Engineering, Southwest University, Chongqing 400715, P. R. of China
E-mail: stezh64@swu.edu.cn

Introduction

Iodoxybenzene (PhIO₂, mp 230 °C) is the first organic compound of iodine(V) prepared by Willgerodt 100 years ago.¹ Specifically, it can be prepared by disproportionation of iodosylbenzene under steam distillation.² Ten years ago iodoxybenzene has been used for the synthesis of cadalenquinone,^{3a} unsaturated carbonyl compounds^{3b-c}, oxidienes,^{3d-e}, epoxides,^{3f} quinine imines,^{3g} and sulfones.^{3h} In this Spotlight recent applications using PhIO₂ in organic syntheses are summarized.

Preparation


Iodoxybenzene (PhIO₂) can be directly prepared using commercial 40% peracetic acid or sodium periodate by dissolving iodobenzene in water (Scheme 1).⁴




Scheme 1

Abstracts


(A) Kita et al. show the solubilization and activation of PhIO₂ by adding catalytic amounts of both cetyltrimethylammonium bromide (CTAB) or MgBr₂ and a chiral tartaric acid derivative. Both reagents were indispensable for the enhancement of chemical and optical yields in the oxidation of sulfides to sulfoxides.⁵




(B) The application of PhIO₂ in the oxidation reaction of phosphorous, phosphorothiono and phosphoseleno compounds into the corresponding ≡P(O) analogues was demonstrated by Lopusinski et al.^{6a} The direct conversion of ammonium dialkyl phosphoro-selenoates into the dialkyl phosphoramidates using iodoxybenzene has been developed by the same group.^{6b}




(C) The reaction performed between *N*-sulfinylphosphoramidate and PhIO_2 in the presence of diethylphosphoramidate results in the formation of *N,N*-bis(diethylphosphor)sulfamide.⁷



(D) Catalytic allylic oxidation of alkenes into enones and oxidation of aryl alkyl ketones into the corresponding ketoacids with a fluorous seleninic acid as catalyst in conjunction with PhIO_2 as reoxidant have been developed by Zou and co-worker.⁸



(E) Recently, air-driven conversion of alcohols into carbonyl compounds in water using catalytic amounts of PhIO_2 has been reported by Liu and co-workers.⁹



References

- (1) Willgerodt, C. *Die Organischen Verbindungen mit Mehrwertigem Jod*; Enke Verlag: Stuttgart, **1914**.
- (2) Willgerodt, C. *J. Prakt. Chem.* **1886**, *33*, 154.
- (3) (a) Reddy, N.; Kesavulu, R. a. o.; G, S. K. *Indian J. Chem., Section B: Org. Chem. Incl. Med. Chem.* **1987**, *26B*, 920. (b) Barton, D. H. R.; Crich, D. *Tetrahedron* **1985**, *41*, 4359. (c) Gleiter, R.; Mueller, G. *J. Org. Chem.* **1988**, *53*, 3912. (d) Künzer, H.; Sauer, G.; Wiechert, R. *Tetrahedron* **1989**, *45*, 6409. (e) Iida, T.; Nishida, S.; Chang, F. C.; Niwa, T.; Goto, J.; Nambara, T. *Chem. Pharm. Bull.* **1993**, *41*, 763. (f) Barret, R.; Sabot, J. P.; Pautet, F.; Cerf, P.; Daudon, M. *Oxidation Communications* **1989**, *12*, 55. (g) Barret, R.; Daudon, M. **1990**, *20*, 1543. (h) Barret, R.; Pautet, F.; Bordat, P.; Tinland, B.; Daudon, M. *Phosphorus, Sulfur and Silicon and the Related Elements* **1989**, *45*, 31.
- (4) (a) Shrefkin, J. G.; Saltzman, H. *Org. Synth.* **1963**, *43*, 65. (b) Kazmierczak, P.; Skulski, L.; Kraszkiewicz, L. *Molecules* **2001**, *6*, 881.
- (5) (a) Tohma, H.; Takizawa, S.; Watanabe, H.; Fukuoka, Y.; Maegawa, T.; Kita, Y. *J. Org. Chem.* **1999**, *64*, 3519. (b) Tohma, H.; Takizawa, S.; Morioka, H.; Maegawa, T.; Kita, Y. *Chem. Pharm. Bull.* **2000**, *48*, 445.
- (6) (a) Mielniczak, G.; Lopusinski, A. *Synlett* **2001**, 505. (b) Mielniczak, G.; Lopusinski, A. *Heteroatom Chem.* **2003**, *14*, 121.
- (7) Mielniczak, G.; Lopusinski, A. *Heteroatom Chem.* **2008**, *19*, 530.
- (8) (a) Crich, D.; Zou, Y. *Org. Lett.* **2004**, *6*, 775. (b) Crich, D.; Zou, Y. *J. Org. Chem.* **2005**, *70*, 3309.
- (9) Mu, R. Z.; Liu, Z. Q.; Yang, Z. J.; Liu, Z. G.; Wu, L. M.; Liu, Z. L. *Adv. Synth. Catal.* **2005**, *347*, 1333.