

# SYNLETT Spotlight 284

## Potassium Thiocyanate (KSCN): A Versatile Reagent

Compiled by Soheil Sayyahi

This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Soheil Sayyahi was born in Ahwaz, Iran in 1975. He received his Bachelor and M.Sc. degrees from Shahid Chamran University of Ahwaz, where he is currently carrying out his Ph.D. under the supervision of Prof. Kiasat and Prof. Badri. His area of research includes the application of polymer-supported reagents and catalysts in organic synthesis.

Chemistry Department, College of Science, Chamran University,  
Ahvaz 61357-4-3169, Iran  
E-mail: sayyahi.soheil@gmail.com




### Introduction

Potassium thiocyanate (KSCN) is a white odorless, crystalline powder, slightly hygroscopic, and commercially available reagent. It is readily soluble in water and stable under normal temperature and pressure (mp: 173 °C,  $d = 1.89 \text{ g/cm}^3$ ). Since sulfur-containing groups serve as an important auxiliary function in synthetic sequences,<sup>1</sup> potassium thiocyanate is widely used as a transfer reagent for sulfur in various organic transformations.<sup>2</sup> The hypervalent iodine(III) in combination with potassium thiocyanate and diphenyl diselenide promoted a multicomponent reaction for the synthesis of phenylsele-


nyl thiocyanates and isothiocyanates from alkenes.<sup>3</sup> Various tosyl and bromo derivatives of Cbz-, Boc-, and Fmoc-protected threonine methyl esters have been subjected to nucleophilic substitution with potassium thiocyanate in acetonitrile for the synthesis of allo- and threo-3,3'-dimethylcystine derivatives.<sup>4</sup> This reagent is supported on silica gel and applied for thiocyanation of  $\beta$ -dicarbonyl compounds and the synthesis of 2-aminothiazoles.<sup>5</sup> Recently, potassium thiocyanate is used for the conversion of alkyl halides into alkyl thiocyanate in water under phase-transfer catalysis.<sup>6</sup> It is also employed for the synthesis of 1-aryloxy-3-(substituted-2-benzothiazolyl)thioureas with antibacterial properties.<sup>7</sup>

### Abstracts


(A) Das et al.<sup>8</sup> reported an efficient and catalyst-free procedure for the synthesis of thiiranes from oxiranes by treatment with KSCN using PEG as a reaction medium at room temperature.





(B) In the presence of a catalytic amount of  $\text{LiClO}_4$ , oxiranes are converted into the corresponding thiiranes by potassium thiocyanate in nonaqueous condition.<sup>9</sup>




(C) Aoyama et al.<sup>10a</sup> introduced a supported reagent system, KSCN/ $\text{SiO}_2$  and  $\text{BnNH}_3\text{OAc}/\text{Al}_2\text{O}_3$ , that has been employed in a one-pot synthesis of *N*-allylthioureas. Also, various  $\alpha$ -halo ketones and allylic bromides were converted into 2-aminothiazoles and *N*-allylthioureas from commercially available materials in one pot by using the supported reagents.<sup>10b</sup>




(D) Rao et al.<sup>11</sup> reported a mild and efficient method for the regio-selective ring opening of aziridines with KSCN in the presence of  $\beta$ -CD as catalyst and water at room temperature with excellent yields ranging from 78 to 90%.




(E) Recently, a practical synthesis of 1,3-disubstituted imidazole-2-thiones via a microwave-promoted reaction of imidazolium salts with potassium thiocyanate or potassium thioacetate under solvent-free conditions has been developed.<sup>12</sup>



(F) Ring-opening reactions of various *N*-tosylaziridines with KSCN proceeded in a silica-water reaction medium in good yield with complete regioselectivity. The system is applicable to a ring expansion of an aziridine with potassium thiocyanate leading to a thiazolidine derivative.<sup>13</sup>



(G) The microwave-assisted nucleophilic substitution of potassium thiocyanate with different halides or tosylates in water has been developed and proved very successful with the formation of alkylthiocyanates in good to excellent yields. Noteworthy, no significant rearrangement to isothiocyanates was observed.<sup>14</sup>



## References

- (1) Kondo, T.; Mitsudo, T. A. *Chem. Rev.* **2000**, *100*, 3205.
- (2) Erian, A. W.; Sherif, S. M. *Tetrahedron* **1999**, *55*, 7957.
- (3) Margarita, R.; Mercanti, C.; Parlanti, L.; Piancatelli, G. *Eur. J. Org. Chem.* **2000**, 1865.
- (4) Nasir Baig, R. B.; Sai Sudhir, V.; Chandrasekaran, S. *Tetrahedron: Asymmetry* **2008**, *19*, 1425.
- (5) Kodomari, M.; Aoyama, T.; Suzuki, Y. *Tetrahedron Lett.* **2002**, *43*, 1717.
- (6) Kiasat, A. R.; Badri, R.; Sayyahi, S. *Chin. Chem. Lett.* **2008**, *19*, 1301.
- (7) Saeed, A.; Rafique, H.; Hameed, A.; Rasheed, S. *Pharm. Chem. J.* **2008**, *42*, 191.
- (8) Das, B.; Saidi Reddy, V.; Krishnaiah, M. *Tetrahedron Lett.* **2006**, *47*, 8471.
- (9) Reddy, S.; Nagavani, S. *Heteroatom Chem.* **2008**, *19*, 97.
- (10) (a) Aoyama, T.; Murata, S.; Nagata, Y.; Takidoa, T.; Kodomari, M. *Tetrahedron Lett.* **2005**, *46*, 4875.  
(b) Aoyama, T.; Murata, S.; Arai, I.; Araki, N.; Takidoa, T.; Suzuki, Y.; Kodomari, M. *Tetrahedron* **2006**, *62*, 3201..
- (11) Reddy, S. M.; Narender, M.; Nageswar, Y. V. D.; Rao, K. R. *Tetrahedron Lett.* **2005**, *46*, 6437.
- (12) Tao, X. L.; Lei, M.; Wang, Y. G. *Synthetic Commun.* **2007**, *37*, 399.
- (13) Minakata, S.; Hotta, T.; Oderaotoshi, Y.; Komatsu, M. *J. Org. Chem.* **2006**, *71*, 7471.
- (14) Ju, Y.; Kumar, D.; Varma, R. S. *J. Org. Chem.* **2006**, *71*, 6697.