
T. NISHIMURA,* Y. YASUHARA, T. SAWANO, T. HAYASHI* (KYOTO UNIVERSITY, JAPAN) Iridium/Chiral Diene Catalyzed Asymmetric 1,6-Addition of Arylboroxines to $\alpha, \beta, \gamma, \delta$ -Unsaturated Carbonyl Compounds

J. Am. Chem. Soc. 2010, 132, 7872-7873.

Iridium-Catalyzed Enantioselective 1,6-Addition



Ir/chiral diene catalyzed asymmetric 1,6-addition to dienones:

Ir/chiral diene catalyzed asymmetric 1,6-addition to dienamides and dienoate:

Synthetic application:

Significance: This work described an iridium-catalyzed asymmetric 1,6-addition of arylboroxines to $\alpha,\beta,\gamma,\delta$ -unsaturated carbonyl compounds by employing an iridium chiral diene complex. Various δ -arylated carbonyl compounds were obtained in high yields and enantioselectivity. Also, the authors elaborated the product in a short sequence to provide nature product and important building blocks.

Comment: Asymmetric 1,6-Michael additions to acyclic systems are challenging as a result of poor control of regioselectivity and enantioselectivity. Recently, the authors reported the hydroxoiridium complex coordinated with COD< was able to catalyze the 1,6-addition of arylboronic acids to unsaturated carbonyl compounds (*Angew. Chem. Int. Ed.* **2006**, *45*, 5164). In this work, they developed an asymmetric version of this reaction. Excellent control of enantioselectivity and regioselectivity was achieved.

 SYNFACTS Contributors: Hisashi Yamamoto, Jiajing Tan

 Synfacts 2010, 9, 1019-1019
 Published online: 23.08.2010

 DOI: 10.1055/s-0030-1257993; Reg-No.: H09710SF

Category

Metal-Catalyzed Asymmetric Synthesis and Stereoselective Reactions

Key words

chiral diene ligands 1,6-addition

iridium

