

Erratum

Cucurbitane-Type Triterpenoids from *Momordica charantia*

Jun Ma, Paul Whittaker, Amy C. Keller, Eugene P. Mazzola, Rahul S. Pawar, Kevin D. White, John H. Callahan, Edward J. Kennelly, Alexander J. Krynitsky, Jeanne I. Rader
 Planta Med 2010; 76: 1758–1761

In the paper by Ma et al., Cucurbitane-type triterpenoids from *Momordica charantia*. *Planta Med* 2010; 76: 1758–1761 [1], compound **1** was named momordicoside U (Fig. 1). Unbeknown to the authors, the same name had been given earlier to a different substance in the article by Nguyen et al., Cucurbitane-type triterpene glycosides from the fruits of *Momordica charantia*. *Magn Reson Chem* 2010; 48: 392–396 [2] (compound **1**) (Fig. 2). Therefore the compound isolated by Ma et al. is to be renamed “Momordicoside X”.

References

- 1 Ma J, Whittaker P, Keller AC, Mazzola EP, Pawar RS, White KD, Callahan JH, Kennelly EJ, Krynitsky AJ, Rader JI. Cucurbitane-type triterpenoids from *Momordica charantia*. *Planta Med* 2010; 76: 1758–1761
- 2 Nguyen XN, Phan VK, Chau VM, Ninh KB, Nguyen XC, Le MH, Bui HT, Tran HQ, Nguyen HT, Kim YH. Cucurbitane-type triterpene glycosides from the fruits of *Momordica charantia*. *Magn Reson Chem* 2010; 48: 392–396




Fig. 1 Structure of momordicoside X (compound **1**) by Ma et al. [1].




Fig. 2 Structure of momordicoside U (compound **1**) by Nguyen et al. [2].