


SYNFORM

People, Trends and Views in Synthetic Organic Chemistry

2012/05

SYNSTORIES

■ **A Highly Diastereo- and Enantioselective Reaction for Constructing Functionalized Cyclohexanes: Six Contiguous Stereocenters in One Step**



■ **Radical Addition of Alkyl Halides to Formaldehyde in the Presence of Cyanoborohydride as a Radical Mediator**

■ **SYNTHESIS/SYNLETT**
Advisory Board Focus: Professor Keisuke Suzuki (Tokyo Institute of Technology, Japan)

CONTACT +++++

Your opinion about SYNFORM is welcome, please correspond if you like:
marketing@thieme-chemistry.com



Dear readers,

This issue of **SYNFORM** is dominated by Asian chemistry and chemists, which is not at all surprising given the impressive development of countries like P. R. of China, India, Singapore, South Korea and others which are now

competing with Japan on the continental scale and synergistically acting with Japan in establishing Asia as the new giant in the global research arena. In the first **SYNSTORY** Professor I. Ryu (Japan) guides us through his recent radical process based on the addition of alkyl halides to an underutilized, at least in radical chemistry, one-carbon building block such as formaldehyde. The second **SYNSTORY** brings us to the P. R. of China where Professor H. Huang has recently established a new world record in the simultaneous formation of six new contiguous stereocenters in one molecule, a highly functionalized cyclohexane, by means of a single synthetic step. Last but not least, the Editorial Advisory Board profile features Professor K. Suzuki (Japan).

Enjoy your reading!

Matteo Zanda


Editor of SYNFORM

IN THIS ISSUE

SYNSTORIES ■ ■ ■ ■

Radical Addition of Alkyl Halides to Formaldehyde in the Presence of Cyanoborohydride as a Radical Mediator..... **A43**

A Highly Diastereo- and Enantioselective Reaction for Constructing Functionalized Cyclohexanes: Six Contiguous Stereocenters in One Step..... **A46**



SYNTHESIS/SYNLETT Advisory Board Focus: Professor Keisuke Suzuki (Tokyo Institute of Technology, Japan)..... **A49**

COMING SOON **A50**

CONTACT +++++

If you have any questions or wish to send feedback, please write to Matteo Zanda at:
Synform@chem.polimi.it

NEWS AND VIEWS ■ ■ NEWS AND VIEWS ■ ■ NEWS AND VIEWS ■ ■


Radical Addition of Alkyl Halides to Formaldehyde in the Presence of Cyanoborohydride as a Radical Mediator

J. Am. Chem. Soc. **2012**, *134*, 875–877

Whereas carbon monoxide, isonitriles, and substituted oxime ethers have found wide applications in radical-mediated multi-component reactions as viable C1 synthons,¹ the utilization of formaldehyde has been only marginally explored.² This is in sharp contrast with the fact that formaldehyde is a quite common reagent in carbanion chemistry, and has been successfully used for one-carbon homologation reactions. Recently, however, the group of Professor Ilhyong Ryu from Osaka Prefecture University (Japan) developed an efficient radical process for the hydroxymethylation of haloalkanes, in which formaldehyde is used as C1 radical synthon. The break-

through is concerned with the use of tetrabutylammonium cyanoborohydride as a radical mediator.

“It has long been known that radical addition to formaldehyde does occur. We thought that formaldehyde is a cheap and potentially abundant feedstock and it was time to develop synthetically useful radical reactions making use of formaldehyde,” said Professor Ryu, who acknowledged that the original work of Fuller and Rust, published in 1958, disclosed that the *tert*-butyl peroxide initiated radical addition of cyclohexane to formaldehyde takes place, affording cyclohexanemethanol.^{2a} Another key paper by Brown, Suzuki, and co-workers, pub-



lished in 1972, demonstrated that in the presence of air, the reaction of tributylborane and formaldehyde gave 1-pentanol.^{2b} “After these pioneering efforts, surprisingly this chemistry has been neglected for nearly four decades,” pointed out Professor Ryu. “Previously, we demonstrated that tetrabutylammonium hydroborate can be used for Giese-type radical reactions and radical carbonylations as a substitute for tributyltin hydride.³ We were naturally tempted to apply formaldehyde in combination with borohydride mediators,” said Professor Ryu. Next, the Ryu group reported that the hydroxymethylation of iodoalkanes proceeds using CO and tetrabutylammonium borohydride;⁴ therefore, this reagent was chosen for the radical reaction of formaldehyde. According to Professor Ryu, the first experiment using tetrabutylammonium borohydride was totally unsuccessful (Scheme, eq. 1). “The key to success was the employment of ‘low-reactive’ borohydride reagents for the hydride addition to formaldehyde, which Takuji Kawamoto, a PhD student in our group, keenly designed and optimized. Takuji examined a milder reagent, tetrabutylammonium cyanoborohydride, which worked beautifully,” said Professor Ryu. Indeed, the reaction of 1-iodo-adamantane gave 85% yield of 1-hydroxymethyladamantane (Scheme, eq. 2). “The reaction of cholesteryl bromide gave a set of products via tandem radical reactions, which is a good sign that further applications to cascade reactions would be promising (Scheme, eq. 3),” added Professor Ryu.

“It is interesting to note that, besides borohydride reagents, radical reactions using borane reagents such as NHC-borane are now widely investigated,” he continued.⁵ “We are quite interested in the reactions of formaldehyde in combination with such newly evolving NHC-borane reagents,” said Professor Ryu, who pointed out that executing kinetic studies surrounding alkyl radical addition to formaldehyde and abstraction of hydrogen of the resulting alkoxy radicals from borate anions would be strongly needed in order to accelerate studies focusing on multicomponent reactions. “In the near future, this methodology will be extended to multicomponent reactions that incorporate formaldehyde and other carbon units, such as alkenes and alkynes. Thus, radical chemistry of formaldehyde will open a new chapter,” concluded Professor Ryu. ■

Matteo Zanda

REFERENCES

- (1) (a) I. Ryu, N. Sonoda, D. P. Curran *Chem. Rev.* **1996**, *96*, 177; (b) S. Kim, S. Kim *Bull. Chem. Soc. Jpn.* **2007**, *80*, 809; (c) H. Miyabe, H. Yoshioka, S. Kohtani *Curr. Org. Chem.* **2010**, *14*, 1254.
- (2) (a) G. Fuller, F. F. Rust *J. Am. Chem. Soc.* **1958**, *80*, 6148; (b) N. Miyaura, M. Itoh, A. Suzuki, H. C. Brown, M. M. Midland, P. Jacob III *J. Am. Chem. Soc.* **1972**, *94*, 6549.
- (3) I. Ryu, S. Uehara, H. Hirao, T. Fukuyama *Org. Lett.* **2008**, *10*, 1005.
- (4) S. Kobayashi, T. Kawamoto, S. Uehara, T. Fukuyama, I. Ryu *Org. Lett.* **2010**, *12*, 1548.
- (5) Recent review: D. P. Curran, A. Solov'yev, M. M. Brahmī, L. Fensterbank, M. Malacria, E. Lacôte *Angew. Chem. Int. Ed.* **2011**, *50*, 10294.

About the authors



T. Kawamoto

Takuji Kawamoto was born in Sakai, Osaka (Japan). He received his Masters degree from Osaka Prefecture University (Japan) in 2011. He is currently a PhD student at the Osaka Prefecture University (Professor Ryu is the supervisor) and will undertake a JSPS (Japan Society of Promotion of Science) Research Fellowship this April.



Dr. T. Fukuyama

Takahide Fukuyama received his Ph.D. in 1999 from Osaka University (Japan) under the direction of Professor Shinji Murai. He spent the year 1999–2000 as a Post-doctoral Fellow of JSPS at Okayama University of Science (Japan) with Professor Junzo Otera. In 2000, he was appointed as an Assistant Professor in Professor Ryu's group at Osaka Prefecture University. He was promoted to Lecturer in 2007 and to Associate Professor in 2010. He had the experience of working with Max Malacria at the University of Pierre and Marie Curie (France, 2006). He is the recipient of the Dainippon Ink & Chemicals Inc. Award of the Society of Synthetic Organic Chemistry, Japan.



Prof. I. Ryu

Ilhyong Ryu received his Ph.D. from Osaka University in 1978 under the direction of Professor Noboru Sonoda. After serving as a JSPS Postdoctoral Fellow and a Research Associate at Osaka University, he was appointed Assistant Professor at Osaka University in 1988 and promoted to Associate Professor in 1995. He also had the experience of working with Professor Howard Alper (1991–1992) at the University of Ottawa (Canada). In

2000, he moved to Osaka Prefecture University as a Full Professor. He has been the recipient of many awards, including the Progress Award in Synthetic Organic Chemistry, Japan (1990) and the Chemical Society of Japan Award for Creative Work (2004).

domino processes that could generate many more stereocenters in one step,” he continued. “Chiral functionalized cyclohexanes with contiguous multiple stereocenters are a highly desirable structural motif found in natural and synthetic bioactive compounds. Previously, the highest number of contiguous stereocenters generated in one single step in functionalized cyclohexanes was five, as reported by Karl Anker Jørgensen (*Angew. Chem. Int. Ed.* **2007**, *46*, 9202) and Bor-Cherng Hong (*Org. Lett.* **2011**, *13*, 1278). A cyclohexane with six contiguous stereocenters was reported as a byproduct by Mikiko Sodeoka and colleagues (*J. Am. Chem. Soc.* **2010**, *132*, 4036; *Tetrahedron: Asymmetry*, **2010**, *21*, 1682) in their Ni-catalyzed conjugate addition reaction, while Jean Rodriguez and co-workers also synthesized only one cyclohexane compound with six chiral centers in lower yield using an organocatalytic method (*Org. Lett.* **2010**, *12*, 5246).” With the background above, Professor Huang and his co-workers designed a cascade process (Figure 1) which was triggered by an active enolate species generated from the α -keto ester. “The principal challenge for establishing such a domino catalytic asymmetric process was to identify a good catalyst for facilitating the formation of the metal-enolate active species **A** and constructing a well-oriented chiral environment around the enolate intermediate (Figure 1),” he explained.

Hypothesizing that the conformationally rigid structure would have good chiral induction abilities, Professor Huang and his co-workers designed and synthesized a series of rigid chiral diamines (*Org. Lett.* **2009**, *11*, 4536) with an intriguing structural motif. “The design of these chiral diamines is based on the structure of DPEN,” he said. “Although DPEN has already been proven to be a good ligand for asymmetric reactions, the free rotation of the phenyl group on the backbone renders it conformationally flexible. We reasoned that

the conformational mobility can be restricted by introducing two additional methylene groups on its scaffold to bridge the phenyl and amino groups. Thus, the chiral amines **5** and **6** were designed (Figure 2),” said Professor Huang, who pointed out that these chiral diamines have subsequently been developed into excellent catalysts for the asymmetric hydrogenation of ketones (*Chem. Eur. J.* **2011**, *17*, 7760).

“Inspired by these results, Dengjian Shi, a third-year graduate student in our group, had the idea to try to prepare a series of Lewis acid catalysts with these chiral diamines for promotion of the above-described enolate-triggered tandem reaction,” acknowledged Professor Huang. “First of all, the catalyst should be highly chemoselective for overcoming the major side reaction that leads to the undesired linear conjugate addition byproduct. Secondly, the catalyst should be highly stereoselective for enantioselectively producing only one enantiomer out of 64 (2^6) possible stereoisomers, since six stereocenters are generated in this cascade asymmetric reaction,” he said. After extensive screening of a number of experimental factors, the Chinese researchers finally found that the catalyst composed of $\text{Cu}(\text{OAc})_2 \cdot \text{H}_2\text{O}$ and chiral diamine **5a** was able to afford the target cyclohexane carboxylates with excellent enantioselectivity and complete diastereoselectivity under mild and environmentally friendly reaction conditions. “The reaction could be performed with 0.1 mol% of Cu catalyst on a gram-scale of substrate to afford the desired product with high yield and excellent selectivity,” he added. “From a practical point of view, we believe that our method represents a record for the synthesis of the highest number of contiguous stereocenters in a molecule made in just one highly selective step,” Professor Huang pointed out. “Although a method for generating chiral compounds with eight contiguous stereocenters has been reported previously, two steps were needed to




Figure 2

perform it (*Angew. Chem. Int. Ed.* **2007**, *46*, 467),” he said. “This research has advanced the state-of-the-art of asymmetric catalytic tandem reactions and, furthermore, confirms that the asymmetric tandem reaction is a powerful and efficient tool in synthetic organic chemistry. It is our opinion that con-

tinuing and extensive research efforts to merge different reaction modes into one domino process will become mainstream in asymmetric synthesis,” concluded Professor Huang. ■

Matteo Zanda

About the authors



Prof. H. Huang



Prof. C. Xia



D. Shi



H. Zhou



Y. Xie

SYNTHESIS/SYNLETT Advisory Board Focus: Professor Keisuke Suzuki (Tokyo Institute of Technology, Japan)

■ **Background and Purpose.** *SYNFORM* will from time to time portrait *SYNTHESIS/SYNLETT* Advisory Board members who answer several questions regarding their research interests and revealing their impressions and views on the developments in organic chemistry as a general research field. In this issue, we present Professor Keisuke Suzuki, Tokyo Institute of Technology (Japan).

BIOGRAPHICAL SKETCH



Prof. K. Suzuki

Keisuke Suzuki was born in 1954 in Chigasaki (Japan). He obtained a Doctor of Science Degree in 1983 at The University of Tokyo (Japan) under the guidance of Professor Teruaki Mukaiyama. He was appointed Research Associate to the late Professor Gen-ichi Tsuchihashi in the Department of Chemistry, Keio University (Japan, 1983), and promoted to Lecturer (1987), Associate Professor

(1989), and Professor (1994) at Keio University. He enjoyed a year as a Visiting Professor at ETH Zürich (Switzerland, 1990–1991) with Professor Dieter Seebach. In 1996, he moved to the Department of Chemistry, Tokyo Institute of Technology, where he remains today. The awards he has received include The Chemical Society of Japan (CSJ) Award for Young Chemists in 1986, the Takasago Award of the Synthetic Organic Chemical Society, Japan (SOCSJ) in 1994, the Japan IBM Award in 1994, the Teshima Award in 1997, the Nagoya Silver Medal in 1999, the SOCSJ Award in 2003, Prizes for Science and Technology, Ministry of Education, Culture, Sport, Science and Technology, Japan in 2006, The CSJ Award in 2008, the Humboldt Research Prize in 2009, and the Medal with Purple Ribbon in 2010.

INTERVIEW

SYNFORM | *What are your main current research interests?*

K. Suzuki | Organic synthesis, focusing the attention to new strategies and tactics in natural/unnatural product synthesis. In spite of the great advances in organic synthesis, certain classes of compounds still remain difficult to access. An example is the densely functionalized polycyclic compounds derived from the type-II polyketide biosynthesis. The synthesis becomes even more challenging when the structure is hybridized with other biosynthetic products, for example, sugars or terpenoids, which we call *hybrid natural products* (Review: K. Suzuki, Lessons from Total Synthesis of Hybrid Natural Products, *Chem. Rec.* **2010**, *10*, 291). Other current targets include the catechin-class polyphenols and their oligomers and the pluramycin-class bis-*C*-glycoside antibiotics.

SYNFORM | *What is your most important scientific achievement to date and why?*

K. Suzuki | The total synthesis of mycinamicin IV, a macrocyclic antibiotic (T. Matsumoto, H. Maeta, K. Suzuki, G. Tsuchihashi, First Total Synthesis of Mycinamicin IV and VII. Successful Application of New Glycosidation Reaction, *Tetrahedron Lett.* **1988**, *29*, 3575). It was my early contribution to the area of natural product synthesis which was completed by using original tactics, including acyclic stereocontrol via stereospecific 1,2-shift for assembling the aglycon and the hafnocene-based *O*-glycoside synthesis. It was also important in my research career because it was through the struggle that I came to notice the importance and synthetic challenge posed by *hybrid natural products*.

SYNFORM | *Do you have hobbies besides chemistry?*

K. Suzuki | Soccer. I used to play in much earlier days, but nowadays I'm just watching games, of course. ■

Matteo Zanda

COMING SOON ► ► COMING SOON ► ►

SYNFORM 2012/06

is available from
May 16, 2012

In the next issues:

SYNSTORIES ■ ■ ■ ■ ■

■ Ruthenium-Catalyzed C–H Bond Arylations of Arenes Bearing Removable Directing Groups via Six-Membered Ruthenacycles

(Focus on an article from the current literature)■ Diastereoselective Rhodium-Catalyzed Ene-Cycloisomerization Reactions of Alkenylidenecyclopropanes: Total Synthesis of (–)- α -Kainic Acid*(Focus on an article from the current literature)*

FURTHER HIGHLIGHTS + + + +

SYNTHESIS

Review on: Detour and Direct Induction of 'Methyl'-Containing Chiral Centers via Catalytic C–H or C–C Bond Formation

(by K. Endo, T. Shibata)

SYNLETT

Account on: Asymmetric Total Synthesis of the Epithilone Sagopilone – From Research to Development

(by U. Klar, J. Platzeck)

SYNFACTS

Synfact of the Month in category "Metal-Catalyzed Asymmetric Synthesis and Stereoselective Reactions":
[Asymmetric Synthesis of Indolines by Pd-Catalyzed C\(sp³\)-H Activation](#)

CONTACT + + + +

Matteo Zanda,
NRP Chair in Medical Technologies
Institute of Medical Sciences
University of Aberdeen
Foresterhill, Aberdeen, AB25 2ZD, UK
andC.N.R. – Istituto di Chimica del Riconoscimento Molecolare,
Via Mancinelli, 7, 20131 Milano, Italy,
e-mail: Synform@chem.polimi.it, fax: +39 02 23993080

Editor

Matteo Zanda, NRP Chair in Medical Technologies, Institute of Medical Sciences, University of Aberdeen, Foresterhill, Aberdeen, AB25 2ZD, UK and
C.N.R. – Istituto di Chimica del Riconoscimento Molecolare
Via Mancinelli, 7, 20131 Milano, Italy
Synform@chem.polimi.it
Fax: +39 02 23993080

Editorial Office

- Managing Editor: Susanne Haak, susanne.haak@thieme.de, phone: +49 711 8931 786
- Scientific Editor: Selena Boothroyd, selena.boothroyd@thieme.de
- Scientific Editor: Stefanie Baumann, stefanie.baumann@thieme.de, phone: +49 711 8931 776
- Assistant Scientific Editor: Christiane Kemper, christiane.kemper@thieme.de, phone: +49 711 8931 768
- Senior Production Editor: Thomas Loop, thomas.loop@thieme.de, phone: +49 711 8931 778
- Production Editor: Helene Deufel, helene.deufel@thieme.de, phone: +49 711 8931 929
- Production Editor: Thorsten Schön, thorsten.schoen@thieme.de, phone: +49 711 8931 781
- Editorial Assistant: Sabine Heller, sabine.heller@thieme.de, phone: +49 711 8931 744
- Marketing: Julia Stötzner, julia.stoetznern@thieme.de, phone: +49 711 8931 771
- Postal Address: SYNTHESIS/SYNLETT/SYNFACTS, Editorial Office, Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany, phone: +49 711 8931 744, fax: +49 711 8931 777
- Homepage: www.thieme-chemistry.com

Publication Information

SYNFORM will be published 12 times in 2012 by Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany, and is an additional online service for SYNTHESIS, SYNLETT and SYNFACTS.

Publication Policy

Product names which are in fact registered trademarks may not have been specifically designated as such in every case. Thus, in those cases where a product has been referred to by its registered trademark it cannot be concluded that the name used is public domain. The same applies as regards patents or registered designs.

Ordering Information for Print Subscriptions to SYNTHESIS, SYNLETT and SYNFACTS

The Americas: Thieme Publishers New York, Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA.

To order: customerservice@thieme.com or use the Web site facilities at www.thieme-chemistry.com, phone: +1 212 760 0888
Order toll-free within the USA: +1 800 782 3488
Fax: +1 212 947 1112

Airfreight and mailing in the USA by Publications Expeditors Inc., 200 Meacham Ave., Elmont NY 11003. Periodicals postage paid at Jamaica NY 11431.

Europe, Africa, Asia, and Australia: Thieme Publishers Stuttgart, Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany.

To order: customerservice@thieme.de or use the Web site facilities at www.thieme-chemistry.com.
Phone: +49 711 8931 421; Fax: +49 711 8931 410Current list prices are available through www.thieme-chemistry.com.

Online Access via Thieme-connect

The online versions of SYNFORM as well SYNTHESIS, SYNLETT and SYNFACTS are available through Thieme-connect (www.thieme-connect.com/ejournals) where you may also register for free trial accounts. For information on multi-site licenses and pricing for corporate customers as well as backfiles please contact our regional offices:The Americas: esales@thieme.com, phone: +1 212 584 4695Europe, Africa, Asia, and Australia: cproducts@thieme.de, phone: +49 711 8931 407

Manuscript Submission to SYNTHESIS and SYNLETT

Please consult the Instructions for Authors before compiling a new manuscript. The current version and the Word template for manuscript preparation are available for download at www.thieme-chemistry.com. Use of the Word template helps to speed up the refereeing and production process.

Copyright

This publication, including all individual contributions and illustrations published therein, is legally protected by copyright for the duration of the copyright period. Any use, exploitation or commercialization outside the narrow limits set by copyright legislation, without the publisher's consent, is illegal and liable to criminal prosecution. This applies translating, copying and reproduction in printed or electronic media forms (databases, online network systems, Internet, broadcasting, telecasting, CD-ROM, hard disk storage, microcopy edition, photomechanical and other reproduction methods) as well as making the material accessible to users of such media (e.g., as online or offline backfiles).

Copyright Permission for Users in the USA

Authorization to photocopy items for internal or personal use, or the internal or personal use of specific clients, is granted by Georg Thieme Verlag KG Stuttgart · New York for libraries and other users registered with the Copyright Clearance Center (CCC) Transactional Reporting Service, provided that the base fee of US\$ 25.00 per copy of each article is paid directly to CCC, 22 Rosewood Drive, Danvers, MA 01923, USA, 0341-0501/02.