

Severity of Chronic Obstructive Pulmonary Disease and Its Relationship to Lung Cancer Prognosis after Surgical Resection*

Yasuo Sekine¹ Hidemi Suzuki¹ Yoshito Yamada² Eitetsu Koh¹ Ichiro Yoshino²

¹Department of Thoracic Surgery, Tokyo Women's Medical University, Yachiyo Medical Center, Yachiyo, Japan

²Department of Thoracic Surgery, Graduate School of Medicine, Chiba University, Chiba, Japan

Address for correspondence Yasuo Sekine, MD, PhD, Department of Thoracic Surgery, Tokyo Women's Medical University, Yachiyo Medical Center, 477-96 OwadaShinden, Yachiyo, Chiba 276-8524, Japan (e-mail: ysekine@tymc.twmu.ac.jp).

Thorac Cardiovasc Surg 2013;61:124–130.

Abstract

Objective The purpose was to determine the rates of postoperative pulmonary complications, and to clarify the impact of COPD on long-term survival in lung cancer patients after surgical resection.

Methods A retrospective chart review was performed on 1,461 patients who had undergone pulmonary resection for lung cancer from 1990 to 2005. Classification of COPD severity was based on spirometric guidelines of the Global Initiative for Chronic Obstructive Lung Disease (GOLD). Postoperative complication rates among the four COPD groups were compared and long-term overall and disease-specific survivals were analyzed.

Results The frequencies of all pulmonary complications in three COPD groups were higher than in the non-COPD group (all $p < 0.05$). Overall and disease-specific survivals were significantly worse in relation to higher COPD grades (all $p \leq 0.05$). Significant prognostic factors were age, body mass index, positive smoking history, tumor size, pneumonectomy, pathologic stage, and COPD grade ($p < 0.05$).

Conclusion Higher COPD grades had higher rates of postoperative pulmonary complications and poorer long-term survivals because of higher rates of cancer-related deaths.

Keywords

- ▶ lung cancer
- ▶ chronic obstructive pulmonary disease
- ▶ prognosis
- ▶ surgery

Introduction

Lung cancer and chronic obstructive pulmonary disease (COPD) are common fatal diseases. Patients with lung cancer who also have COPD are frequently deemed inoperable because of low cardiopulmonary reserve. Preoperative evaluation and risk stratification of patients undergoing lung resection include pulmonary function testing, arterial blood

gas analysis, exercise testing, and split perfusion-ventilation radionuclide scanning.^{1,2}

Pulmonary complications after lung cancer surgery are major causes of morbidity for patients with COPD.^{3,4} Pulmonary resection decreases the already limited respiratory reserve and causes hypoventilation, hypoxia, hypercapnia, and retention of secretions, which can lead to respiratory failure,⁵ and poor short-term survival and quality of life (QOL). However, the effects of different degrees of COPD severity on short- and long-term survival have not been well studied. In particular, the risk of lung cancer recurrence in patients with COPD has been insufficiently studied.

*This paper was presented at the 54th Annual Meeting of the Thomas L. Petty Aspen Lung Conference in Colorado, USA, 2011.

received
September 2, 2011
accepted after revision
November 21, 2011
published online
April 25, 2012

© 2013 Georg Thieme Verlag KG
Stuttgart · New York

DOI <http://dx.doi.org/10.1055/s-0032-1304543>.
ISSN 0171-6425.

Studies evaluating risk factors for postoperative complications and survival in patients with COPD have been published.^{5,6} However, the prevalences of different types of complications and their association with long-term survival of lung cancer patients with COPD have not been determined, especially for patients with postoperative pulmonary complications, in whom compromised lung function affects survival as well as quality of life.

The purpose of this study was to: (a) determine the prevalences of various types of postoperative complications in nonsmall cell lung cancer patients undergoing curative surgery in the various stages of COPD; (b) assess the impact of COPD on long-term survival of these patients; and (c) enumerate the causes of death in our patient population.

Patients and Methods

The study was approved by the Institutional Review Board of our hospital. Patient-informed consent was waived because this was a retrospective study. The medical records of 1,461 consecutive patients with lung cancer who underwent pulmonary resection between January 1990 and April 2005 were retrospectively reviewed. Data were collected from our institutional cancer registry database and from patient follow-up medical visits, and consisted of information on preoperative patient characteristics, disease status, operative procedures, postoperative complications, pathologic diagnosis, and follow-up examinations. The data were collected at the time of hospital discharge and from outpatient charts in a prospective manner.

Of the 1,461 patients, 363 (24.8%) had been diagnosed with COPD. There were 89 patients with mild, 238 with moderate, and 36 with severe COPD, based on spirometric criteria of the Global Initiative for Chronic Obstructive Lung Disease (GOLD) as follows⁷: (1) non-COPD: forced expiratory volume in 1 second (FEV1)/forced vital capacity (FVC) \geq 70%, (2) mild COPD: FEV1/FVC $<$ 70% and FEV1 \geq 80% predicted, (3) moderate COPD: FEV1/FVC $<$ 70 and 50% \leq FEV1 $<$ 80% predicted, (4) severe COPD: FEV1/FVC $<$ 70 and 30% \leq FEV1 $<$ 50% predicted. The patients with mild to severe COPD constituted the COPD groups. All participants were involved perioperatively in a pulmonary rehabilitation program based on the severity of COPD.

Preoperative evaluation for all patients also included a detailed medical history, physical examination, blood and urine examinations, and a 12-lead electrocardiogram (ECG). Incentive spirometry and nebulization with distilled water with or without a bronchodilator were routinely encouraged to enhance lung expansion and airway clearance \sim 1 week before and after the surgery. Postoperative morbidities included bacterial pneumonia (confirmed by infiltrative shadows on chest X-ray, positive sputum culture, body temperature \geq 37.5°C and white blood cell count $>$ 10,000/ μ L); acute interstitial pneumonia (AIP) (aggravation of dyspnea on exertion, deterioration of arterial blood gases, and diffuse interstitial abnormalities compatible with acute interstitial pneumonia); mechanical ventilation \geq 3 days, bronchial stump

dehiscence; empyema (positive bacterial infection of pleural effusion); tracheostomy; and postoperative home oxygen therapy (HOT) for patients with PaO₂ $<$ 60 torr on exercise at the time of hospital discharge.

After the hospital discharge, patients visited our outpatient clinic regularly every 1 to 6 months, unless there was a tumor recurrence or any other problem. Tumor recurrence or metastasis was routinely evaluated by chest computed tomography (CT), brain magnetic resonance imaging (MRI), 2-[18F]fluoro-2-deoxy-D-glucose positron emission tomography (FDG-PET), bone scintigraphy, and several serum tumor markers every 6 to 12 months. When patients returned to referring doctors or were referred to other doctors, the status of the patient and causes of death, if death occurred, were confirmed by mail or phone with the patient, patient's family or referring doctors every 2 years. Overall and disease-specific survivals were analyzed for each patient group.

Statistical Analysis

Data were analyzed using Stat View version 5.0 (Statistical Analysis Systems; Cary, North Carolina, United States). To compare differences between the four groups, Tukey-Kramer's multiple comparison procedure was used to analyze continuous variables, and the χ^2 test used for categorical variables. Survival curves were estimated using the Kaplan-Meier method, and differences in survival times between the four groups were calculated by the log-rank test. Cox's proportional hazards model was used to identify prognostic factors.

Results

Preoperative Patient Characteristics

Preoperative patient characteristics are summarized in **Table 1**. Males and positive smoking history were more predominant in the mild to severe COPD groups than in the non-COPD group. Values for FVC, %FVC, FEV1, and FEV1/FVC were significantly lower in the mild to severe COPD groups than in the non-COPD group ($p <$ 0.05). PaO₂ values in the non-COPD and mild COPD groups were significantly higher than values in the moderate and severe COPD groups ($p <$ 0.05).

Perioperative characteristics are listed in **Table 2**. Pneumonectomy was performed in six patients of the severe COPD group despite severe pulmonary dysfunction because in those patients, the main bronchus was obstructed with tumor, and it was thought that the reduction in pulmonary function would be minimal even after pneumonectomy. Adenocarcinoma was predominant in the non-COPD and mild COPD groups, whereas squamous cell carcinoma was predominant in the moderate and severe COPD groups. Advanced cancer was more frequent in relation to the severity of COPD ($p <$ 0.0001).

Postoperative Morbidity and Mortality

Postoperative pulmonary complications and mortality are summarized in **Table 3**. The frequencies of all specific pulmonary complications and 30-day mortality were significantly higher in the severe COPD group than in any other

Table 1 Preoperative patient characteristics

Characteristics	Non-COPD	Mild COPD	Moderate COPD	Severe COPD	p-Value
	(n = 1,098)	(n = 89)	(n = 238)	(n = 36)	
Gender male, n (%)	656 (59.7)	79 (88.8)	220 (92.4)	36 (100)	< 0.0001
Gender Female	442	10	18	0	
Age (y) ^a	62.5 ± 10.3	69.1 ± 7.7 ^b	67.6 ± 7.1 ^b	67.9 ± 5.7 ^b	
BMI (kg/m ²) ^a	22.5 ± 3.2	22.3 ± 2.4	21.9 ± 3.0 ^b	21.1 ± 3.4	
PSH, n (%)	666 (60.7)	80 (89.9)	225 (94.5)	36 (100)	< 0.0001
Smoking index ^a (pack-year)	46.4 ± 28.0	57.6 ± 33.3	55.5 ± 31.1	53.5 ± 21.5	
Pulmonary function tests ^a					
FVC (L)	2.90 ± 0.76	3.59 ± 0.71 ^b	2.98 ± 0.56 ^{b,c}	2.39 ± 0.41	
%FVC (%)	97.1 ± 18.0	114.0 ± 12.7 ^b	92.4 ± 12.8 ^{b,c}	73.4 ± 10.4 ^{b,c}	
FEV1 (L)	2.38 ± 2.51	2.37 ± 0.47	1.85 ± 0.36 ^{b,c}	1.24 ± 0.25 ^{b,c}	
%FEV1 (%)	90.0 ± 17.6	90.0 ± 8.7	66.3 ± 8.3 ^{b,c}	43.0 ± 6.7 ^{b,c}	
FEV1/FVC (%)	80.0 ± 6.1	65.8 ± 3.7 ^a	62.2 ± 6.2 ^{b,c}	52.0 ± 10.1 ^{b,c}	
PaO ₂ (torr)	88.0 ± 9.6	86.7 ± 9.3	84.1 ± 9.3 ^b	82.1 ± 11.6 ^b	
PaCO ₂ (torr)	41.1 ± 3.7	40.7 ± 3.2	40.6 ± 3.5	40.6 ± 3.8	

Abbreviations: BMI, body mass index; COPD, chronic obstructive pulmonary disease; FEV1, forced expiratory volume in 1 second; forced vital capacity; PSH, positive smoking history.

^aData are presented as mean ± SD.

^b*p* < 0.05 for Tukey–Kramer's multiple comparison with data from the non-COPD group.

^c*p* < 0.05 for Tukey–Kramer's multiple comparison with data from the mild COPD group.

group. Pneumonia occurred more frequently in the mild to severe COPD groups than in the non-COPD group (*p* < 0.0001). The causes of 30-day mortality included empyema with bronchial fistula (*n* = 2), bacterial pneumonia (*n* = 1), interstitial pneumonia (*n* = 1), empyema (*n* = 1), myocardial infarction (*n* = 1), massive bleeding (*n* = 1), brain infarction (*n* = 1), deteriorating interstitial pneumonia (*n* = 1), and acute respiratory distress syndrome (*n* = 2).

Survival Analyses

► **Fig. 1** shows overall survival following the surgery. Cumulative 5-year survival rates of the four groups were 61.5% (non-COPD), 50.2% (mild COPD), 55.3% (moderate COPD), and 25.1% (severe COPD) (*p* < 0.0001). In patients with pathological stage IA disease, overall 5-year survival rates were 89.0% (non-COPD), 52.5% (mild COPD), 81.1% (moderate COPD), and 57.9% (severe COPD) (*p* < 0.0001, ► **Fig. 2**). Although there was also a significant difference in survival rates in pathological stage IB patients (*p* = 0.0038), there were no statistical differences in survival in stages IIA to IV. Although there was no significant difference in disease-specific survival for the entire study population (*p* = 0.194, ► **Fig. 3**), patients with mild to severe COPD had significantly lower disease-specific survival rates than patients with non-COPD for stage IA (*p* < 0.0001, ► **Fig. 4**). The 5-year disease-specific survival rates were 93.0% in the non-COPD, 60.8% in the mild COPD, 82.6% in the moderate COPD, and 85.7% in the severe COPD group.

Multivariate Analyses of Risk Factors for Long-Term Overall Mortality

We attempted to identify risk factors for long-term overall mortality. By univariate analysis, male gender, increased age,

decreased BMI, positive smoking history, increased tumor size, pneumonectomy, squamous cell carcinoma, pathologic stage, FEV1/FVC, % predicted FEV1, and severity of COPD were all significant factors (*p* < 0.001). By multivariate analysis, increased age, decreased BMI, positive smoking history, increased tumor size, pneumonectomy, squamous cell carcinoma, advanced pathologic stage, and severe COPD were identified as independent risk factors (► **Table 4**).

Discussion

We previously reported that stage IA lung cancer patients with COPD had poorer long-term survivals compared with those without COPD, because they had higher rates of tumor recurrence and metastasis.⁸ In this current study, we extended patient selection from stage IA to all pathologic stages and divided patients into four groups based on the severity of COPD. It was confirmed that in more severe COPD, there were higher rates of postoperative complications and poorer long-term survival than in less severe COPD. Only survival rate of patients with mild COPD seemed to be relatively worse than that of patients with moderate COPD (statistically not significant in ► **Figs. 1** to **4**). This may be due to the small number of patients with mild COPD. However, in patients with advanced lung cancer, COPD was not an important prognostic factor because of rapid disease progression.

COPD and lung cancer are closely related, and COPD is a significant risk factor for lung cancer development. Reports have shown that between 50 and 80% of patients diagnosed with lung cancer had pre-existing COPD,^{9,10} and that the prevalence of COPD in lung cancer varied from 8 to 50%.^{11,12} Furthermore, the prognosis for lung cancer patients with

Table 2 Perioperative characteristics of lung cancer patients

Characteristics	Non-COPD	Mild COPD	Moderate COPD	Severe COPD	p-Value
	(n = 1,098)	(n = 89)	(n = 238)	(n = 36)	
Operation methods					< 0.0001
Pneumonectomy, n (%)	56 (5.1)	14 (15.7)	13 (5.5)	6 (16.7)	
Lobectomy, n (%)	980 (89.3)	71 (79.8)	205 (86.1)	26 (72.2)	
Segmentectomy/partial resection, n (%)	52 (4.7)	4 (4.5)	18 (7.6)	4 (11.1)	
Tumor size ^a (cm) (maximal diameter)	3.2 ± 1.9	3.2 ± 1.9	3.7 ± 2.0	3.9 ± 1.8	
Histologic diagnoses, n (%)					< 0.0001
Ad	736 (67.6)	55 (62.5)	98 (41.4)	10 (27.8)	
Sq	239 (21.9)	30 (34.1)	113 (47.7)	23 (63.9)	
La	46 (4.2)	2 (2.3)	15 (6.3)	2 (5.6)	
Others	77 (7.0)	2 (2.3)	12 (5.0)	1 (2.8)	
Pathologic T status, n (%)					0.0001
1	488 (45.1)	38 (43.2)	84 (35.7)	10 (27.8)	
2	267 (24.7)	28 (31.8)	83 (35.3)	21 (58.3)	
3	95 (8.8)	9 (10.2)	33 (14.0)	4 (11.1)	
4	232 (21.4)	13 (14.8)	35 (14.9)	1 (2.8)	
Pathologic nodal status, n (%)					0.789
0	679 (63.1)	60 (69.0)	144 (61.8)	22 (61.1)	
1	146 (13.6)		10 (11.5)	42 (18.0)	6 (16.7)
2	218 (20.3)	15 (17.2)	45 (19.3)	7 (19.4)	
3	32 (3.0)		2 (2.3)	2 (0.9)	1 (2.8)
Pathologic staging, n (%)					0.0001
IA	396 (36.1)	29 (32.6)	63 (26.5)	10 (27.8)	
IB	132 (12.0)		18 (20.2)	46 (19.3)	10 (27.8)
IIA	47 (4.3)	1 (1.1)	8 (3.4)	0 (0)	
IIB	99 (9.0)	8 (9.0)	34 (14.3)	7 (19.4)	
IIIA	126 (11.5)	13 (14.6)	42 (17.6)	7 (19.4)	
IIIB	230 (21.0)	15 (16.9)	33 (13.9)	1 (2.8)	
IV	51 (4.7)	4 (4.5)	8 (3.4)	1 (2.8)	

Abbreviations: Ad, adenocarcinoma; COPD, chronic obstructive pulmonary disease; La, large cell carcinoma; Sq, squamous cell carcinoma.

^aSize was measured on resected specimen.

COPD is worse than for the patients without COPD because of inadequate cancer treatments and poorer pulmonary function and QOL.¹² Sin et al summarized several series on the underlying causes of death in COPD patients.¹³ They reported that the main causes of death in those with mild or moderate COPD were lung cancer and cardiovascular diseases, whereas in those with more advanced COPD (< 60% of predicted FEV1) respiratory failure was the predominant cause. This study has confirmed that poorer cancer survival was seen in patients with more advanced COPD than in non-COPD patients. This suggests that early detection of both COPD and lung cancer is needed for improving survival and maintaining QOL in patients with COPD. In this study, at least for patients who did

not have severe COPD, standard lung cancer surgery could be performed with acceptable rates of postoperative complications.

Since Cooper et al reported favorable results after lung volume reduction surgery (LVRS) for severe emphysema,¹⁴ the functional limitation for lung cancer surgery in patients with COPD has also been extended.¹⁵ However, the effect of LVRS has been thought to be limited in patients with heterogeneous bullous emphysema with regard to expectations for improved exercise tolerance, better bronchial clearance, improved quality of life, and longer survival.^{16,17} In contrast, the effects of pulmonary resection on postoperative lung function for lung cancer patients with COPD have been clarified and

Table 3 Postoperative complications and mortality

Characteristics	Non-COPD (n = 1,098)	Mild COPD (n = 89)	Moderate COPD (n = 238)	Severe COPD (n = 36)	p-Value
Characteristics	(n = 1,098)	(n = 89)	(n = 238)	(n = 36)	
Pneumonia, n (%)	47 (4.2)	9 (10.1)	31 (13.0)	5 (13.9)	< 0.0001
AIP, n (%)	32 (2.9)	3 (3.4)	5 (2.1)	4 (11.1)	0.0316
Bronchial fistula, n (%)	19 (1.7)	1 (1.1)	4 (1.7)	3 (8.3)	0.0332
Empyema, n (%)	19 (1.7)	2 (2.2)	2 (0.8)	3 (8.3)	0.0171
Prolonged mech. vent., n (%)	22 (2.0)	2 (2.2)	7 (2.9)	5 (13.9)	0.0001
HOT, n (%)	6 (0.5)	0 (0)	2 (0.8)	0 (0)	0.7860
Tracheostomy, n (%)	67 (0.6)	12 (13.5)	24 (10.1)	9 (25)	< 0.0001
SVT, n (%)	144 (13.1)	15 (16.9)	41 (17.2)	6 (16.7)	0.3138
30-day mortality, n (%)	6 (0.5)	1 (1.1)	1 (0.4)	3 (8.3)	< 0.0001

Abbreviations: AIP, acute interstitial pneumonia; COPD, chronic obstructive pulmonary disease; HOT, home oxygen therapy; prolonged mech. vent., prolonged mechanical ventilation > 3 days; SVT, supraventricular tachycardia.




Fig. 1 Overall survival in all patients after pulmonary resection for lung cancer. The 5-year survivals for the non-COPD, mild, moderate, and severe COPD groups were 61.5, 50.2, 55.3, and 25.1%, respectively. A statistically significant difference was found according to COPD grade (log-rank $p < 0.0001$).




Fig. 2 Overall survival in patients with pathological stage IA cancer after pulmonary resection for lung cancer. The 5-year survivals for the non-COPD, mild, moderate, and severe COPD groups were 89.0, 52.5, 81.0, and 57.9%, respectively. A statistically significant difference was found according to COPD grade (log-rank $p < 0.0001$).



Fig. 3 Disease-specific survival in all patients after pulmonary resection for lung cancer. The 5-year survivals for the non-COPD, mild, moderate, and severe COPD groups were 67.0, 55.7, 63.6, and 48.4%, respectively. No statistically significant difference was found according to COPD grade (log-rank $p = 0.1937$).




Fig. 4 Disease-specific survival in patients with pathological stage IA cancer after pulmonary resection for lung cancer. The 5-year survivals for the non-COPD, mild, moderate, and severe COPD groups were 93.0, 60.8, 82.6, and 85.7%, respectively. A statistically significant difference was found according to COPD grade (log-rank $p < 0.0001$).

actual postoperative function appeared to be better than predicted.^{18,19} Therefore, although predicted lung functions were not at preferred levels, we attempted as far as possible to perform standard lung cancer operations in patients with moderate to severe COPD.

By Cox's proportional hazards analyses, pneumonectomy was a significant negative risk factor for long-term survival. On the other hand, limited resection such as segmentectomy or partial resection was not a significant factor for reduced survival. It is well known that limited pulmonary resection did not confer improved perioperative morbidity, mortality, or late postoperative pulmonary function, but instead resulted in increased death and locoregional recurrence rates in patients with T1N0 nonsmall cell lung cancer.²⁰ However,

Table 4 Multivariate analysis of risk factors for overall mortality by proportional hazards model

Variables	χ^2 test	Hazard ratio	95% CI	p-Value
Gender (male)	0.811	1.139	0.858–1.514	0.3678
Age	12.85	1.019	1.008–1.029	0.0003
BMI	13.00	0.946	0.917–0.975	0.0003
Positive smoking history	5.845	1.454	1.073–1.969	0.0016
Tumor size	38.21	1.153	1.102–1.206	< 0.0001
Operation method	11.05			0.0040
Segmentectomy/partial resection		1	Reference	
Lobectomy	2.039	1.437	0.874–2.358	0.1533
Pneumonectomy	10.26	1.686	1.224–2.320	0.0014
Histology	10.57			0.0143
Ad		1	Reference	
Sq	3.864	0.808	0.653–0.999	0.0493
La	0.069	0.945	0.619–1.443	0.7934
Others	4.855	1.524	1.048–2.218	0.0276
Pathologic stage	175.0			< 0.0001
IA		1	Reference	
IB	7.402	1.639	1.148–2.340	0.0065
IIA	24.36	3.238	2.031–5.163	< 0.0001
IIB	22.17	2.381	1.659–3.418	< 0.0001
IIIA	62.82	3.739	2.698–5.180	< 0.0001
IIIB	97.37	4.483	3.328–6.040	< 0.0001
IV	115.1	8.641	5.827–12.81	< 0.0001
COPD grade	21.34			< 0.0001
Non-COPD		1	Reference	
Mild COPD	2.087	1.292	0.913–1.829	0.1486
Moderate COPD	0.156	1.049	0.827–1.331	0.6931
Severe COPD	20.23	2.774	1.778–4.328	< 0.0001

Abbreviations: Ad, adenocarcinoma; BMI, body mass index; CI, confidence interval; COPD, chronic obstructive pulmonary disease; La, large cell carcinoma; Sq, squamous cell carcinoma.

there have been many reports that segmentectomy did not cause decreased survival benefit in stage IA lung cancer patients.^{21,22} Especially in patients with severely compromised lung function, Griffin et al reported that wedge resection for stage I disease produced a survival rate similar to the rate for standard operation.²³ This suggests that early detection of lung cancer and limited resection may lead to cure despite the presence of severe COPD.

This retrospective study has certain limitations and biases. First, some of the moderate-to-severe COPD patients could not undergo postoperative adjuvant therapy and additional therapy at the time of tumor recurrence because of poor cardiopulmonary reserve. This might have affected the prognosis. Volpino et al reported that moderate-to-severe COPD patients have increased risk for noncancer death due to cardiovascular or pulmonary causes.²⁴ Second, COPD should not be defined by functional limitations only. Other factors, such as diffusion capacity, radiological findings, and clinical criteria should have been taken into consideration.²⁵ It is possible that the use of other criteria for COPD could produce different results. Third, this was a small dataset from a single-center study, and there were small numbers of patients with mild and severe COPD. Therefore, there may be several confounding variables.

In conclusion, although COPD carries a high risk for postoperative complications and poor long-term survival, appropriate selection of operative method and careful perioperative management can provide a better than expected prognosis. Patients with severe COPD should be considered carefully with regard to indications for lung cancer surgery.

References

- Brunelli A, Charloux A, Bolliger CT, et al; European Respiratory Society and European Society of Thoracic Surgeons joint task force on fitness for radical therapy. ERS/ESTS clinical guidelines on fitness for radical therapy in lung cancer patients (surgery and chemo-radiotherapy). *Eur Respir J* 2009;34(1):17–41
- Colice GL, Shafazand S, Griffin JP, Keenan R, Bolliger CT; American College of Chest Physicians. Physiologic evaluation of the patient with lung cancer being considered for resectional surgery: ACCP evidenced-based clinical practice guidelines (2nd edition). *Chest* 2007;132(3 Suppl):161S–177S
- Sekine Y, Kesler KA, Behnia M, Brooks-Brunn J, Sekine E, Brown JW. COPD may increase the incidence of refractory supraventricular arrhythmias following pulmonary resection for non-small cell lung cancer. *Chest* 2001;120(6):1783–1790
- Licker MJ, Widikker I, Robert J, et al. Operative mortality and respiratory complications after lung resection for cancer: impact of chronic obstructive pulmonary disease and time trends. *Ann Thorac Surg* 2006;81(5):1830–1837
- Sekine Y, Behnia M, Fujisawa T. Impact of COPD on pulmonary complications and on long-term survival of patients undergoing surgery for NSCLC. *Lung Cancer* 2002;37(1):95–101
- López-Encuentra A, Astudillo J, Cerezal J, Gonzalez-Aragoneses F, Novoa N, Sánchez-Palencia A; Bronchogenic Carcinoma Cooperative Group of the Spanish Society of Pneumology and Thoracic Surgery (GCCB-S). Prognostic value of chronic obstructive pulmonary disease in 2994 cases of lung cancer. *Eur J Cardiothorac Surg* 2005;27(1):8–13
- American Thoracic Society. Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 1995;152(5 Pt 2):S77–S121
- Sekine Y, Yamada Y, Chiyo M, et al. Association of chronic obstructive pulmonary disease and tumor recurrence in patients with stage IA lung cancer after complete resection. *Ann Thorac Surg* 2007;84(3):946–950
- Young RP, Hopkins RJ, Christmas T, Black PN, Metcalf P, Gamble GD. COPD prevalence is increased in lung cancer, independent of age, sex and smoking history. *Eur Respir J* 2009;34(2):380–386
- Wilson DO, Weissfeld JL, Balkan A, et al. Association of radiographic emphysema and airflow obstruction with lung cancer. *Am J Respir Crit Care Med* 2008;178(7):738–744
- Congleton J, Muers MF. The incidence of airflow obstruction in bronchial carcinoma, its relation to breathlessness, and response to bronchodilator therapy. *Respir Med* 1995;89(4):291–296
- Kurishima K, Satoh H, Ishikawa H, et al. Lung cancer patients with chronic obstructive pulmonary disease. *Oncol Rep* 2001;8(1):63–65
- Sin DD, Anthonisen NR, Soriano JB, Agusti AG. Mortality in COPD: Role of comorbidities. *Eur Respir J* 2006;28(6):1245–1257
- Cooper JD, Trulock EP, Triantafillou AN, et al. Bilateral pneumectomy (volume reduction) for chronic obstructive pulmonary disease. *J Thorac Cardiovasc Surg* 1995;109(1):106–116, discussion 116–119
- Choong CK, Meyers BF, Battafarano RJ, et al. Lung cancer resection combined with lung volume reduction in patients with severe emphysema. *J Thorac Cardiovasc Surg* 2004;127(5):1323–1331
- Benzo R, Farrell MH, Chang CC, et al; NETT Research Group. Integrating health status and survival data: the palliative effect of lung volume reduction surgery. *Am J Respir Crit Care Med* 2009;180(3):239–246
- Fessler HE, Scharf SM, Ingenito EP, McKenna RJ Jr, Sharafkhaneh A. Physiologic basis for improved pulmonary function after lung volume reduction. *Proc Am Thorac Soc* 2008;5(4):416–420
- Sekine Y, Iwata T, Chiyo M, et al. Minimal alteration of pulmonary function after lobectomy in lung cancer patients with chronic obstructive pulmonary disease. *Ann Thorac Surg* 2003;76(2):356–361, discussion 362
- Baldi S, Ruffini E, Harari S, et al. Does lobectomy for lung cancer in patients with chronic obstructive pulmonary disease affect lung function? A multicenter national study. *J Thorac Cardiovasc Surg* 2005;130(6):1616–1622
- Ginsberg RJ, Rubinstein LV; Lung Cancer Study Group. Randomized trial of lobectomy versus limited resection for T1 N0 non-small cell lung cancer. *Ann Thorac Surg* 1995;60(3):615–622, discussion 622–623
- Nakamura H, Kawasaki N, Taguchi M, Kabasawa K. Survival following lobectomy vs limited resection for stage I lung cancer: a meta-analysis. *Br J Cancer* 2005;92(6):1033–1037
- Koike T, Yamato Y, Yoshiya K, Shimoyama T, Suzuki R. Intentional limited pulmonary resection for peripheral T1 N0 M0 small-sized lung cancer. *J Thorac Cardiovasc Surg* 2003;125(4):924–928
- Griffin JP, Eastridge CE, Tolley EA, Pate JW. Wedge resection for non-small cell lung cancer in patients with pulmonary insufficiency: prospective ten-year survival. *J Thorac Oncol* 2006;1(9):960–964
- Volpino P, Cangemi R, Fiori E, et al. Risk of mortality from cardiovascular and respiratory causes in patients with chronic obstructive pulmonary disease submitted to follow-up after lung resection for non-small cell lung cancer. *J Cardiovasc Surg (Torino)* 2007;48(3):375–383
- Celli BR, Halbert RJ, Isonaka S, Schau B. Population impact of different definitions of airway obstruction. *Eur Respir J* 2003;22(2):268–273