



P. HE, X. LI, H. ZHENG, W. LI, L. LIN, X. FENG* (SICHUAN UNIVERSITY, CHENGDU, P. R OF CHINA)







Asymmetric 1,2-Reduction of Enones with Potassium Borohydride Catalyzed by Chiral *N,N'*-Dioxide–Scandium(III) Complexes

Org. Lett. **2012**, *14*, 5134–5137.

Scandium-Catalyzed Asymmetric Reduction with Potassium Borohydride



Selected examples:



Significance: As an extension on previous work using chiral *N,N'*-dioxide–metal complexes for asymmetric catalysis (see Review), the authors now describe the scandium-catalyzed asymmetric reaction of enones and ketones with KBH_4 . The resulting chiral alcohols are obtained with good yield and enantioselectivity.

Review: X. Liu, L. Lin, X. Feng *Acc. Chem. Res.* **2011**, *44*, 574–587.

Comment: Chiral allylic alcohols are important motifs widely present in natural products and biologically active molecules. The enantioselective reduction of enones is known as the most straightforward access to such motifs. Herein, the first example of catalytic enantioselective reduction of enones and ketones by using KBH_4 is reported. The utilization of an aqueous solution of KBH_4 was found to be crucial for obtaining high yield and enantioselectivity as the presence of water is believed to benefit proton transfer to accelerate the catalytic cycle. In this case, the reaction was performed in a homogeneous catalyst system. The HRMS spectra experiments indicated that the initial reducing species is KBH_3OH .

SYNFACTS Contributors: Hisashi Yamamoto, Jiajing Tan
Synfacts 2013, 9(1), 0070 Published online: 17.12.2012
DOI: 10.1055/s-0032-1317785; **Reg-No.:** H16212SF

2013 © THIEME STUTTGART • NEW YORK