D. CAMP,* M. CAMPITELLI, G. R. HANSON, I. D. JENKINS (GRIFFITH UNIVERSITY, NATHAN AND UNIVERSITY OF QUEENSLAND, ST. LUCIA, AUSTRALIA)

Formation of an Unusual Four-Membered Nitrogen Ring (Tetrazetidine) Radical Cation *J. Am. Chem. Soc.* **2012**, *134*, 16188–16196.

Formation of an Unusual Four-Membered Nitrogen Ring

Significance: Above is one of three proposed mechanisms for the formation of the new radical cationic nitrogen four-membered ring **5**. This tetrazetidine forms from allowing triphenylphosphine and diisopropyl azodicarboxylate to react. Previous studies had shown cursory evidence for the formation of a tetrazetidine by other methods, while this work offers more conclusive evidence in the form of EPR spectra and DFT calculations.

Comment: The reaction was monitored by EPR spectroscopy, which showed the presence and disappearance of **3** followed by the formation of **5**. Thus, the authors claim that the above mechanism is the most plausible. The DFT calculations show electron density on the methine or methylene protons (R = Et, *i*-Pr), which helps to rationalize hydrogen hyperfine coupling seen on the EPR spectra.

Category

Synthesis of Materials and Unnatural Products

Key words

diquinone-based cavitands molecular machines resorcin[4]arenes

SYNFACTS Contributors: Timothy M. Swager, Markrete Krikorian Synfacts 2013, 9(1), 0039 Published online: 17.12.2012

DOI: 10.1055/s-0032-1317888; Reg-No.: S13612SF