


M. A. KHALILY, O. USTAHUSEYIN, R. GARIFULLIN, R. GENÇ, M. O. GÜLER* (BILKENT UNIVERSITY, ANKARA, TURKEY)

A Supramolecular Peptide Nanofiber Templated Pd Nanocatalyst for Efficient Suzuki Coupling Reactions Under Aqueous Conditions

Chem. Commun. **2012**, 48, 11358–11360.

The Suzuki–Miyaura Coupling with Pd@Peptide



Significance: Palladium nanoparticles supported on peptide nanofiber (Pd@Peptide) were prepared by complexation of peptide nanofiber **2**, prepared via self-assembly of peptide amphiphile **1**, with Na₂PdCl₄ in aqueous NaOH followed by reduction with ascorbic acid (eq. 1). Pd@Peptide catalyzed the Suzuki–Miyaura coupling of aryl halides with arylboronic acids in water to give the corresponding biaryls in up to 99% conversion (10 examples, eq. 2)

Comment: Pd@Peptide were characterized with TEM, SEM, XRD, and TGA. In the coupling reaction of bromobenzene and 4-methoxyphenylboronic acid, the catalyst was reused four times without significant loss of catalytic activity (1st reuse: 97% conversion, 2nd reuse: 97% conversion, 3rd reuse: 97% conversion, 4th reuse: 95% conversion).

SYNFACTS Contributors: Yasuhiro Uozumi, Hiroaki Tsuji
Synfacts 2013, 9(1), 0110 Published online: 17.12.2012

DOI: 10.1055/s-0032-1317909; **Reg-No.:** Y13712SF