


Successful use of Hemospray to control refractory post-polypectomy bleeding

A 66-year-old man was referred to our department for screening colonoscopy. A polyp was found in the sigmoid colon (size 25 mm, Paris classification 0-Ip, narrow-band imaging international colorectal endoscopic [NICE] classification 2). After the submucosal injection of diluted epinephrine and lifting (▶ Fig. 1), endoscopic mucosal resection was done.

Immediately after the procedure, spurting post-polypectomy bleeding occurred. Because attempts to control the bleeding with clips were unsuccessful (▶ Fig. 2), a decision was made to apply Hemospray (Cook Medical, Bloomington, Indiana, USA). More than 20 g of hemostatic powder was applied through the 10-Fr catheter, and the bleeding was arrested (▶ Fig. 3, ▶ Video 1).

At the 24-hour second-look colonoscopy, coagulated blood was noted to be retained in the colonic lumen, but no fresh blood was seen at the polypectomy site. At the 7-day follow-up colonoscopy, a post-polypectomy ulcer with a large-caliber blood vessel was seen (▶ Fig. 4), which was subsequently closed with the combined use of clips and a nylon snare in a “tulip bundle” fashion. No perforation, symptomatic systemic embolism, or bowel obstruction was observed during the 30-day follow-up period.

Post-polypectomy bleeding is a well-known complication of colonic polypectomy, with a reported incidence ranging from 0.6% to 8.6% [1]. The risk is higher in patients who have a sessile polyp larger than 1 cm or a pedunculated polyp, such as the one in our patient [2]. Acute post-polypectomy bleeding is often immediately apparent and amenable to endoscopic therapy, and the use of hemoclips is a preferred treatment modality [3].



However, in our patient, a failure of hemostasis was observed, possibly caused by the retraction of a blood vessel deeper into the colonic wall and the subsequent inability to grasp the vessel with a clip. The use of Hemospray for the control of colonic hemorrhage was demonstrated recently [4]. However, the efficacy of this hemostatic agent in controlling spurting post-polypectomy bleeding was limited [4]. Our case contrasts with this finding, possibly because a larger amount of the powder was applied to the site of bleeding.

Although the use of Hemospray in the lower gastrointestinal tract is off label, we believe that our case adds to the experience with this novel treatment modality, in which Hemospray is applied for “rescue” hemostasis in the context of spurting post-polypectomy bleeding after failure of the primary hemostatic modality.

Endoscopy_UCTN_Code_TTT_1AQ_2AZ

Competing interests: None




Fig. 1 A polyp with a thick stalk located in the sigmoid colon, detected in a 66-year-old man undergoing screening colonoscopy. Diluted epinephrine was injected.




Fig. 2 Refractory post-polypectomy bleeding. Attempts to arrest the bleeding with clips failed.




Fig. 3 Application of Hemospray.




Fig. 4 Follow-up colonoscopy at 7 days revealed a post-polypectomy ulcer.

**Hrvoje Ivekovic¹, Branko Bilic¹,
Pave Markos¹, Nadan Rustemovic¹,
Rajko Ostojic¹, Klaus Mönkemüller²**

¹ University Hospital Centre Zagreb,
Department of Gastroenterology
and Hepatology, Zagreb, Croatia

² Division of Gastroenterology & Hepa-
tology, UAB School of Medicine,
Birmingham, Alabama, USA

References

- 1 *Kapetanios D, Beltsis A, Chatzimavroudis G et al.* Postpolypectomy bleeding: incidence, risk factors, prevention, and management. *Surg Laparosc Endosc Percutan Tech* 2012; 22: 102–107
- 2 *Kim HS, Kim TI, Kim WH et al.* Risk factors for immediate postpolypectomy bleeding of the colon: a multicenter study. *Am J Gastroenterol* 2006; 101: 1333–1341
- 3 *Conway JD, Adler DG.* ASGE Technology Committee. et al. Endoscopic hemostatic devices. *Gastrointest Endosc* 2009; 69: 987–996
- 4 *Holster IL, Brullet E, Kuipers EJ et al.* Hemo-spray treatment is effective for lower gastro-intestinal bleeding. *Endoscopy* 2014; 46: 75–78

Bibliography

DOI <http://dx.doi.org/10.1055/s-0034-1392865>
Endoscopy 2015; 47: E466–E467
© Georg Thieme Verlag KG
Stuttgart · New York
ISSN 0013-726X

Corresponding author

Hrvoje Ivekovic, MD, PhD
University Hospital Centre Zagreb
Department of Gastroenterology and Hepatology
Kispaticeva 12
10000 Zagreb
Croatia
Fax: +385-1-2388200
hrvoje.ivekovic@gmail.com