


Z.-A. HUANG, C. CHEN, X.-D. YANG,\* X.-B. FAN, W. ZHOU, C.-H. TUNG, L.-Z. WU, H. CONG\* (FUDAN UNIVERSITY, SHANGHAI, TECHNICAL INSTITUTE OF PHYSICS AND CHEMISTRY & UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING, AND PEKING UNIVERSITY, BEIJING, P. R. OF CHINA)

Synthesis of Oligoparaphenylene-Derived Nano hoops Employing an Anthracene Photodimerization–Cycloreversion Strategy

*J. Am. Chem. Soc.* **2016**, *138*, 11144–11147.

## Nanohoop through Anthracene Photodimerization–Cycloreversion



**Significance:** Nano hoops composed of highly strained aromatic hydrocarbons are fascinating carbon-rich structures, and creative strategies are required for their efficient syntheses. The authors utilize an anthracene photodimerization–cycloreversion strategy to make a novel nano hoop with interesting photophysical properties.

**Comment:** The rigid dianthracene scaffold approach generates the target nano hoop in excellent yields over six steps. The often critical ring-closing step was achieved in a particularly efficient manner under nickel-mediated Yamamoto coupling conditions.

**SYNFACTS Contributors:** Timothy M. Swager, Constantin-Christian A. Voll  
Synfacts 2016, 12(11), 1141 Published online: 18.10.2016

DOI: 10.1055/s-0036-1589379; Reg-No.: S10116SF