

Subcutaneous Dissemination in Highly Aggressive Intracranial Glioblastoma: Case Report and Literature Review

Disseminação subcutânea de glioblastoma intracraniano altamente agressivo: caso clínico e revisão da literatura

Pedro Ribeiro da Cunha¹ Gonçalo Costa¹ Marcos Barbosa¹

¹Serviço de Neurocirurgia, Centro Hospitalar Universitário de Coimbra, Coimbra, Coimbra, Portugal

Address for correspondence Pedro Ribeiro da Cunha, MD, Centro Hospitalar Universitário de Coimbra, Coimbra, Coimbra, Portugal (e-mail: pedrorcferreira@gmail.com).

Arq Bras Neurocir 2017;36:47–53.

Abstract

Glioblastoma (GBM) is a highly invasive and incurable primary central nervous system (CNS) tumor. Despite its aggressive behavior, extracranial metastases are rare, with an estimated incidence of less than 2%. In our literature review, we found only 21 reported cases of skin and soft tissue dissemination. We report a case of an early (two and a half months) postoperative skin and muscle flap-associated dissemination of a temporal glioblastoma. The particular aspect of this case, besides its rarity, is that the clinical presentation, the image reports and even the surgical findings were always in favor of a postoperative subdural empyema and epicranial abscesses. The diagnosis of soft tissue dissemination was only possible after negative microbiological cultures and histopathological confirmation of muscle and skin invasion by the tumor. This case illustrates the rare but potential risk of myocutaneous flap tumor dissemination through the durotomy/craniotomy site that can mimic a much more common, post-surgical infection.

Keywords

- ▶ soft tissue dissemination
- ▶ glioblastoma
- ▶ subdural empyema
- ▶ brain abscess

Resumo

O glioblastoma é um tumor do sistema nervoso central extremamente invasivo e incurável. Apesar do comportamento agressivo desses tumores, a metastização extracraniana é rara, apresentando uma incidência inferior a 2%. A nossa revisão da literatura revelou apenas 21 casos descritos de disseminação de glioblastoma para a pele e tecidos moles. Nós descrevemos um caso de disseminação precoce de um glioblastoma temporal para o retalho miocutâneo associado à cirurgia, cerca de 2 meses e meio após a intervenção. Além da raridade, este caso é peculiar, uma vez que tanto a forma de apresentação clínica quanto a descrição imagiológica e os achados intraoperatórios foram sempre muito sugestivos de um empiema subdural e de abscessos epicranianos. O diagnóstico definitivo de disseminação tumoral para os tecidos moles apenas foi possível após o resultado negativo das culturas microbiológicas e a confirmação histológica de invasão do músculo e da pele pelo tumor. Este caso ilustra o raro, mas potencial risco de disseminação tumoral de um glioblastoma através do retalho miocutâneo de acesso cirúrgico e do local de craniotomia/durotomia, que se pode facilmente confundir com uma situação mais frequente de infecção após cirurgia.

Palavras-chave

- ▶ disseminação nos tecidos moles
- ▶ glioblastoma
- ▶ empiema subdural
- ▶ abscesso cerebral

received
September 9, 2016
accepted
November 9, 2016
published online
December 30, 2016

DOI <http://dx.doi.org/10.1055/s-0036-1597571>.
ISSN 0103-5355.

Copyright © 2017 by Thieme Revinter Publicações Ltda, Rio de Janeiro, Brazil

License terms



Introduction

Glioblastoma (GBM) is a highly malignant tumor typically associated with a poor prognosis, with a 5-year survival rate of 4.7%.¹⁻⁶ While brain metastases secondary to neoplasms in other organs are quite common and occur in ~ 10% of cases, the opposite is much less frequent. Like the majority of the primary brain tumors, GBM does not frequently metastasize outside of the central nervous system (CNS). Death usually occurs as a consequence of the growing mass within the skull, resulting in increased intracranial pressure and herniation.

There are few reports of extracranial metastasis from GBM, and most mention an overall occurrence rate of less than 2%.¹⁻⁵ Metastases are postulated to occur by hematogenous dissemination or direct extension of the tumor.⁷ The reported sites of metastases include lung and pleura, bone, bone marrow, lymph node, and liver.⁸

Skin metastases secondary to GBM are extremely rare. To the best of our knowledge, ~ 21 cases have been reported in the literature so far (►Table 1).⁹⁻²²

In our case, the review of the radiological and intra-operative findings points to the hypothesis of soft tissue tumor dissemination through the surgical wound site.

Case Report

Presentation A 36-year-old woman with dyslipidemia, esophageal reflux and smoker of 1 pack of cigarettes a day, presented with a 3-week persistent headache. The neurological examination was normal. A computed tomography (CT) scan revealed an intra-axial irregular ring contrast-enhanced right temporal mass lesion (►Fig. 1).

Treatment The patient was submitted to surgery, and a gross total resection was accomplished. The histopathological exam confirmed giant cell GBM (grade IV). It was a solid tumor made up mainly of pleomorphic cells, with frequent multinucleated giant cells, extensive necrosis and microvascular proliferation. The mitotic index was high, and there was a markedly positive expression for vimentin, NSE, p53 and EGFR. There was no MGMT methylation or IDH1 mutation.

Outcome and follow-up The patient fulfilled the Stupp Protocol, but 3 months after surgery she had a recurrence of symptoms, with strong headaches and a right hemispheric and facial mass with an associated edema. Image studies showed the recurrence of a previous lesion in the temporal lobe, a new right frontal lobe hemorrhagic lesion, and an extra-axial antero-inferior frontal lesion with peripheral enhancing signal extending epicranially through the craniotomy, suggesting an empyema and small epicranial abscesses (►Fig. 2A-C). The differential diagnosis included abscesses, radiation-associated necrosis and soft tissue tumor dissemination.

A second surgery was performed with the enlargement of the craniotomy and the removal of purulent and necrotic subcutaneous/subdural and intra cerebral tissue. The microbiological tests were negative, with no confirmation of abscesses.

After one month, the patient was submitted to a third surgery because of extensive local recurrence with mass effect. Magnetic resonance imaging (MRI) showed extensive tumor invasion of the subcutaneous tissue and temporal muscle (►Fig. 3A-C). The surgical intervention consisted of the reopening of the previous surgical wound with extensive removal of the tumor infiltrating the temporal muscle and a wide resection of tumor recurrence in the temporal and frontal lobes.

The histology finally confirmed duramater, temporal muscle and subcutaneous tissue invasion by GBM cells (►Fig. 4B). The immunohistochemical analysis revealed no MGMT methylation, no IDH 1 mutation, but there was EGFR amplification.

She repeated the treatment with temozolomide, but died before completing it, seven months after the initial diagnosis.

Discussion

Extracranial metastases of high-grade gliomas, especially to soft tissue, skin or muscle are truly rare. They are estimated to occur in less than 2% of patients. The first report in 1988 by Shuangshoti²³ described a case of skin metastasis in a surgically removed GBM. About 21 cases of skin and soft tissue metastases have been published to date (►Table 1).

The average time for recognition of metastases since the initial tumor diagnosis has been reported to be 12.1 months.²⁴

In the majority of cases of GBM extracranial metastases previously described in the literature, patients had already undergone a neurosurgical procedure, especially craniotomies. However, there is one case of GBM metastasis outside the central nervous system, without previous surgical procedures.²⁵

There are several theories on why GBM is hardly ever correlated with extracranial metastases:

1. Because of the aggressiveness of the tumor, patients die of the primary illness before metastasis can occur.²
2. The presence of physical barriers, exclusive to the brain, as the thick dura and thickened basement membranes on blood vessels preclude both hematogenous and lymphatic spreads.^{6,26}
3. Glioma malignant cells may lack the capacity to disseminate through connective tissue and blood vessels outside the central nervous system, because within the brain parenchyma there is no collagen and fibronectin.^{7,27} Since these cells do not have the means necessary to invade connective tissue situated outside the central nervous system, they lack the ability to develop systemic metastases.

The pathophysiology of extracranial metastases is still not fully understood. The common feature of the majority of cases reported in the literature is that most patients were previously submitted to surgery. The surgical procedure does provide a break in the physical barrier, allowing seeding or hematogenous dissemination. Moreover, Müller et al²⁸ recently described that ~ 20% of GBM patients have significant

Table 1 Review of the case reports of subcutaneous and soft tissue metastasis of high-grade intracranial gliomas presented in the literature

	Sex	Age	1 st surgery	Histology (WHO grade)	Metastatic site	Time* (mo.)	Radiation/chemotherapy prior to metastasis	Second line chemotherapy
Shuangshoti et al ²³	F	69	Subtotal resection	GBM (IV)	NS	Before Su.	NS/ NS	-
Carvalho et al ⁹	F	26	Subtotal resection	GBM (IV)	Temporal/parietal areas of the scalp	16	+/- NS	-
Figueroa et al ¹⁰	M	34	Partial excision	GBM (IV)	2 cm posterior to the frontal scalp line	7	+/-	-
Santos et al ¹¹	M	42	GT resection	GBM (IV)	Skin/subcutaneous tumor seeding	36	+/-NS	-
Allan ³¹	M	60	NS	GBM (IV)	Scalp over craniotomy	12	NS/NS	-
Jain et al ¹²	M	49	"Decompression"	GBM (IV)	Adjacent to scar	10	+/-	-
Bouillot-Eimer et al ¹³	F	60	ST biopsy	GBM (IV)	Subcutaneous occipital region	8	+/- Carmustine	-
Schultz et al ¹⁴	F	74	Partial excision	GBM (IV)	Left parietal scalp	12	+/-	Stereotactic DTI-015
Saad et al ¹⁵	M	13	Subtotal resection	GBM (IV)	Scalp through skull defect, spinal leptomeninges	9	+/-Temozolamide	Thalidomide; celecoxib; etoposide
Mentrikoski et al ⁷	M	41	NS	Anap Olig (III)	Adjacent to the craniotomy site	2	-/-	-
Mentrikoski et al ⁷	F	58	NS	GBM (IV)	Adjacent to the craniotomy site	16	+/-Temozolamide	Bevacizumab
Miliaras et al ¹⁶	M	63	GT resection	GBM (IV)	Scapular subcutaneous mass	7	+/-Temozolamide	-
Senetta et al ¹⁷	F	48	GT resection	GBM (IV)	Scalp metastasis close to the surgical scar	NS	+/-Temozolamide	-
Senetta et al ¹⁷	F	54	Partial excision	GBM (IV)	Scalp. 2 cm away from the surgical scar	NS	+/-Temozolamide	-
Torres et al ²⁶	F	63	GT resection	GBM (IV)	Scalp/subcutaneous tissue near the craniotomy edge	6	+/-Temozolamide	Carmustine implants; bevacizumab; irinotecan
Armstrong et al ¹⁸	F	30	Partial excision	GBM (IV)	Below scalp, several centimeters posterior to the craniotomy scar	NS	+/-Temozolamide	Sorafenib; erlotinib; irinotecan; bevacizumab
Guo et al ¹⁹	F	19	NS	GBM (IV)	Ipsilateral posterior cervical region	8	+/-Temozolamide	-
Dawar et al ²⁷	F	57	GT resection	GBM (IV)	Subcutaneous pre-auricular mass	57	+/-Temozolamide	Mesna; doxorubicin; isofosfamide
Ginat et al ²⁰	M	62	Subtotal resection	GBM (IV)	Subcutaneous, adjacent to the craniotomy	10	+/-Temozolamide	Bevacizumab; lomustine; erlotinib
Forsyth et al ²¹	F	59	NS	GBM (IV)	Bone, subdermal over the craniotomy	6	+/-Temozolamide	Bevacizumab
Ray et al ²²	M	29	Subtotal resection	Anap Olig (III)	Posterior neck mass	13	+/-Procarbazine; lomustine; vincristine	-
Ferreira et al (present report)	F	36	GT resection	GBM (IV)	Subcutaneous and temporal muscle over the craniotomy site	3	+/-Temozolamide	-

Abbreviations: Anap Olig; F, female; GBM, glioblastoma; GT, gross total; M, male; NS, non specified; ST, stereotactic; Su., surgery. *time from first surgery to metastasis (months).




Fig. 1 Post-contrast axial CT scan showing a right temporal intra-axial irregular ring contrast-enhanced mass lesion, suggesting a high-grade glioma.

levels of circulating tumor cells (CTC) in their blood, which supports their potential for extra-CNS dissemination. Even more relevant, these CTCs were discovered both before and after surgical intervention, thus revealing that the presence of CTCs does not depend on surgical manipulation.

The pathogenic dissemination mechanism most commonly described in previous reports is the seeding of tumor cells after surgical treatment.

Additionally, reported cases of sarcomatous transformation of the tumor allows for successful implantation and growth in a non-neuronal environment.^{7,27} The current case, however, did not show any evidence of sarcomatous features.

In our case, there was an initial misdiagnosis of subdural empyema and soft tissue abscesses over the craniotomy site. This was based on the three-month previous clinical manifestations of our patient, with facial/cranial mass and edema, and also with neuroradiology reports suggesting local infection. Due to this high suspicion of infection, demanding urgent surgery, no MRI was obtained before the surgical procedure. During the second surgery, we removed a lot of necrotic tissue and what seemed to be purulent tissue invading the subcutaneous tissue and temporal muscle. After




Fig. 2 Post-contrast axial CT scans depicting a recurrence of the previous lesion in the temporal lobe (A), a new right frontal lobe hemorrhagic lesion (B), and an extra-axial antero-inferior frontal lesion with the peripheral enhancing signal extending epicranially through the craniotomy, suggesting an empyema and small epicranial abscesses (C).




Fig. 3 Post-contrast axial (A), coronal (B) and sagittal (C) MRI showing new right temporal tumor recurrence with direct extension and invasion of the subcutaneous tissue and temporal muscle, through the borders of the craniotomy.




Fig. 4 Photomicrographs (H&E) showing a giant cell glioblastoma with multinuclear cells with a bizarre aspect (A) and soft tissue infiltration by the same tumor (B).

opening the previous cranial flap and the dura, we could see the same kind of tissue continuing to the subdural space and also invading the temporal and frontal lobes.

We were quite convinced by the smell and consistency of the tissue removed that it was most likely an infection, so the collected sample was only sent to microbiological analysis, and no pathological analysis was made in this second procedure.

The microbiology results were negative, and one month later the patient was submitted to a third surgery because of extensive local recurrence with mass effect. Magnetic resonance imaging showed extensive tumor invasion of the subcutaneous tissue and temporal muscle that was linked to the intracranial tumor through the margins of the craniotomy. There was no diffusion restriction of the soft tissue lesions, and they were similar to the intracerebral lesions in all MRI sequences, excluding the possibility of abscesses and thus confirming the presence of extracranial metastases of the known glioblastoma (–Fig. 3A-C). Histology confirmed dura, temporal muscle and subcutaneous tissue invasion by the tumor (–Fig. 4A,B). The immunohistochemical analysis in our case revealed no MGMT methylation, as well as no IDH1 mutation, but there was EGFR amplification. This last genetic marker seems to promote the extracranial growth of GBM, and it has been previously associated with circulating GBM tumor cells.²⁸

The same way that MGMT methylation²⁹ and IDH ½ mutations³⁰ are known to be associated with long-term survival and better treatment responses, their absence, associated with other specific molecular factors, such as EGFR amplification, as it happened with our patient, may predict the potential for metastatic dissemination. All of these molecular signatures should probably be routinely searched in each patient in the near future, since there is a higher chance for tumor metastases with prolonged survival.

Our case illustrates the rare occurrence of soft tissue dissemination in a patient submitted to surgery for a glioblastoma. As most of the previous reported cases, the most likely mechanism of dissemination points to a migration of the glioblastoma tumor cells through the durotomy incision and the craniotomy, spreading into the temporal muscle and subcutaneous tissue. The similar tumoral aspect of the dura and soft tissue over the craniotomy site also supports the previous described theories of possible mesenchymal differentiation of the tumor when in contact with other tissues.

Thus, greater awareness of the potential for GBM metastases can contribute to the development of surgical protocols to minimize seeding, namely watertight dural closing, calvarial reconstruction, or altering instruments between the intradural and extradural parts of the surgical procedure.³¹

Furthermore, an important lesson learned is that one should consider the possibility of extracranial metastases of these tumors in the differential diagnosis of the more common surgical complications of wound infection, abscesses and subdural empyema. For this matter, an MRI exam is better than the contrast-enhanced CT scan, and should be obtained before proceeding with the surgical intervention.

Conclusions

Soft tissue dissemination in GBM is rare, and seems to be associated with younger ages and even more aggressive tumoral lesions.

The possibility of GBM soft tissue metastases should always be considered in the differential diagnosis of a possible post-surgical infection, and an MRI exam is mandatory to help with diagnosis.

The pathogenesis of GBM soft tissue metastases, the best treatment to offer to these patients and possible preventive surgical measures to avoid this complication are issues to consider for further investigation.

Disclosure

The authors have no conflicts of interest to declare. The patient's family has consented to the submission of the case report to the journal.

References

- Cuddapah VA, Robel S, Watkins S, Sontheimer H. A neurocentric perspective on glioma invasion. *Nat Rev Neurosci* 2014;15(7):455–465
- Beauchesne P. Extra-neural metastases of malignant gliomas: myth or reality? *Cancers (Basel)* 2011;3(1):461–477
- Hamilton JD, Rapp M, Schneiderhan T, et al. Glioblastoma multiforme metastasis outside the CNS: three case reports and possible mechanisms of escape. *J Clin Oncol* 2014;32(22):e80–e84
- Lun M, Lok E, Gautam S, Wu E, Wong ET. The natural history of extracranial metastasis from glioblastoma multiforme. *J Neurooncol* 2011;105(2):261–273
- Beaumont TL, Kupsky WJ, Barger GR, Sloan AE. Gliosarcoma with multiple extracranial metastases: case report and review of the literature. *J Neurooncol* 2007;83(1):39–46
- Omuro A, DeAngelis LM. Glioblastoma and other malignant gliomas: a clinical review. *JAMA* 2013;310(17):1842–1850
- Mentrikoski M, Johnson MD, Korones DN, Scott GA. Glioblastoma multiforme in skin: a report of 2 cases and review of the literature. *Am J Dermatopathol* 2008;30(4):381–384
- Cervio A, Piedimonte F, Salaberry J, et al. Bone metastases from secondary glioblastoma multiforme: a case report. *J Neurooncol* 2001;52(2):141–148
- Carvalho PA, Schwartz RB, Alexander E III, et al. Extracranial metastatic glioblastoma: appearance on thallium-201-chloride/technetium-99m-HMPAO SPECT images. *J Nucl Med* 1991;32(2):322–324
- Figuerola P, Lupton JR, Remington T, et al. Cutaneous metastasis from an intracranial glioblastoma multiforme. *J Am Acad Dermatol* 2002;46(2):297–300
- Santos AV, Saraiva PF, Santiago B. [Extracranial metastasis of glioblastoma multiforme]. *Acta Med Port* 2003;16(3):209–211
- Jain N, Mirakhor M, Flynn P, Choudhary KA. Cutaneous metastasis from glioblastoma. *Br J Neurosurg* 2005;19(1):65–68
- Bouillot-Eimer S, Loiseau H, Vital A. Subcutaneous tumoral seeding from a glioblastoma following stereotactic biopsy: case report and review of the literature. *Clin Neuropathol* 2005;24(6):247–251
- Schultz S, Pinsky GS, Wu NC, Chamberlain MC, Rodrigo AS, Martin SE. Fine needle aspiration diagnosis of extracranial glioblastoma multiforme: Case report and review of the literature. *Cytojournal* 2005;2:19
- Saad AG, Sachs J, Turner CD, et al. Extracranial metastases of glioblastoma in a child: case report and review of the literature. *J Pediatr Hematol Oncol* 2007;29(3):190–194

- 16 Miliaras G, Tsitsopoulos PP, Markoula S, Kyritsis A, Polyzoidis KS, Malamou-Mitsi V. Multifocal glioblastoma with remote cutaneous metastasis: a case report and review of the literature. *Cent Eur Neurosurg* 2009;70(1):39–42
- 17 Senetta R, Trevisan E, Rudà R, Benech F, Soffietti R, Cassoni P. Skin metastases of glioblastoma in the absence of intracranial progression are associated with a shift towards a mesenchymal immunophenotype: report of two cases. *Acta Neuropathol* 2009;118(2):313–316
- 18 Armstrong TS, Prabhu S, Aldape K, et al. A case of soft tissue metastasis from glioblastoma and review of the literature. *J Neurooncol* 2011;103(1):167–172
- 19 Guo L, Qiu Y, Ge J, Zhou D. Glioblastoma multiforme with subcutaneous metastases, case report and literature review. *J Korean Neurosurg Soc* 2012;52(5):484–487
- 20 Ginat DT, Kelly HR, Schaefer PW, Davidson CJ, Curry W. Recurrent scalp metastasis from glioblastoma following resection. *Clin Neurol Neurosurg* 2013;115(4):461–463
- 21 Forsyth TM, Bi WL, Abedalthagafi M, Dunn IF, Chiocca EA. Extracranial growth of glioblastoma multiforme. *J Clin Neurosci* 2015; 22(9):1521–1523
- 22 Ray A, Manjila S, Hdeib AM, et al. Extracranial metastasis of glioblastoma: Three illustrative cases and current review of the molecular pathology and management strategies. *Mol Clin Oncol* 2015;3(3):479–486
- 23 Shuangshoti S, Taecholarn C, Kasantikul V. Metastasizing glioblastoma multiforme diagnosed during life of patients: case report and review of literature. *J Med Assoc Thai* 1988;71(6):329–339
- 24 Vertosick FT Jr, Selker RG. Brain stem and spinal metastases of supratentorial glioblastoma multiforme: a clinical series. *Neurosurgery* 1990;27(4):516–521, discussion 521–522
- 25 Hulbanni S, Goodman PA. Glioblastoma multiforme with extraneural metastases in the absence of previous surgery. *Cancer* 1976;37(3):1577–1583
- 26 Torres IJ, Fernandez PJ, Zufiria JO, Rodriguez Barbero JM. Skin spread from an intracranial glioblastoma: case report and review of the literature. *BMJ Case Rep* 2011;20(11):1–5
- 27 Dawar R, Fabiano AJ, Qiu J, Khushalani NI. Secondary gliosarcoma with extra-cranial metastases: a report and review of the literature. *Clin Neurol Neurosurg* 2013;115(4):375–380
- 28 Müller C, Holtschmidt J, Auer M, et al. Hematogenous dissemination of glioblastoma multiforme. *Sci Transl Med* 2014;6(247): 247ra101
- 29 Hegi ME, Diserens AC, Gorlia T, et al. MGMT gene silencing and benefit from temozolomide in glioblastoma. *N Engl J Med* 2005; 352(10):997–1003
- 30 Yan H, Parsons DW, Jin G, et al. IDH1 and IDH2 mutations in gliomas. *N Engl J Med* 2009;360(8):765–773
- 31 Allan RS. Scalp metastasis from glioblastoma. *J Neurol Neurosurg Psychiatry* 2004;75(4):559