


S. HASHIGUCHI, A. FUJII, J. TAKEHARA, T. IKARIYA, R. NOYORI* (NAGOYA UNIVERSITY, JAPAN)


Asymmetric Transfer Hydrogenation of Aromatic Ketones Catalyzed by Chiral Ruthenium(II) Complex

J. Am. Chem. Soc. **1995**, *117*, 7562–7563.


Ruthenium-Catalyzed Enantioselective Transfer Hydrogenation



Selected examples:



Scale-up:



Significance: The work reported by Noyori and co-workers regarding ruthenium-catalyzed asymmetric transfer hydrogenation, represented one of the first reports describing this process with high asymmetric induction and turnover number. This transformation offers advantages over traditional hydrogenation methods that employ hydrogen gas, such as i) avoidance of pressurized H₂ gas, ii) inexpensive and synthetically convenient hydrogen sources, and iii) recyclable byproduct from the oxidation of the hydrogen surrogate (D. Wang, D. Astruc *Chem. Rev.* **2015**, *115*, 6621–6686).

Comment: Isopropanol was used as the solvent and serves as the hydrogen surrogate, generating acetone as a byproduct. The chiral alcohols were obtained in good to excellent yields, and excellent enantioselectivities. The scalability of this process was demonstrated by carrying out the reaction in a 20 mmol scale, yielding the product without erosion of either enantioselectivity or yield. The authors note that extended reaction times and higher concentrations (1 M as opposed to 0.1 M) result in the deterioration of enantioselectivity.

SYNFACTS Contributors: Mark Lautens, José F. Rodríguez
Synfacts 2019, 15(02), 0151 Published online: 18.01.2019
DOI: 10.1055/s-0037-1611998; Reg-No.: L16918SF

2019 © Georg Thieme Verlag Stuttgart · New York

Category

Metals in Synthesis

Key words

ruthenium catalysis

transfer hydrogenation

alcohols

Synfact
Classic