

β_2 -Glycoprotein I Is Proteolytically Cleaved *In Vivo* upon Activation of Fibrinolysis

Daniëlle A. Horbach^{1, 2, 3}, Erica van Oort^{1, 2}, Ton Lisman¹, Joost C. M. Meijers^{1, 3}, Ronald H. W. M. Derksen², Philip G. de Groot^{1, 3}

From the Departments of ¹Haematology and ²Rheumatology and Clinical Immunology, University Hospital Utrecht, The Netherlands and the ³Institute of Biomembranes, Utrecht University, The Netherlands

Summary

β_2 -glycoprotein I (β_2 GPI) is a plasma glycoprotein with unknown physiological function(s). *In vitro* experiments it has been demonstrated that β_2 GPI has both anticoagulant properties, such as the inhibition of factor X and prothrombin activation and procoagulant properties, such as the inhibition of the anticoagulant activity of activated protein C. Besides this, β_2 GPI bound to cardiolipin is recognized by anti-phospholipid antibodies (aPL).

In this study we demonstrate that β_2 GPI is very sensitive for cleavage between Lys317 and Thr318 by plasmin, resulting in two immunologically different cleaved forms. *In vitro* experiments show that these plasmin cleaved forms of β_2 GPI bind to negatively charged phospholipids with much lower affinity compared to intact β_2 GPI. Similar to plasmin, trypsin and elastase can also induce this proteolytical cleavage in β_2 GPI, whereas thrombin and factor Xa do not cleave β_2 GPI. The *in vivo* occurrence of the proteolytical cleavage was demonstrated by the finding that in plasmas of patients with disseminated intravascular coagulation (DIC) and in plasmas of patients treated with streptokinase, significant amounts of cleaved β_2 GPI (up to 12 μ g/ml) are present. During the development of DIC, the increase in levels of cleaved β_2 GPI is accompanied by a 70% decrease in the levels of intact β_2 GPI, whereas in streptokinase treated patients levels of intact β_2 GPI stay within the normal range.

This study demonstrates for the first time that during *in vivo* activation of fibrinolysis β_2 GPI is cleaved, which results in the formation of a form of β_2 GPI with much lower affinity for negatively charged phospholipids. Plasmin is most likely responsible for this modification.

Introduction

The presence of antiphospholipid antibodies (aPL) in plasmas of patients with systemic lupus erythematosus (SLE) is strongly correlated with a history of venous and arterial thrombosis, recurrent fetal loss and thrombocytopenia (1-4). This clinical-serological entity is called the antiphospholipid syndrome (APS). Till 1990, it was generally accepted that aPL are directed against negatively charged phospholipids and that aPL could be divided into different subclasses named anticardiolipin antibodies (aCL) and lupus anticoagulant (LAC), based on the detec-

tion method used. This idea changed by the work of different groups, who pointed to the importance of phospholipid binding plasma proteins, such as β_2 -glycoprotein I (β_2 GPI), for the binding of aPL (5-8).

β_2 GPI, also known as apolipoprotein H, is a plasma glycoprotein, which consists of 326 amino acids, has a molecular mass of approximately 50 kD and has five potential glycosylation sites (9). The protein comprises 5 repeating domains of about 60-80 amino acids, in which cysteine, proline and tryptophane residues appear to be highly conserved. Each domain resembles the so called short consensus repeats (SCR) or Sushi domains (10, 11).

Although there is considerable insight in the structure of β_2 GPI, little is known about the physiological function of the protein in human plasma. The plasma concentration of β_2 GPI is approximately 160 μ g/ml (3.8 μ M) of which 40% is bound to lipoproteins, especially chylomicrons and VLDL (12, 13). Besides binding to lipoproteins, β_2 GPI can also bind to negatively charged surfaces (14-16) such as heparin, negatively charged phospholipids and DNA. *In vitro*, β_2 GPI shows a variety of anticoagulant properties. β_2 GPI inhibits contact activation (17-19), it inhibits tenase and prothrombinase activity on activated human platelets or phospholipid vesicles (20-22) and it inhibits ADP-induced platelet aggregation (14, 16). β_2 GPI also has procoagulant activities, because it was shown that β_2 GPI can inhibit the anticoagulant activity of activated protein C (23, 24). However, the physiological importance of β_2 GPI as a regulator of coagulation is disputable, because individuals without detectable levels of β_2 GPI do not suffer from clotting or bleeding complications (25, 26).

In this study, we describe that during purification of β_2 GPI from human plasma using heparin-Sepharose affinity chromatography, proteolytical cleavage of β_2 GPI, between residues Lys317 and Thr318, occurs. We found that plasmin, trypsin and elastase can cause this cleavage *in vitro*. With monoclonal antibodies raised against cleaved and intact β_2 GPI, levels of the different forms of β_2 GPI were measured in plasmas of healthy controls, in plasmas of patients with disseminated intravascular coagulation (DIC) and in plasmas of patients treated with streptokinase. From the results it can be concluded that β_2 GPI is cleaved *in vivo* upon activation of fibrinolysis and that plasmin is most likely to be responsible for this *in vivo* modification of β_2 GPI.

Methods

Materials

The different Sepharoses used and horseradish peroxidase labeled sheep anti-M13 antibody (α -M13-PO) were from Pharmacia Biotech (Uppsala, Sweden). High binding ELISA plates (strip plate, cat.no. 9102) and regular 96-well ELISA plates were from Costar (Cambridge, MA, USA). PNGase F was from New England Biolabs (Beverly, MA, USA). BCA protein assay kit was from Pierce (Rockford, IL, USA). MaxiSorp tubes were from Nunc (Roskilde, Den-

This work was supported in part by a grant from "The Dutch League against Rheumatism" (NR 642). J. C. M. Meijers is an Established Investigator of the Netherlands Heart Foundation.

Correspondence to: Daniëlle A. Horbach, Department of Haematology, G03.647, University Hospital Utrecht, P.O.Box 85500, 3508 GA Utrecht, The Netherlands – Tel: #31-30-2506510; FAX Number: #31-30-2511893

mark). Protifar was from Nutricia (Zoetermeer, The Netherlands). *E. coli* strain XL1-Blue and helper phage VCS-M13 were from Stratagene (La Jolla, CA, USA). Polyclonal rabbit anti- β_2 GPI-antibodies (PoAb) were from Behring (Amersfoort, The Netherlands). Peroxidase labeled rabbit anti mouse immunoglobulins (RAMPO or α M) and peroxidase labeled swine anti rabbit immunoglobulins (SWARPO) were from Dako (Glostrup, Denmark). Soybean Trypsin Inhibitor (SBTI), benzamidine, polybrene, phosphatidylserine, phosphatidylcholine and phosphatidylethanolamine were from Sigma (St. Louis, MO, USA). PVDF Immobilon-P Transfer membranes were from Millipore Corporation (Bedford, MA, USA). S2251 was from Chromogenix (Mölnådal, Sweden). Streptokinase (Streptase®) was from Hoechst (Amsterdam, The Netherlands). Alkaline phosphatase conjugated goat anti human IgG was from Tago (Burlingame, CA, USA). Trypsin was from Worthington Biochemical Corp. (Freehold, NJ, USA). Elastase and automated STA APTT kit was from Boehringer Mannheim (Mannheim, Germany). Nyocard D-dimer kit was from Nycomed UK Ltd (Birmingham, UK).

Blood Collection

Blood samples were collected by venipuncture using evacuated tubes containing 3.8% trisodium citrate (0.129 M) as the anticoagulant (9:1, v/v). To obtain platelet poor plasma the samples were centrifuged twice at $2000 \times g$ for 10 min at 4° C. Plasma samples were stored at -50° C until use.

Healthy Controls and Patients

Healthy controls. Plasmas of 65 healthy controls (40 females and 25 males) were used in this study. The mean age was 32 years (range 23–51 years).

Sepsis/DIC patients. 15 sepsis/DIC patients were recruited from a primary medical-surgical intensive care unit. Their median age was 60 years (range 23–80 years). Patients were identified as having sepsis on the basis of temperature ($>38.3^\circ\text{C}$ or $<35.0^\circ\text{C}$) with a systolic blood pressure 20 mmHg below their usual level. The sepsis syndrome was verified by positive blood cultures and/or a specific site of infection. Mean platelet count was $130 \times 10^9/l$ (range 18×10^9 – 372×10^9) (27). All sepsis patients included in this study suffered from disseminated intravascular coagulation (DIC), as monitored by the presence of a prolonged activated partial thromboplastin time (APTT), prolonged prothrombin time (PT), prolonged thrombin time (TT) and elevated levels of D-dimer and F1+2.

Streptokinase treated patients. Plasmas of four patients who were treated with streptokinase (1.5×10^6 units) after acute myocardial infarction were included in this study. Samples were taken 1 h after streptokinase administration.

An almost drowned person. Multiple plasma samples were taken from a person (male, age 22 years) who was hospitalized immediately after being rescued from drowning. The patient developed severe DIC.

Activated Partial Thromboplastin Time (APTT)

APTT clotting times were measured in citrated plasma using the automated STA APTT kit, according to instructions of the manufacturer.

Levels of D-dimer

D-dimer levels were measured in citrated plasma using NyoCard D-dimer kit, according to the instructions of the manufacturer.

Proteins

β_2 GPI was purified from freshly frozen citrated human plasma (obtained from the local blood bank) as described before (3), with a few modifications. Briefly, human plasma was quickly thawed at 37° C and 40 $\mu\text{g/ml}$ Soybean Trypsin Inhibitor (SBTI), 1 mM benzamidine, 1 mM EDTA and 0.0025% polybrene were added to the plasma. The plasma was successively applied to a DEAE-Sephadex A50 column, protein G-Sepharose column and a S-Sepharose column. Afterwards, eluted proteins were applied to a heparin-Sepharose col-

umn. Bound phospholipid contaminations were eluted using 20 mM Tris/50 mM NaCl/0.1% Triton X-100 (pH 8.3). The column was washed with the same Tris buffer without Triton X-100 and bound proteins were eluted with a linear salt gradient from 50 mM to 350 mM NaCl in 20 mM Tris (pH 8.3). The peak fractions were analyzed by SDS-PAGE, and the presence of β_2 GPI was demonstrated after the proteins were transferred to a PVDF Immobilon-P Transfer membrane by blotting with rabbit anti-human β_2 GPI-antibodies. Protein concentration was determined using a BCA protein assay kit according to the manufacturer's instructions with bovine serum albumin (BSA) as a standard.

Plasminogen [purified from freshly frozen citrated human plasma, as described elsewhere (28)] was a kind gift from Dr. Peter A. Kr. von dem Borne (University Hospital Utrecht, The Netherlands). Plasminogen was activated by using a plasminogen-streptokinase complex (29). Plasmin concentration was determined using a BCA protein assay kit according to the manufacturer's instructions with BSA as a standard. The final preparation was homogeneous by SDS-PAGE.

Factor X was purified from human plasma with a monoclonal anti-factor X-antibody as described by Hackeng et al. (30). Factor X was activated by incubation with immobilized Russell viper venom as described by Bock et al. (31).

PNGase F Treatment

To remove all N-linked carbohydrate side chains from β_2 GPI, the protein (150 $\mu\text{g/ml}$) was incubated overnight at 37° C with PNGase F under reducing conditions, according to the instructions of the manufacturer.

N-terminal Sequence Analysis

β_2 GPI was subjected to 12% SDS-PAGE followed by Western blotting, according to instructions from the Sequence Centre Utrecht (Centre for Biomembranes and Lipid Enzymology, Department Biochemistry of Lipids, Utrecht, The Netherlands). The protein band was cut from the blot and the N-terminal sequence was determined by automated Edman degradation using a gas phase sequenator (Applied Biosystems model 476A) by the Sequence Centre Utrecht.

Production and Purification of Monoclonal Antibodies against β_2 GPI

BALB/c mice were immunized with a 1:1 mixture of β_2 GPI^{intact} and β_2 GPI^{intact} according to standard methods. Spleen cells were fused with Ag 8.653 myeloma cells. Fusion and hybridoma selection were performed according to standard procedures. Culture supernatants were screened for the presence of specific antibodies by an ELISA in which β_2 GPI was used as antigen. Bound antibodies were detected with peroxidase-labeled rabbit antibodies against mouse immunoglobulins (RAMPO). Positive clones were subcloned by limiting dilution and tested. This resulted in two clones (from different fusions), one named NIK-13A10 and one named OCK-4F3. MoAbs were purified from culture medium by passing the medium through a 3.5 ml α M (rabbit antibodies against mouse immunoglobulins)-Sepharose column. 9 mg α M was coupled to 3.5 ml CNBr-activated Sepharose according to the instructions of the manufacturer. Bound proteins were eluted with 3M KSCN in TBS (50 mM Tris/150 mM NaCl). Antibody containing fractions were combined, dialyzed against TBS and applied to a protein G-Sepharose FF-column (10 ml) for concentration. Bound proteins were eluted with 0.1 M glycine (pH 2.7) and antibody containing fractions were pooled and dialyzed against TBS. Purified antibodies were stored at -20° C until use.

Selection of Phage-antibodies against β_2 GPI

The phage display library used for this study was constructed as described before (32) and was kindly provided by Dr. J. de Kruif (Department of Immunology, Utrecht University, The Netherlands). Phages which express human single chain Fv antibody fragments against β_2 GPI^{intact} on the surface were selected from the phage display library using a selection procedure as described before (32), in which β_2 GPI^{intact} (50 $\mu\text{g/ml}$ in TBS, pH 7.4) was coated on

MaxiSorp tubes overnight at 4° C. The selected phages, that recognize coated β_2 GPI^{intact} were used to infect bacteria. *E. coli* strain XL1-Blue were infected with the eluted phage antibodies and grown as described before (33). The assembling of new phage particles by the infected XL1-Blue strain was accomplished using helper phage VCS-M13. Phage antibodies were isolated from the XL1-Blue culture by polyethylene glycol (PEG)/NaCl precipitation. After the last precipitation the phage containing pellet was dissolved in 3 ml PBS/1% BSA and stored at 4° C.

After each selection round, obtained phage antibodies were screened for their specific binding to β_2 GPI^{intact} using an ELISA, in which β_2 GPI^{intact} (10 μ g/ml in TBS, 100 μ l/well) was coated on high binding 96 wells ELISA plates. Wells were blocked with TBS/3% BSA/0.1% Tween-20 (150 μ l/well) for 2 h at 37° C and wells were incubated with phage antibodies at various dilutions for 1.5 h at 37° C. Afterwards bound phage antibodies were detected with 1:2500 diluted horseradish peroxidase labeled sheep anti-M13 antibody (α -M13-PO) for 1.5 h at 37° C and developed with staining solution (100 μ l/well) consisting of 0.4 mg/ml o-phenylenediamine (OPD) and 0.002% H₂O₂ in 100 mM phosphate/50 mM citric acid buffer (pH 5.0). The reaction was stopped by the addition of 1 M H₂SO₄ (50 μ l/well) and absorbance was measured at 490 nm in a multiscan photometer (Vmax reader, Molecular Devices, CA, USA). Dilutions were made in TBS/3% BSA/0.1% Tween-20 and after each incubation step, the wells were washed three times with TBS/0.1% Tween-20.

To obtain highly specific anti- β_2 GPI^{intact}-phage antibodies, the selection procedure was repeated three times. After the last selection round, individual colonies were used to prepare monoclonal phage antibodies, as described before (32). This procedure resulted in a monoclonal anti- β_2 GPI^{intact}-phage antibody, designated phage-F3.

Specificity of Different Monoclonal anti- β_2 GPI-antibodies for Cleaved and Intact β_2 GPI Preparations

Recognition of cleaved and intact β_2 GPI by phage-F3 antibodies, murine MoAbs 13A10, 4F3 and 2B2 (the latter was a generous gift of Dr. A. Tincani, Spedali Civili, Brescia, Italy) or polyclonal rabbit anti- β_2 GPI-antibodies (PoAb) was tested in an ELISA. High binding ELISA plates were coated with increasing concentrations of purified β_2 GPI (100 μ l/well) in TBS. Wells were blocked with 150 μ l/well blocking buffer (TBS/3% BSA/0.1% Tween-20) for 2 h at 37° C. Afterwards wells were incubated with 50 μ l/well phage-F3 antibodies (1:50 diluted), 13A10 (3 μ g/ml), 4F3 (3 μ g/ml), 2B2 (3 μ g/ml) or PoAb (3 μ g/ml) for 1.5 h at 37° C. Bound phage-F3 antibodies were incubated with α -M13-PO (1:2500 diluted), bound MoAbs with RAMPO (1:1000 diluted) and bound PoAbs with peroxidase labeled swine anti rabbit antibodies (SWARPO, 1:1000 diluted) for 1.5 h at 37° C. Staining procedure was performed with staining solution (100 μ l/well) consisting of 0.4 mg/ml o-phenylenediamine (OPD) and 0.002% H₂O₂ in 100 mM phosphate/50 mM citric acid buffer (pH 5.0). The reaction was stopped by the addition of 1 M H₂SO₄ (50 μ l/well) and absorbance was measured at 490 nm. Between every incubation, wells were washed three times with TBS/0.1% Tween-20. All dilutions were made in blocking buffer.

Purification of IgG from Plasma

From the plasma of a SLE-patient with high levels of anti- β_2 GPI-IgG (122%) and aCL-IgG (>100 GPL), without anti- β_2 GPI-IgM and aCL-IgM antibodies (3) total IgG was purified with protein G-Sepharose according to the manufacturer's instructions.

Binding of Patient IgG to Immobilized β_2 GPI

The binding of purified patient IgG to immobilized β_2 GPI was tested as described before (3), using the ELISA for the detection of anti- β_2 GPI-antibodies in plasma. Briefly, β_2 GPI was coated on high binding ELISA plates. After blocking aspecific binding sites, wells were incubated with 100 μ g/ml purified patient IgG, followed by incubation with alkaline phosphatase conjugated goat

antihuman IgG. Colour development was initiated by the addition of p-nitrophenyl phosphate and was stopped by the addition of 2.4 M NaOH. Absorbance was measured at 405 nm in a multiscan photometer.

Detection of β_2 GPI Levels in Plasma with ELISA

Total β_2 GPI (cleaved and intact β_2 GPI) levels in plasma were detected with an ELISA in which 100 μ l/well of a solution containing 3 μ g/ml MoAb 2B2 (in 15 mM Na₂CO₃·10H₂O, 35 mM NaHCO₃, pH 9.6) was coated on 96-wells ELISA plates overnight at 4° C. Afterwards, wells were blocked with TBS/3% BSA/0.1% Tween-20, followed by successive incubations with 1:1000 diluted plasmas (50 μ l/well, 1.5 h at 37° C) and 50 μ l/well PoAb (3 μ g/ml; 1 h at 37° C). Bound proteins were detected with 1:1000 diluted SWARPO (50 μ l/well) and developed using OPD. Purified β_2 GPI^{intact} (5 ng/ml-1 μ g/ml) was used as a standard.

For the selective detection of β_2 GPI^{intact} a similar approach was used, in which phage-F3 antibodies (1:50 diluted) instead of PoAb and α -M13-PO (1:2500 diluted) instead of SWARPO were used.

For the selective detection of β_2 GPI¹ levels in plasma, 96-wells ELISA plates were coated with MoAb 13A10 instead of MoAb 2B2. After blocking, wells were incubated with 50 μ l/well 1:5 diluted plasmas for 1.5 h at 37° C, followed by 50 μ l/well 3 μ g/ml PoAb for 1 h at 37° C. Bound antibodies were detected with 100 μ l/well 1:1000 diluted SWARPO and developed using OPD. Purified β_2 GPI¹ (10 ng/ml-1 μ g/ml) was as a standard.

Dilutions were made in TBS/3% BSA/0.1% Tween-20 and after each incubation, wells were washed three times with TBS/0.1% Tween-20.

Proteolytical Cleavage of β_2 GPI

In order to investigate the proteolytical cleavage of β_2 GPI, the effect of plasmin on β_2 GPI was studied in solution and when β_2 GPI was immobilized on high binding ELISA plates.

1. *The effect of plasmin on β_2 GPI in solution.* β_2 GPI^{intact} (200 μ g/ml) was incubated with 0.2 μ M plasmin in TBS at 37° C. At various time intervals, 10 μ l samples were taken from the incubation mixture, treated with PNGase F and subjected to 12% SDS-PAGE under reducing conditions. The proteins were visualized by Coomassie Brilliant Blue staining.

2. *The effect of plasmin, trypsin, elastase, factor Xa and α -thrombin on immobilized β_2 GPI.* β_2 GPI^{intact} (5 μ g/ml) was coated on high binding 96-wells ELISA plates as described before. Afterwards, the wells were incubated with increasing concentrations of plasmin (0-0.2 μ M), trypsin (0-0.5 μ M), elastase (0-0.5 μ M), factor Xa (0-50 nM) or α -thrombin (0-0.5 μ M) (a generous gift of Dr. Walter Kiesel, University of New Mexico, Albuquerque, NM, USA) in TBS/3% BSA/3 mM CaCl₂ for 90 min at 37° C. The different forms of β_2 GPI on the surface were measured using MoAb 4F3, MoAb 13A10 or phage-F3, followed by RAMPO or anti-M13-PO, as described before.

Preparation of Phospholipid Vesicles

Phospholipid vesicles containing 20% phosphatidylserine, 40% phosphatidylcholine and 40% phosphatidylethanolamine were prepared according to Brunner et al. (34), with some modifications as described by Van Wijnen et al. (35). The phospholipid content of the fractions was determined by phosphate analysis (36).

Binding of β_2 GPI to Phospholipids

Binding of β_2 GPI to PS/PC/PE vesicles was tested in a solid phase binding assay. High binding ELISA plates were coated with PS/PC/PE vesicles (25 μ M in TBS; 50 μ l/well) overnight at 4° C. Wells were blocked with 150 μ l/well TBS/0.5% gelatine for 2 h at 37° C. After blocking, wells were incubated with the different β_2 GPI preparations (50 μ l/well) for 1.5 h at 37° C, followed by incubation with MoAb 2B2 (3 μ g/ml; 50 μ l/well; 1.5 h at 37° C). Afterwards, the wells were incubated with RAMPO (1:1000 diluted) followed by staining procedure using OPD. All samples were diluted in TBS/0.5% gelatine and after each

incubation step, wells were washed three times with TBS. Non-specific binding was determined in wells where phospholipids were absent.

Binding of Human anti- β_2 GPI-antibodies to Cardiolipin Bound β_2 GPI

To determine whether anti- β_2 GPI-antibodies recognize β_2 GPI which is bound to coated cardiolipin (CL), an aCL-ELISA was performed as described previously (3), with a few modifications. Briefly, CL (5 μ g/well) was coated on high binding 96-well ELISA plates. Wells were blocked with blocking buffer (TBS/0.1% gelatine/0.5 mg/ml purified horse-IgG) and incubated with 100 μ g/ml purified patient IgG in the presence of the different forms of β_2 GPI (0-50 μ g/ml) for 1.5 h at 37° C. Binding of human-IgGs was visualized as described before. All dilutions were made in blocking buffer and after each incubation the wells were washed three times with TBS.

Results

Purification and Characterization of β_2 GPI from Normal Human Plasma

The final step in our purification of β_2 GPI from plasma is the elution of β_2 GPI from a heparin-Sepharose column with a linear salt gradient. This procedure results in three separate peaks at 115 mM (β_2 GPI-1, fraction 16-26), 150 mM (β_2 GPI-2, fraction 33-36) and 200 mM NaCl




Fig. 1 Heparin-Sepharose chromatography of β_2 GPI. The dotted line indicates the linear salt gradient from 50-350 mM NaCl. Closed symbols (●) correspond to the right y-axis and show levels of β_2 GPI (μ g/ml), as measured with ELISA. Open symbols (○) correspond to the left y-axis and show protein levels, as measured as the absorption at 280 nm. The horizontal bars indicate which fractions are pooled for further experiments: β_2 GPI-1 = fraction 16-26, β_2 GPI-2 = fraction 33-36 and β_2 GPI-3 = fraction 47-56




Fig. 2 SDS-PAGE and Western blot analysis of the different β_2 GPI preparations. Purified β_2 GPI preparations were separated by SDS-PAGE under non-reducing conditions, under reducing conditions and after PNGase F treatment, and visualized by Coomassie Brilliant Blue staining. Binding of rabbit anti- β_2 GPI-antibody to purified β_2 GPI was analyzed on Western blot. Lane 1: β_2 GPI-1, lane 2: β_2 GPI-2, lane 3: β_2 GPI-3. Molecular markers (in kDa) are shown on the left

(β_2 GPI-3, fraction 47-56), respectively (Fig. 1). The mutual ratio of the three peaks varied between purifications.

The three protein preparations were analyzed by SDS-PAGE under (non) reducing conditions, after PNGase F treatment (which removes all N-linked carbohydrate side chains) and by Western blotting (Fig. 2). Under non-reducing and reducing conditions the three protein preparations showed a triplet of a molecular weight of about 42 kDa and 51 kDa, respectively. Western blotting analysis showed that all three protein preparations reacted with polyclonal rabbit anti human β_2 GPI antibodies. After treatment with PNGase F under reducing conditions the three β_2 GPI preparations showed a single band with a molecular weight of approximately 38 kDa for β_2 GPI-1 and β_2 GPI-2, whereas for β_2 GPI-3 the molecular weight was about 1 kDa higher. The enzyme PNGase F is represented by the band with a molecular weight of 36 kDa.

N-terminal sequence analysis of the three β_2 GPI preparations revealed a single sequence for β_2 GPI-3 (Gly-Arg-Thr-Cys-Pro-Lys), consistent with normal intact β_2 GPI (9). Analysis of β_2 GPI-1 and β_2 GPI-2 revealed for both preparations two different N-terminal amino acid sequences, with similar molar concentrations, indicating that in these β_2 GPI preparations two N-termini are present. One sequence is identical to the sequence found for β_2 GPI-3. The other sequence (Thr-Asp-Ala-Ser-Asp-Val) is identical to that which appears when β_2 GPI is cleaved in the fifth domain between Lys317 and Thr318, as described by Hunt et al. (37).

In conclusion, heparin-Sepharose affinity chromatography results in three different forms of β_2 GPI: two "cleaved" forms β_2 GPI-1 and β_2 GPI-2, which are designated as β_2 GPI^{*1} and β_2 GPI^{*2}, respectively, and one intact form β_2 GPI-3, which is designated as β_2 GPI^{intact}.

Characterization of the Monoclonal anti- β_2 GPI-antibodies

The specificity of MoAbs 13A10, 4F3, 2B2 and phage-F3 for the three β_2 GPI forms was tested in an ELISA in which purified β_2 GPI was coated on high-binding 96 wells plates. As shown in Fig. 3, MoAb 13A10 only recognized β_2 GPI^{*1}, phage-F3 only recognized β_2 GPI^{intact}, whereas MoAb 4F3 recognized all three forms of β_2 GPI with comparable affinity. β_2 GPI^{*2} is not recognized by MoAb 13A10 nor by phage-F3. MoAb 2B2 also recognized all three forms of β_2 GPI (data not shown). Unfortunately, we were unable to raise specific MoAb for β_2 GPI^{*2}. With the different anti- β_2 GPI-antibodies, ELISAs were developed with which plasma levels of total β_2 GPI, β_2 GPI^{intact} and β_2 GPI^{*1} were measured.

Levels of Cleaved and Intact β_2 GPI in Plasmas

1. *Healthy controls.* Mean levels (\pm SD) of total β_2 GPI and β_2 GPI^{intact} in plasmas of healthy individuals were 142 ± 30 μ g/ml and 161 ± 31 μ g/ml, respectively. Only trace amounts of β_2 GPI^{*1} (0.1 ± 0.05 μ g/ml) were detectable (Fig. 4).

2. *Sepsis/DIC patients and patients treated with streptokinase.* In plasmas of the patients with sepsis/DIC levels of total β_2 GPI (73 ± 38 μ g/ml) and β_2 GPI^{intact} (65 ± 44 μ g/ml) were significantly decreased ($p < 0.0001$ and $p < 0.0001$, respectively; student's t-test) compared to the healthy control population (Fig. 4). Levels of β_2 GPI^{*1} (1.1 ± 1.2 μ g/ml) were significantly increased ($p < 0.0001$, student's t-test) compared to the healthy control population. In plasmas of four streptokinase treated patients levels of total β_2 GPI (165 ± 23 μ g/ml) and β_2 GPI^{intact} (157 ± 15 μ g/ml) were similar to their respective levels in the control population, whereas levels of β_2 GPI^{*1} (10 ± 2 μ g/ml) were




Fig. 3 Recognition of the different β_2 GPI forms by MoAb 13A10 (A), MoAb 4F3 (B) and phage F3 (C). Increasing concentrations of purified $\beta_2\text{GPI}^{\text{intact}}$ (\bullet), $\beta_2\text{GPI}^{\text{intact}}$ (\square) and $\beta_2\text{GPI}^{\text{intact}}$ (Δ), were coated on high binding 96-wells ELISA plates and were assayed for antibody recognition

approximately 100 times increased ($p < 0.0001$, student's t-test) compared to the control population (Fig. 4).

3. *A patient rescued from drowning.* APTT clotting times, levels of D-dimers (as measure of the fibrinolytic activity) and levels of total β_2 GPI, $\beta_2\text{GPI}^{\text{intact}}$ and $\beta_2\text{GPI}^{\text{intact}}$ were measured in plasma samples of a patient who was hospitalized immediately after being rescued from drowning. Within 10 h after hospitalization the APTT prolonged up to 120 s, paralleled by an increase of plasma D-dimer levels up to 24 mg/ml (Fig. 5A). During the next 5 days both APTT clotting times and D-dimer levels gradually decrease to normal values. Simultaneous with this activation of the coagulation and fibrinolysis, there was a decrease in levels of total β_2 GPI and $\beta_2\text{GPI}^{\text{intact}}$ to about 30% of the value at admission to the hospital, and an increase in levels of $\beta_2\text{GPI}^{\text{intact}}$




Fig. 4 Plasma levels of total β_2 GPI, $\beta_2\text{GPI}^{\text{intact}}$ and $\beta_2\text{GPI}^{\text{intact}}$ in sepsis patients ($n = 15$), streptokinase treated patients ($n = 4$) and healthy controls ($n = 65$). Open symbols (\circ) correspond to the left y-axis and closed symbols (\bullet) correspond to the right y-axis. Horizontal bars indicate mean β_2 GPI levels

(Fig. 5B). When coagulation and fibrinolytic activities were normalizing, there was an increase in levels of total β_2 GPI and $\beta_2\text{GPI}^{\text{intact}}$ and a decrease of levels of $\beta_2\text{GPI}^{\text{intact}}$, returning to levels found at hospitalization.

Proteolytical Cleavage of β_2 GPI

To investigate which enzymes are responsible for the cleavage of $\beta_2\text{GPI}^{\text{intact}}$, purified $\beta_2\text{GPI}^{\text{intact}}$ (200 $\mu\text{g/ml}$) was incubated with plasmin (0.2 μM), a protease with known specificity for Lys residues. At different time points samples were taken, treated with PNGase F and subjected to SDS-PAGE. Purified $\beta_2\text{GPI}^{\text{intact}}$ was used as a control. As shown in Fig. 6, during plasmin incubation a protein band with a molecular weight of approximately 1 kDa below $\beta_2\text{GPI}^{\text{intact}}$ appeared, indicating that cleaved β_2 GPI ($\beta_2\text{GPI}^{\text{intact}}$ and/or $\beta_2\text{GPI}^{\text{intact}}$) is formed. Within 60 min incubation more than 50% of the $\beta_2\text{GPI}^{\text{intact}}$ was cleaved. N-terminal sequence analysis confirmed that this plasmin treated β_2 GPI is cleaved between residues Lys317 and Thr318. In addition, heparin-Sepharose chromatographic analysis of the plasmin cleaved β_2 GPI product suggests that the plasmin cleaved form of β_2 GPI is similar to $\beta_2\text{GPI}^{\text{intact}}$ (data not shown).

We also tested the effect of plasmin on $\beta_2\text{GPI}^{\text{intact}}$ bound to a plastic surface. β_2 GPI coated on an ELISA plate, was incubated with increasing concentrations of plasmin for 90 min. Subsequently, coated β_2 GPI was incubated with MoAbs with specificity for the different forms of β_2 GPI. As shown in Fig. 7, the recognition of coated β_2 GPI by MoAb 4F3 decreases to 55% (at the highest plasmin concentration) of the originally coated β_2 GPI, indicating that upon plasmin incubation part of the coated β_2 GPI is removed from the plastic surface. The recognition by phage-F3 antibodies totally disappeared after incubation with 0.15 μM plasmin, indicating that with this plasmin concentration, coated β_2 GPI is totally cleaved by plasmin. However, cleavage of coated $\beta_2\text{GPI}^{\text{intact}}$ by plasmin does not result in the formation of $\beta_2\text{GPI}^{\text{intact}}$, because MoAb 13A10 did not recognize the plasmin treated protein. These results strongly suggest that cleavage of coated β_2 GPI by plasmin results in the formation of $\beta_2\text{GPI}^{\text{intact}}$ and not in the formation of $\beta_2\text{GPI}^{\text{intact}}$.

We also tested the proteolytical effect of trypsin, elastase, α -thrombin and factor Xa on β_2 GPI coated to an ELISA plate (data not shown). Both trypsin and elastase had similar effect as plasmin, whereas α -thrombin and activated factor Xa did not cause a cleavage of β_2 GPI.




Fig. 5 (A) APTT and D-dimer levels and (B) levels of different forms of β_2 GPI in plasmas of a patient who was rescued from drowning. Plasmas were taken immediately after hospitalization (t = 0) at various times. (A) Closed circles (●) show levels of D-dimer (right y-axis) and open circles (○) indicate APTT clotting time (left y-axis). (B) Closed squares (■) indicate levels of β_2 GPI^{*1} (left y-axis), open squares (□) indicate levels of total β_2 GPI (right




Fig. 6 Cleavage of β_2 GPI^{intact} by plasmin in solution. β_2 GPI^{intact} (200 μ g/ml) was incubated with 0.2 μ M plasmin and at various times samples were taken. After PNGase F treatment, samples were subjected to SDS-PAGE and visualized by Coomassie Brilliant Blue staining. Purified β_2 GPI^{*1} served as a control

Characteristics of the Different β_2 GPI Preparations

Binding of β_2 GPI to phospholipids. Binding of the three different β_2 GPI preparations to PS/PC/PE (20%/40%/40%) was studied in a solid-phase binding assay, in which PS/PC/PE vesicles were coated on high binding ELISA plates. Fig. 8 shows binding curves of β_2 GPI^{*1}, β_2 GPI^{*2} and β_2 GPI^{intact} to immobilized PS/PC/PE. Analysis of the specific binding revealed a β_2 GPI concentration at which half maximal binding was observed of 0.02 μ M (1 μ g/ml) for β_2 GPI^{intact} and 0.13 μ M (6.6 μ g/ml) for β_2 GPI^{*2}. For β_2 GPI^{*1} this concentration could




Fig. 7 Plasmin cleavage of immobilized β_2 GPI^{intact}. Wells of high binding ELISA plates were coated with β_2 GPI^{intact} and were incubated with increasing concentrations plasmin (0-0.2 μ M) for 90 min at 37° C. Afterwards, coated β_2 GPI was either detected by MoAb 13A10 (□), which recognizes only β_2 GPI^{*1}, MoAb 4F3 (△), which recognizes all three forms of β_2 GPI or phage F3 (●), which recognizes only β_2 GPI^{intact}. Measured OD490 nm values are expressed as percentage of originally coated β_2 GPI^{intact}. Results are expressed as mean \pm SD of three individual experiments




Fig. 8 Binding of different forms of β_2 GPI to immobilized phospholipids. PS/PC/PE (20%/40%/40%) vesicles were immobilized on high binding 96-well plates and incubated with increasing concentrations β_2 GPI^{intact} (△), β_2 GPI^{*2} (□) or β_2 GPI^{*1} (●). Afterwards, bound β_2 GPI was detected with MoAb 2B2. Results are expressed as mean \pm SD of three individual experiments

not be determined, because no saturation of binding occurred with the β_2 GPI concentrations used.

Binding of human anti- β_2 GPI-IgG to β_2 GPI. The recognition of the three different β_2 GPI preparations by anti- β_2 GPI-IgG from a SLE-patient was tested in an ELISA using total IgG purified from the plasma of a patient with high anti- β_2 GPI-antibody titres. As shown in Fig. 9A, patient anti- β_2 GPI-antibodies bind to all three β_2 GPI preparations, immobilized directly to the ELISA plates, indicating that the epitope for anti- β_2 GPI-antibodies is present on all three forms of β_2 GPI.

We next studied the binding of patient anti- β_2 GPI-antibodies to the different β_2 GPI preparations bound to cardiolipin. Fig. 9B shows that the amount of bound anti- β_2 GPI-antibodies reflects the amount of β_2 GPI bound to the cardiolipin coated well.

Discussion

In 1990 it was shown that the presence of a plasma protein, identified as β_2 GPI, was required for the binding of aPL to negatively charged phospholipids (6, 7). At that time β_2 GPI was already known as a natural anticoagulant in *in vitro* experiments. It inhibits the contact ac-




Fig. 9 Recognition of different forms of β_2 GPI by human anti- β_2 GPI-antibodies. (A) Increasing concentrations purified $\beta_2\text{GPI}^{\text{intact}}$ (Δ), $\beta_2\text{GPI}^{*2}$ (\square) or $\beta_2\text{GPI}^{*1}$ (\bullet) were coated on high binding 96-wells ELISA plates and were incubated with 100 $\mu\text{g/ml}$ purified patient IgG. Bound anti- β_2 GPI-antibodies were detected with alkaline phosphatase conjugated goat anti-human-IgG. (B) High binding 96-wells ELISA plates were coated with cardiolipin (5 $\mu\text{g/well}$) and were incubated with 100 $\mu\text{g/ml}$ purified patient-IgG, in the presence of increasing concentrations $\beta_2\text{GPI}^{\text{intact}}$ (Δ), $\beta_2\text{GPI}^{*2}$ (\square) or $\beta_2\text{GPI}^{*1}$ (\bullet). Bound anti- β_2 GPI-antibodies were detected with alkaline phosphatase conjugated goat anti-human-IgG. Results are expressed as mean \pm SD of three individual experiments

tivation of the intrinsic coagulation pathway, both tenase and prothrombinase activity and ADP-induced platelet aggregation (14, 16-22). Apart from these anticoagulant effects, β_2 GPI also inhibits the activity of APC, which is one of the most important natural anticoagulants (23, 24). Whether β_2 GPI is an important anticoagulant or procoagulant factor *in vivo* is still uncertain, because persons with a β_2 GPI deficiency lack the clinical symptoms of bleeding or thrombosis (25, 26).

In this study we have found that purified β_2 GPI is very sensitive for proteolytical cleavage, resulting in β_2 GPI which is cleaved in the fifth domain between Lys317 and Thr318. In purified systems, it was found that plasmin, trypsin and elastase are able to cause this specific cleavage. Upon proteolytical cleavage, the phospholipid binding affinity of β_2 GPI decreases (Fig. 8), which may have consequences for the proposed function(s) of β_2 GPI. It is thought that β_2 GPI inhibits pro- and anticoagulant reactions by shielding of the negatively charged phospholipid surface, which is essential as a template for most coagulation reactions. After plasmin digestion β_2 GPI might be released from these catalytic surfaces, which then results in continuation of different pro- and anticoagulant reactions. To prevent unwanted cleavage of β_2 GPI during purification, pretreatment of plasma with perchloric acid precipitation is advised. Such treatment will protect β_2 GPI against proteolytical cleavage.

Although we initially discovered the proteolytical cleavage of β_2 GPI during purification (Fig. 1), we next demonstrated that under certain pathological conditions proteolytically cleaved forms of β_2 GPI can be detected in plasma of patients (Fig. 4). Clinical syndromes in which these cleaved forms of β_2 GPI circulate were patients suffering from DIC and patients treated with streptokinase. After induction of DIC, illustrated by a prolonged APTT and increase in D-dimer levels in a patient who was rescued from drowning, the plasma level of cleaved β_2 GPI increased up to 6 $\mu\text{g/ml}$ (Fig. 5). Concomitant with the formation of cleaved β_2 GPI a drop of intact β_2 GPI was noticed. The drop of intact β_2 GPI in plasma exceeds the increase in cleaved β_2 GPI. The simultaneous occurrence of an increase of cleaved β_2 GPI and a decrease of intact β_2 GPI, might suggest that the cleavage of β_2 GPI induces a clearance of β_2 GPI. However, in plasmas of the patients treated with streptokinase strongly increased levels of cleaved β_2 GPI are present, in absence of a drop in levels of intact β_2 GPI. This suggests that cleavage of β_2 GPI itself is not the explanation for clearance of β_2 GPI in patients with DIC. In previous reports, it has been suggested that β_2 GPI is involved in the clearance of negatively charged phospholipid liposomes (38-42). It was described that *in vitro* β_2 GPI binds to different kinds of particles expressing negatively charged phospholipids, like apoptotic bodies and oxidised LDL. Therefore, a simple explanation for the disappearance of β_2 GPI from the circulation in patients with DIC is that β_2 GPI binds to negatively charged phospholipids, which are exposed abundantly on apoptotic bodies. These bodies are known to be present in patients with sepsis/DIC (43-46).

The observation of high levels of cleaved β_2 GPI in streptokinase treated patients and the observation that *in vitro* plasmin is able to cleave β_2 GPI, suggests that *in vivo* plasmin is also responsible for the proteolytical cleavage. Our *in vitro* experiments showed that cleavage of β_2 GPI by plasmin results in the formation of $\beta_2\text{GPI}^{*2}$ (Fig. 6-7), which is immunological different from the cleaved form of β_2 GPI ($\beta_2\text{GPI}^{*1}$) that we demonstrated *in vivo* (Fig. 4). These findings can point to the presence of an unknown factor in plasma which *in vivo* immediately converts plasmin cleaved $\beta_2\text{GPI}^{*2}$ into $\beta_2\text{GPI}^{*1}$. This assumption is strengthened by the observation that during purification of β_2 GPI from human plasma, both cleaved forms of β_2 GPI are formed (Fig. 1). Unfortunately, the lack of a specific antibody against $\beta_2\text{GPI}^{*2}$ made it impossible to measure the presence of $\beta_2\text{GPI}^{*2}$ directly in plasma.

Based on N-terminal sequence analysis and molecular weight determination, however, $\beta_2\text{GPI}^{*1}$ and $\beta_2\text{GPI}^{*2}$ were found to be identical (Fig. 2). At this time, we can only speculate about the differences between $\beta_2\text{GPI}^{*1}$ and $\beta_2\text{GPI}^{*2}$. It is possible that the difference between the cleaved forms of β_2 GPI is caused by a modification in glycosylation. Alternatively, the immunological differences between $\beta_2\text{GPI}^{*1}$ and $\beta_2\text{GPI}^{*2}$, however, may result from differences in tertiary structure. Another explanation might be that the differences between $\beta_2\text{GPI}^{*1}$ and $\beta_2\text{GPI}^{*2}$ are caused by the presence of different β_2 GPI allotypes.

In conclusion, we have demonstrated for the first time that *in vivo* β_2 GPI is cleaved between Lys317 and Thr318 during fibrinolysis and that *in vitro* plasmin could induce this specific proteolytical cleavage. The proteolytical cleavage of β_2 GPI is not required for the observed clearance of β_2 GPI from the circulation in patients with DIC. β_2 GPI is probably removed from the circulation by binding to negatively charged phospholipids, which are exposed on apoptotic cell particles.

After finishing this study, a study from Ohkura et al. (47) was published in which comparable proteolytical effect of plasmin on β_2 GPI was demonstrated. However, the authors did not measure the *in vivo* occurrence of the intact and cleaved forms of β_2 GPI.

Acknowledgements

The authors thank Dr. R. Fijnheer (department of Haematology, University Hospital Utrecht, The Netherlands) for the plasmas of the sepsis/DIC patients, Dr. M. Levi (department of Hemostasis and Thrombosis, University Hospital Amsterdam, The Netherlands) for the plasmas of patients treated with streptokinase and Dr. A. Tincani (Spedali Civili, Brescia, Italy) for MoAb 2B2.

References

- Love PE, Santoro SA. Antiphospholipid antibodies: anticardiolipin and the lupus anticoagulant in systemic lupus erythematosus (SLE) and in non-SLE disorders. Prevalence and clinical significance. *Ann Intern Med* 1990; 112: 682-98.
- Harris EN, Asherson RA, Hughes GR. Antiphospholipid antibodies-autoantibodies with a difference. *Annu Rev Med* 1988; 39: 261-71.
- Horbach DA, Van Oort E, Donders RCJM, Derksen RHW, De Groot PG. Lupus anticoagulant is the strongest risk factor for both venous and arterial thrombosis in patients with systemic lupus erythematosus – Comparison between different assays for the detection of antiphospholipid antibodies. *Thromb Haemost* 1996; 76: 916-24.
- Roubey RAS. Autoantibodies to phospholipid-binding plasma proteins: a new view of lupus anticoagulants and other “antiphospholipid” autoantibodies. *Blood* 1994; 84: 2854-67.
- Bevens EM, Galli M, Barbui T, Comfurius P, Zwaal RFA. Lupus anticoagulant IgG's (LA) are not directed to phospholipids only, but to a complex of lipid-bound human prothrombin. *Thromb Haemost* 1991; 66: 629-32.
- McNeil HP, Simpson RJ, Chesterman CN, Krilis SA. Antiphospholipid antibodies are directed against a complex antigen that includes a lipid binding inhibitor of coagulation: β_2 -glycoprotein I (apolipoprotein H). *Proc Natl Acad Sci USA* 1990; 87: 4120-4.
- Galli M, Comfurius P, Maassen C, Hemker HC, de Baets MH, van Breda-Vriesman PJC, Barbui T, Zwaal RFA, Bevers EM. Anticardiolipin antibodies (ACA) directed not to cardiolipin but to a plasma protein factor. *Lancet* 1990; 335: 1544-7.
- Oosting JD, Derksen RHW, Entjes HTI, Bouma BN, De Groot PG. Lupus anticoagulant activity is frequently dependent on the presence of β_2 -glycoprotein I. *Thromb Haemost* 1992; 67: 499-502.
- Lozier J, Takahashi N, Putnam FW. Complete amino acid sequence of human plasma β_2 -glycoprotein. *Proc Natl Acad Sci USA* 1984; 81: 3640-4.
- Steinkasserer A, Barlow PN, Willis AC, Kertesz Z, Campbell ID, Sim RB, Norman DG. Activity, disulphide mapping and structural modelling of the fifth domain of human β_2 -glycoprotein I. *FEBS Lett* 1992; 313: 193-7.
- Kato H, Enjyoji K. Amino acid sequence and location of the disulfide bonds in bovine β_2 -glycoprotein I: the presence of five sushi domains. *Biochemistry* 1991; 30: 11687-94.
- Polz E, Wurm H, Kostner GM. Investigations on β_2 -glycoprotein-I in the rat: isolation from serum and demonstration in lipoprotein density fractions. *Int J Biochem* 1980; 11: 265-70.
- Polz E, Kostner GM. The binding of β_2 -glycoprotein-I to human serum lipoproteins: distribution among density fractions. *FEBS Lett* 1979; 102: 183-6.
- Nimpf J, Wurm H, Kostner GM. Interaction of β_2 -glycoprotein-I with human blood platelets: influence upon the ADP-induced aggregation. *Thromb Haemost* 1985; 54: 397-401.
- Wurm H. β_2 -glycoprotein-I (apolipoprotein H) interactions with phospholipid vesicles. *Int J Biochem* 1984; 16: 511-5.
- Schousboe I. Binding of β_2 -glycoprotein I to platelets: effect of adenylate cyclase activity. *Thromb Res* 1980; 19: 225-37.
- Schousboe I. β_2 -glycoproteine I: A plasma inhibitor of the contact activation of the intrinsic blood coagulation pathway. *Blood* 1985; 66: 1086-91.
- Schousboe I, Rasmussen MS. Synchronized inhibition of the phospholipid mediated autoactivation of factor XII in plasma by β_2 -glycoprotein I and anti- β_2 -glycoprotein I. *Thromb Haemost* 1995; 73: 798-804.
- McNally T, Mackie IJ, Isenberg DA, Machin SJ. β_2 -glycoprotein-I inhibits factor XII activation on triglyceride rich lipoproteins: The effect of antibodies from plasma of patients with antiphospholipid syndrome. *Thromb Haemost* 1996; 76: 220-5.
- Goldsmith GH, Pierangeli SS, Branch DW, Gharavi AE, Harris EN. Inhibition of prothrombin activation by antiphospholipid antibodies and β_2 -glycoprotein I. *Br J Haematol* 1994; 87: 548-54.
- Shi W, Chong BH, Hogg PJ, Chesterman CN. Anticardiolipin antibodies block the inhibition by β_2 -glycoprotein I of the factor Xa generating activity of platelets. *Thromb Haemost* 1993; 70: 342-5.
- Nimpf J, Bevers EM, Bomans PHH, Till U, Wurm H, Kostner GM, Zwaal RFA. Prothrombinase activity of human platelets is inhibited by β_2 -glycoprotein-I. *Biochim Biophys Acta Gen Subj* 1986; 884: 142-9.
- Mori T, Takeya H, Nishioka J, Gabazza EC, Suzuki K. β_2 -Glycoprotein I modulates the anticoagulant activity of activated protein C on the phospholipid surface. *Thromb Haemost* 1996; 75: 49-55.
- Matsuda J, Gohchi K, Kawasugi K, Gotoh M, Saitoh N, Tsukamoto M. Inhibitory activity of anti- β_2 -glycoprotein I antibody on factor Va degradation by activated-protein C and its cofactor protein S. *Am J Hematol* 1995; 49: 89-91.
- Bancsi LF, van der Linden IK, Bertina RM. β_2 -glycoprotein I deficiency and the risk of thrombosis. *Thromb Haemost* 1992; 67: 649-53.
- Lane DA, Mannucci PM, Bauer KA, Bertina RM, Bochkov NP, Boulyjenkov V, Chandry M, Dahlbäck B, Ginter EK, Miletich JP, et al. Inherited thrombophilia: part I. *Thromb Haemost* 1996; 76: 651-62.
- Bone RC, Balk RA, Cerra FB, Dellinger RP, Fein AM, Knaus WA, Schein RMH, Sibbald WJ. Definitions for sepsis and organ failure and guidelines for the use innovative therapies in sepsis. *Chest* 1992; 101: 1644-55.
- Deutsch DG, Mertz ET. Plasminogen: purification from human plasma by affinity chromatography. *Science* 1970; 170: 1095-6.
- Lijnen HR, Van Hoef B, De Cock F, Okada K, Ueshima S, Matsuo O, Collen D. On the mechanism of fibrin-specific plasminogen activation by staphylokinase. *J Biol Chem* 1991; 266: 11826-32.
- Hackeng TM, Hessing M, Van't Veer C, Meijer-Huizinga F, Meijers JCM, De Groot PG, Van Mourik JA, Bouma BN. Protein S binding to human endothelial cells is required for expression of cofactor activity for activated protein C. *J Biol Chem* 1993; 268: 3993-4000.
- Bock PE, Craig PA, Olson ST, Singh P. Isolation of Human Blood Coagulation α -Factor X_a by Soybean Trypsin Inhibitor-Sepharose Chromatography and Its Active-Site Titration with Fluorescein Mono-*p*-guanidinobenzoate. *Arch Biochem Biophys* 1989; 273: 375-88.
- De Kruijff J, Terstappen L, Boel E, Logtenberg T. Rapid selection of cell subpopulation-specific human monoclonal antibodies from a synthetic phage antibody library. *Proc Natl Acad Sci USA* 1995; 92: 3938-42.
- Marks JD, Hoogenboom HR, Bonnert TP, McCafferty J, Griffiths AD, Winter G. By-passing immunization. Human antibodies from V-gene libraries displayed on phage. *J Mol Biol* 1991; 222: 581-97.
- Brunner J, Skrabal P, Hauser H. Single bilayer vesicles prepared without sonication. Physico-chemical properties. *Biochim Biophys Acta* 1976; 455: 322-31.
- Van Wijnen M, Stam JG, Van't Veer C, Meijers JCM, Reitsma PH, Bertina RM, Bouma BN. The interaction of protein S with the phospholipid surface is essential for the activated protein C-independent activity of protein S. *Thromb Haemost* 1996; 76: 397-403.
- Rouser G, Fkeischer S, Yamamoto A. Two dimensional thin layer chromatographic separation of polar lipids and determination of phospholipids by phosphorus analysis of spots. *Lipids* 1970; 5: 494-6.
- Hunt JE, Simpson RJ, Krilis SA. Identification of a region of β_2 -glycoprotein I critical for lipid binding and anti-cardiolipin antibody cofactor activity. *Proc Natl Acad Sci USA* 1993; 90: 2141-5.
- Chonn A, Semple SC, Cullis PR. β_2 -Glycoprotein I is a major protein associated with very rapidly cleared liposomes in vivo, suggesting a significant role in the immune clearance of “non-self” particles. *J Biol Chem* 1995; 270: 25845-9.

39. Wurm H, Polz E, Holasek A, Kostner GM. Studies on the possible function of β_2 -glycoprotein I: influence in the triglyceride metabolism in the rat. *Metabolism* 1982; 31: 484-6.
40. Balasubramanian K, Chandra J, Schroit AJ. Immune clearance of phosphatidylserine-expressing cells by phagocytes. The role of β_2 -glycoprotein I in macrophage recognition. *J Biol Chem* 1997; 272: 31113-7.
41. Hasunuma Y, Matsuura E, Makita Z, Katahira T, Nishi S, Koike T. Involvement of β_2 -glycoprotein I and anticardiolipin antibodies in oxidatively modified low-density lipoprotein uptake by macrophages. Involvement of β_2 -glycoprotein I and anticardiolipin antibodies in oxidatively modified low-density lipoprotein uptake by macrophages. *Clin Exp Immunol* 1997; 107: 569-73.
42. Manfredi AA, Rovere P, Heltai S, Galati G, Nebbia G, Tincani A, Balestrieri G, Sabbadini MG. Apoptotic cell clearance in systemic lupus erythematosus – II. Role of β_2 -glycoprotein I. *Arthritis Rheum* 1998; 41: 215-23.
43. Ayala A, Urbanich MA, Herdon CD, Chaudry IH. Is sepsis-induced apoptosis associated with macrophage dysfunction? *J Trauma* 1996; 40: 568-73.
44. Ayala A, Herdon CD, Lehman DL, Ayala CA, Chaudry IH. Differential induction of apoptosis in lymphoid tissues during sepsis: variation in onset, frequency, and the nature of the mediators. *Blood* 1996; 87: 4261-75.
45. Ayala A, Karr SM, Evans TA, Chaudry IH. Factors responsible for peritoneal granulocyte apoptosis during sepsis. *J Surg Res* 1997; 69: 67-75.
46. Williams TE, Ayala A, Chaudry IH. Inducible macrophage apoptosis following sepsis is mediated by cysteine protease activation and nitric oxide release. *J Surg Res* 1997; 70: 113-8.
47. Ohkura N, Hagihara Y, Yoshimura T, Goto Y, Kato H. Plasmin can reduce the function of human β_2 -glycoprotein I by cleaving domain V into a nicked form. *Blood* 1998; 91: 4173-9.

Received July 13, 1998 Accepted after revision September 10, 1998

DISCLAIMER Published materials reflect the authors' opinions only, not those of the editor of the journal or the publisher. Neither the editor nor the publisher assume any warranty for the correctness of the opinions or representations made by the authors. No conclusion as to whether it is a question of registered or non-registered trade marks may be drawn from the mentioning of trade names or marks in this journal.