

Errata

Facial reanimation by hypoglossal-facial neuroorrhaphy

Yvens Barbosa Fernandes, Ricardo Ramina, Hélio Sérgio Fernandes Cyrino, Marcílio Silva Prôa Júnior

Arq Bras Neurocir 33(1): 17-21, 2014

Onde se lê:

Table 2 – Clinical summary and results of 12 patients treated by hypoglossal-facial neuroorrhaphy

Case	Age, Sex	Duration of palsy (months)	Pre-op grade*	Post-op grade*
1	62, F	36	VI	III
2	65, F	12	VI	III
3	53, F	18	VI	III
4	45, F	30	VI	IV
5	51, F	3	VI	III
6	29, M	96	VI	VI
7	45, F	6	VI	III
8	46, M	8	VI	III
9	19, F	3	VI	III
10	59, F	3	VI	III
11	46, F	6	VI	III
12	55, F	3	VI	III

*Pre-operative and post-operative.

Leia-se:

Table 2 – Clinical summary and results of 12 patients treated by hypoglossal-facial neuroorrhaphy

Case	Age, Sex	Duration of palsy (months)	Pre-op grade*	Post-op grade*
1	62, F	36	I	III
2	65, F	12	I	III
3	53, F	18	I	III
4	45, F	30	I	IV
5	51, F	3	VI	III
6	29, M	96	I	VI
7	45, F	6	I	III
8	46, M	8	I	III
9	19, F	3	I	III
10	59, F	3	I	III
11	46, F	6	I	III
12	55, F	3	I	III

*Pre-operative and post-operative.