

2028

ASPIRIN AND THE KIDNEY IN PATIENTS WITH CEREBRAL ISCHEMIA.

M.Monreal (1), E.Lafoz (1), M.Foz (1), and J.Monasterio (2).  
Hospital de Badalona Germans Trias i Pujol, Spain (1), Facul-  
Tad de Medicina, Universidad Autónoma Barcelona (1), and S.  
Hemostasia, Hospital Valle Hebrón, Barcelona, Spain (2).

The acceptance of aspirin therapy for prevention of cere-  
bral ischemia is based on positive results of several large  
clinical trials, and the usual dose is 1000-1500 mg/day. Se-  
veral recent reports emphasize the adverse effects of aspirin  
and other nonsteroidal anti-inflammatory agents on renal func-  
tion. We examined wether two extreme doses of aspirin (60 mg  
vs 1200 mg/day) could alter renal function in patients recent-  
ly admitted to hospital with cerebral ischemia. We studied 33  
patients with cerebral ischemia and no history of ingestion  
of aspirin nor other drugs two weeks prior to admission. Du-  
ring the first 5 days all patients received a 50 mEq sodium  
diet and no drugs, while during the second 5 days (trial) the  
patients were randomly assigned (double blind) to placebo, as-  
pirin 20 mg or aspirin 400 mg, 3 times daily, with meals.

Overall, body weight increased by 650 g in patients taking  
1400 mg/day of aspirin ( $p < 0.01$ ), but not in patients taking  
60 mg/day. Also increases in systolic and diastolic blood pre-  
ssure did not reach significant differences. While waiting  
more reports, aspirin at doses near 1000 mg should be adminis-  
tered cautiously in sodium restricted patients with cerebral  
ischemia. Acute water retention may be specially troublesome.

2030

LIPYOXYGENASE PRODUCTS CHANGES IN 'IN VITRO' AND 'IN VIVO'  
ASPIRINISED PLATELETS UNDER THE INFLUENCE OF PAF AND EPINEPHRINE  
P.E.Makris, A.Papadopoulos, D.A.Tsakiris. First Medical Propae-  
deutic Dept, University of Thessaloniki, Greece.

We aimed to investigate the changes of lipoxygenase products  
in platelets and the simultaneous behaviour of 'in vivo' or 'in  
vitro' aspirinised platelets, stimulated by two agonists, PAF  
and epinephrine (EPI). 12 healthy were included. 6 received 20mg  
of aspirin (ASA) per os for 7 days (group A), and in 6 (group B)  
platelets were aspirinised 'in vitro' (5 or 10min incubation at  
37°C with ASA 1M). In group A blood was drawn once at the begin-  
ning and once at the end of the trial, while in group B just once.  
First, platelet aggregation was studied using two agonists simu-  
ltaneously (0.6 µM EPI and 20 nM PAF). We incubated then all pla-  
telet samples with 0.5 M of the substance BW755C (kind offer of  
Dr Moncada) for 3 min at 37°C. Second we measured PLO products  
according to Takayama et al (1980), in platelets with or without  
ASA, and in platelets with ASA and after treatment with BW755C,  
always after addition of both agonists. Our results showed: a)  
Irreversible aggregation was slightly enhanced by the simultane-  
ous addition of PAF and EPI in both groups and in non-aspirinised  
platelets. After ASA treatment, each agonist alone did not induce  
irreversible aggregation, whereas their combination overcame this  
inhibition, a fact not noticed under BW755C (a known PLO inhibi-  
tor). b) PLO products were measured in nmol TBRS/10 platelets:

	PRIME	ASA	ASA+BW755C
A ( $\bar{X} \pm SD$ )	1.44+0.37	1.70+0.44	0.80+0.30
B ( $\bar{X} \pm SD$ )	1.32+0.33	1.34+0.29	0.27+0.07

Our results agree with Cerletti et al (1986) and confirm that  
the two agonists combined are capable of overcoming the inhibi-  
tion caused by ASA, possibly by activating the PLO pathway (Cer-  
letti et al, 1986). Respectively the quantitative determination  
of PLO products (about which we did not notice any other report  
sofar) confirm the above assumption, since inhibition by  
BW755C coincides with the steep fall of PLO levels, which for  
group A is statistically significant ( $p < 0.01$ , paired t-test)  
and for group B entirely significant ( $p < 0.001$ ).

2029

CYCLOOXYGENASE INDEPENDENT PLATELET AGGREGATION:RELATION WITH ASPI-  
RIN CONCENTRATION. C. Cordova, F. Violi, D. Praticò, A. Ghiselli,  
C. Alessandri and F. Balsano. Institute of Clinical Medicine I,  
University of Rome "La Sapienza" 00161 Rome, Italy.


Low doses of aspirin (20 mg/day) were previously reported to  
be uneffective in preventing platelet aggregation (PA) induced by  
pairs of aggregating agents such as PAF and adrenalin. This was in  
part attributed to the inability of such treatment to inhibit lipo  
oxygenase-dependent PA. The latter can be observed in vitro in "aspi-  
rinated" platelets stimulated with high quantities of aggregating  
agents. The aim of this study was to evaluate if the lipooxygenase-  
dependent PA was influenced by aspirin in a dose-dependent fashion.  
PA was studied in platelet rich plasma (PRP) (Born's method) by  
using threshold doses of aggregating agents (TDA) such as PAF(4-75  
nM), epinephrine(0.6-2 µM) and collagen(2-4 µg/ml). PA performed  
in PRP pretrated with 100 µM aspirin was fully prevented; in the same  
samples thromboxane (Tx)  $A_2$  evaluated by its metabolite Tx  $B_2$  was  
almost absent. Increasing amount of PAF(20 fold TDA), epinephrine(20  
fold TDA) and collagen (36 fold TDA) do aggregate "aspirinated" pla-  
telets; similarly "aspirinated" platelets aggregate when stimulated  
with a pair of aggregating agents (TDA of PAF+epinephrine). This  
phenomenon was not detected if platelets were incubated with higher  
amounts of aspirin (250-500 µM). The study suggests that aspirin  
could influence lipooxygenase-dependent PA. This hypothesis is sup-  
ported by a research showing the aspirin inhibits dose-dependently  
platelet HETE formation. A further study is now in progress to eva-  
luate the influence of high doses of aspirin on cyclooxygenase-in-  
dependent PA in vivo.

2031

INFLUENCE OF CELL CYCLING ON MAGNITUDE OF RESPONSE OF HUMAN  
UMBILICAL VEIN ENDOTHELIAL CELLS (EC) TO NOREPINEPHRINE (NE) AND  
HISTAMINE (H). M. Gupta and G.J. Stewart. Temple Medical  
School, Philadelphia, PA 19140, USA.

EC have several roles in preventing thrombosis. Efficiency  
with which EC fulfill these might well influence outcome of  
potentially thrombotic stimuli. Thus, factors that influence EC  
function are of interest. As a general measure of EC response  
to NE and H the level of cAMP was measured. EC were harvested  
and passaged by enzymatic digestion and grown in DMEM plus 10%  
fetal calf serum and antibiotics. cAMP was assayed by  
radioimmunoassay. Percentage cycling cells was determined by  
autoradiography of cultures pulse labeled with tritiated  
thymidine and was increased by hydroxyurea treatment.

Basal cAMP was remarkably consistent (less than 10%  
difference) for cells isolated from different cords, studied  
after different number of passages, fresh or frozen and with  
varying percentage of cells cycling. However, magnitude of  
response to agonists was highly variable. Within the same cell  
lot between 5-8 passages, magnitude of response was directly  
proportional to percentage of cycling cells but magnitude of  
response for different cell lots varied over several fold.



EC response to NE was mediated through B-adrenergic receptors  
and that to H through H<sub>2</sub> receptors. Variability might be caused  
by differences in receptor number or some intracellular process.