


O. G. KULINKOVICH\*, S. V. SVIRIDOV, D. A. VASILEVSKI (BELORUSSIAN STATE UNIVERSITY, MINSK, USSR)







Titanium(IV) Isopropoxide-Catalyzed Formation of 1-Substituted Cyclopropanols in the Reaction of Ethylmagnesium Bromide with Methyl Alkanecarboxylates

*Synthesis* **1991**, 234.


## The Catalytic Kulinkovich Reaction



### Selected examples:



### Proposed mechanism:



**Significance:** The Kulinkovich reaction generates cyclopropanols from simple Grignard reagents and esters in the presence of a titanium(IV) alkoxide catalyst. This reaction has been subsequently expanded to a wide range of substrates (see Review below) and an asymmetric version was also demonstrated by the group of Corey (*J. Am. Chem. Soc.* **1994**, 116, 9345).

**Review:** O. G. Kulinkovich, A. de Meijere *Chem. Rev.* **2000**, 100, 2789–2834.

**Comment:** Although the group of Kulinkovich previously reported the synthesis of cyclopropanols through a titanium(IV) alkoxide mediated reaction (*Zh. Org. Khim.* **1989**, 25, 2244), the current report demonstrated a method that was catalytic in titanium.

SYNFACTS Contributors: Mark Lautens, Egor M. Larin  
 Synfacts 2019, 15(08), 0890 Published online: 18.07.2019  
 DOI: 10.1055/s-0039-1689804; Reg-No.: L08519SF

© 2019, Thieme. All rights reserved.  
 Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany