

J. LI, J. CHEN, R. SANG, W.-S. HAM, M. B. PLUTSCHACK, F. BERGER, S. CHABBRA, A. SCHNEGG, C. GENICOT, T. RITTER* (MAX-PLANCK-INSTITUT FÜR KOHLENFORSCHUNG, MÜLHEIM AN DER RUHR, GERMANY)
 Photoredox Catalysis with Aryl Sulfonium Salts Enables Site-Selective Late-Stage Fluorination
Nat. Chem. **2020**, *12*, 56–62.

Copper-Mediated Late-Stage Fluorination of Arenes Using Photoredox Catalysis

Category

Metals in Synthesis


Key words

photoredox catalysis

sulfonium salts

fluorination

Synfact
of the
Month



Significance: The authors report a site-selective late-stage fluorination of aryl sulfonium salts using photoredox catalysis. This newly introduced leaving group (sulfonium salt) has a relatively low redox potential and therefore provides easy access to various aryl radicals. The sulfonium group can be easily attached through thianthrene of even complex arenes, which enables a late-stage fluorination of biological active molecules.

Comment: This method was applied to fluorinate over 30 arenes, including 12 drug-like molecules. Based on detailed experimental studies, Ritter and co-workers propose that the reaction proceeds through two perfectly meshing catalytic cycles. Therein, the sulfonium leaving group plays an important part in the recycling of both the iridium and the copper catalyst.

SYNFACTS Contributors: Paul Knochel, Simon Graßl
 Synfacts 2020, 16(02), 0163 Published online: 21.01.2020
 DOI: 10.1055/s-0039-1691473; Reg-No.: P17719SF

© 2020, Thieme. All rights reserved.
 Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany