

Implementation of Transforaminal Endoscopic Lumbar Sequestrectomy in a German University Hospital Setting: A Long and Rocky Road

Michael Bender¹ Carolin Gramsch¹ Lukas Herrmann¹ Seong Woong Kim¹ Eberhard Uhl¹ Karsten Schöller¹

¹Department of Neurosurgery, Universitätsklinikum Giessen und Marburg, Giessen, Germany

²Department of Neuroradiology, Justus-Liebig-University Giessen, Giessen, Germany

Address for correspondence Dr. Michael Bender, Department of Neurosurgery, Universitätsklinikum Giessen und Marburg, Klinikstrasse 33 Giessen 35392, Germany (e-mail: michael.bender@neuro.med.uni-giessen.de).

J Neurol Surg A 2020;81:17–27.

Abstract

Objective Microsurgical diskectomy/sequestrectomy is the standard procedure for the surgical treatment of lumbar disk herniations. The transforaminal endoscopic sequestrectomy technique is a minimally invasive alternative with potential advantages such as minimal blood loss and tissue damage, as well as early mobilization of the patient. We report the implementation of this technique in a German university hospital setting.

Methods One single surgeon performed transforaminal endoscopic sequestrectomy from February 2013 to July 2016 for lumbar disk herniation in 44 patients. Demographic as well as perioperative, clinical, and radiologic data were analyzed from electronic records. Furthermore, we investigated complications, intraoperative change of the procedure to microsurgery, and reoperations. The postoperative course was analyzed using the Macnab criteria, supplemented by a questionnaire for follow-up. Pre- and postoperative magnetic resonance imaging volumetric analyses were performed to assess the radiologic efficacy of the technique.

Results Our study population had a median age of 52 years. The median follow-up was 15 months, and the median length of hospital stay was 4 days. Median duration of surgery was 100 minutes with a median blood loss of 50 mL. Surgery was most commonly performed at the L4–L5 level (63%) and in caudally migrated disk herniations (44%). In six patients, surgery was performed for recurrent disk herniations. The procedure had to be changed to conventional microsurgery in four patients. We observed no major complications. Minor complications occurred in six patients, and in four patients a reoperation was performed. Furthermore, a significantly lower Oswestry Disability Index score ($p = 0.03$), a lower Short Form 8 Health Survey (SF-8) score ($p = 0.001$), a lower visual analog scale (VAS) lower back pain score ($p = 0.03$) and VAS leg pain score ($p = 0.0008$) at the 12-month follow-up were observed in comparison with the preoperative examination. In MRI volumetry, we detected a median postoperative volume reduction of the disk herniation of 57.1% ($p = 0.02$).

Conclusions The transforaminal endoscopic sequestrectomy can be safely implemented in a university hospital setting in selected patients with primary and recurrent lumbar disk herniations, and it leads to good clinical and radiologic results. However, learning curve, caseload, and residents' microsurgical training requirements clearly affect the implementation process.

Keywords

- ▶ transforaminal endoscopic surgery
- ▶ lumbar disk herniation
- ▶ learning curve
- ▶ university hospital setting

received
October 18, 2018
accepted after revision
February 20, 2019
published online
August 29, 2019

© 2020 Georg Thieme Verlag KG
Stuttgart · New York

DOI <https://doi.org/10.1055/s-0039-1694040>.
ISSN 2193-6315.

Introduction

Microdiskectomy is a widely used and effective operation to treat lumbar disk herniations.^{1–3} However, muscular injury, facet joint violation, and epidural fibrosis can cause late sequelae.⁴ To overcome these problems, minimally invasive techniques such as transforaminal endoscopic sequestrectomy using the transforaminal endoscopic surgical system (TESSYS; Joimax GmbH, Karlsruhe, Germany) were have been established in recent years and are getting increasingly popular.^{5–9} Potential advantages of endoscopic diskectomy are less blood loss, lower rate of cerebrospinal fluid (CSF) fistulas and wound infections, as well as shorter length of hospital stay and recovery time compared with microsurgical diskectomy.^{10–14} Furthermore, in a systematic review by Nellensteijn et al, clinical results were comparable between transforaminal endoscopic surgery and microdiskectomy.⁹ Despite several advantages, endoscopic diskectomy has a significant learning curve (e.g., with longer durations of surgery and a higher reoperation rate).^{11,14} The technique is mostly performed in private practice or highly specialized spine surgery departments with or without only limited resident teaching obligations.

In contrast, microdiskectomy is the standard procedure for lumbar disk herniation surgery at our university hospital, and it is a requirement for our residents to be proficient in this procedure at the end of their training. However, due to its potential advantages, we recognized the need to innovate and integrate contemporary spinal endoscopy into our residency training program. This study was designed to demonstrate our experiences and pitfalls from the first 3 years after implementation of transforaminal endoscopic diskectomy in a German university hospital setting.

Patients and Methods

Study Design

In February 2013, transforaminal endoscopic sequestrectomy using TESSYS was introduced in our department. The current study includes 44 patients who were operated on between February 2013 and July 2016. The study protocol was approved by the institutional ethics board (No. 192/14). We included all patients with a lumbar disk herniation and an indication for surgery who (1) were judged to be suitable for a TESSYS operation by the senior spine surgeon (K.S.), and (2) gave their consent after information about the alternative of a microdiskectomy. Demographic, perioperative, and radiologic data as well as complications, intraoperative changes of the procedure to microsurgery, and reoperations were extracted from electronic records and Picture Archiving and Communication System (PACS).

Follow-up

Postoperative outcome was analyzed using the Macnab criteria,¹⁵ the return-to-work rate, the analgesic medication, and the overall satisfaction with help of a questionnaire. Overall satisfaction was assessed with a two-point scale: (1) I

would consider the same operation again, or (2) I would not want to be operated again with the TESSYS technique. The mode of analgesic medication was determined by using a 4-point scale: (1) no analgesic medication, (2) less analgesic medication, (3) same analgesic medication, and (4) more analgesic medication.

The final 11 patients of our study cohort operated on between June 2015 and July 2016 were additionally analyzed in a prospective manner. The Oswestry Disability Index (ODI), horizontal visual analog scale (VAS) for lower back pain and leg pain, and Short Form 8 (SF-8) Health Survey for quality-of-life evaluation were used at discharge, 6 weeks after surgery, as well as at 12 months after surgery.^{16–18}

Procedure

In this study all endoscopic transforaminal lumbar sequestrectomies were performed with the TESSYS method under general anesthesia by one senior neurosurgeon with a subspecialization in spine surgery and fellowship training in minimally invasive spinal procedures (K.S.). All patients were treated in the prone position.

After detection of the correct surgical level with the C-arm (Veradius, Philips GmbH, Hamburg, Germany) and adjustment of the operating table (radiograph criteria: spinous processes in line and pedicle eyes symmetrical and clearly visible in anteroposterior [AP] views, with no double configuration of posterior vertebral body walls or pedicles in lateral views), the skin was sterilized and draped in a standard fashion. A hollow needle was inserted posterolaterally between 10 and 15 cm from the midline depending on the surgical level, the localization of the sequester, and the physiognomy of the patient and was advanced into the lateral neuroforamen under radiographic control. A Seldinger wire was then introduced into the needle, and, after removal of the needle, a linear skin incision of ~0.8 cm was performed. A bent rod was then advanced transforaminally with the Seldinger technique into the anterior spinal canal (►Fig. 1) just passing the midline, followed by sequential dilation of the transforaminal trajectory with different tubes and sequential reaming of the caudo-posterior neuroforamen. Finally, the working cannula was placed in the vicinity of the pathology (►Fig. 2). All approach steps were conducted under repetitive radiographic control.

As a next step, an endoscope with a 30-degree view (6.3 mm outer diameter), and a working, an irrigation and a suction channel was then inserted through the working cannula. After identification of the pedicle, the dural sac, and the exiting nerve root, the herniated disk was identified, mobilized, and removed with different hooks and forceps (►Figs. 3 and 4). Hemostasis was performed using a radiofrequency probe (VaporFlex; Joimax GmbH, Karlsruhe, Germany). After complete removal of the herniated disk tissue, the endoscope and the working cannula were removed. The skin was closed by single sutures (►Fig. 5). No specific measures were taken in cases with a dural tear.

Mobilization was started immediately after surgery, and physiotherapy was initiated on postoperative day 1. Preoperative analgesic medication was continued 3 days after




Fig. 1 Intraoperative lateral radiographic view after transforaminal insertion of a bent rod into the anterior epidural space at level L4–L5.

surgery and was subsequently reduced as required thereafter.

MRI Volumetric Analysis

Imaging

Magnetic resonance imaging (MRI) was performed on a 3-T MR Scanner (Verio; Siemens, Erlangen, Germany) equipped with 4 channels of a 12-channel surface coil. The thoracolumbar spine of each patient was imaged using sagittal T2-weighted sequences (TR/TE, 3,000/104 ms, field of view




Fig. 2 Intraoperative anteroposterior radiographic visualization of the working cannula at level L5–S1 with its opening just medial to the pedicle.



Fig. 3 Intraoperative setting. Removal of a disk herniation with endoscopic forceps.




Fig. 4 Intraoperative anteroposterior radiographic visualization of an endoscopic telescope hook used to mobilize a disk herniation just medial to the S1 pedicle.



Fig. 5 The ~0.8-cm-long skin incision was closed by two single sutures.




Fig. 7 The sagittal slices of the T2-weighted sequences. A radiologist manually outlined the herniated disk in these slices.

[FOV] 280×280 , slice thickness 2 mm) and constructive interference in steady state (CISS) sequences (TR/TE 5,51/2,39 ms, FOV 0.7 mm isotropic voxels) preoperatively and between postoperative day 1 and 3 in the final 11 patients of our cohort in a prospective manner.

Volumetric Analysis

MR images were viewed and postprocessed by an experienced neuroradiologist (C.G.) on an PACS workstation (INFINITT Healthcare Co., Brussels, Belgium). The sagittal slices of the T2-weighted sequences were matched with the axial slices of the CISS sequence. With the aid of the CISS sequence, the individual midline was identified in each data set (►Fig. 6). Only those sagittal slices that represented herniated disk material of the most affected side were used for volumetry (i.e., slices were identified from the midline to the most lateral slice just containing herniated disk material). The neuroradiologist manually outlined the herniated disk in these slices as

shown in ►Fig. 7. The volume of the disk material initially measured pre- and postoperatively as an area in sagittal slices was calculated as the product of these areas, the number of the respective adjacent slices, and the slice thickness (2 mm).

Statistical Analysis

For data analysis and graphic illustration, Graph Pad Prism v.5 (GraphPad Software, Inc., La Jolla, California, United States) was used. Data were expressed as median plus or minus range. The chi-square test (categorical variables) and the Mann-Whitney rank sum test (continuous variables) were used for intergroup comparisons. A $p < 0.05$ was defined as the level of significance.

Results

Patients of our study group had a median age of 52 years (range: 25–75 years). The median duration of hospital stay




Fig. 6 The sagittal and axial slices of the T2-weighted constructive interference in steady state (CISS) sequences. With help of the CISS sequence, the individual midline was identified in each data set.

was 4 days (range: 1–13 days), and the median follow-up was 15 months (range: 7–30 months). A total of 40 patients were available for follow-up (one patient had died, with no association to surgery, two patients did not respond to the questionnaire, and one patient was lost to follow-up). Baseline data are summarized in ► **Table 1**.

Perioperative Data

Surgery was mostly performed at level L4–L5 (63%) and in caudally migrated disk herniations (44%). In 42 patients, surgery was performed at one level, and 2 patients were operated on at two levels. Neurophysiologic monitoring was used in seven patients; however, monitoring was abandoned later because we did not find any benefit from it. In the first two patients, operations were supported by an external surgeon with vast experience in transforaminal endoscopic surgery; 13 further operations were supported by an experienced application specialist from the Joimax Company with a scrub nurse background.

Table 1 Baseline data of the study population^a

	Study population (n = 44)
Demographics	
Median age, y (range)	52 (25–78)
Women, n (%)	18 (41)
Median BMI, kg/m ² (range)	28 (20–47)
Median ASA score (range)	2 (1–3)
Median duration of symptoms, wk (range)	13 (1–53)
Spinal level of disk herniation, n (%)	
L1–L2	1 (2)
L2–L3	1 (2)
L3–L4	5 (11)
L4–L5	29 (63)
L5–S1	10 (22)
Localization of disk herniation, n (%)	
Right side	26 (59)
Left side	18 (41)
Cranial sequestration	1 (2)
Caudal sequestration	23 (50)
Mediolateral sequestration	16 (35)
Intraforaminal sequestration	2 (4)
Intra-extraforaminal sequestration	1 (2)
Extraforaminal sequestration	3 (7)
Type of disk herniation, n (%)	
Primary disk herniation	38 (86)
Recurrent disk herniation	6 (14)

Abbreviations: ASA, American Society of Anesthesiologists; BMI, body mass index.

^aIn two patients, surgery was performed at two levels.

The median duration of surgery was 100 minutes (range: 40–234 minutes) with a median blood loss of 50 mL (range: 5–1,005 mL). We observed no significant difference regarding duration of surgery ($p = 0.50$) and blood loss ($p = 0.56$) between the first 22 and last 22 patients. We observed a median intraoperative radiation dose of 15.8 cGy cm² (range: 3.3–318 cGy cm²).

The procedure had to be changed to microsurgery in four patients: In two patients who were operated on at level L4–L5, a transmuscular tubular approach was applied due to intraspinal bleeding that was not adequately controlled by endoscopy. In two further patients who were operated at level L5–S1, a high iliac crest in conjunction with neuroforaminal stenosis and ligamentous hypertrophy prohibited a controlled endoscopic removal of the disk herniation, and the procedure was changed to a subperiosteal specular approach.

Our study population experienced no major complications. In six patients the following minor complications occurred: five patients exhibited a temporary neurologic deficit (three patients with worsened and two patients with new paresis), and one patient experienced an early recurrent disk herniation 2 days after surgery. A dural tear occurred in six patients; however, there was no postoperative CSF fistula in our series.

Reoperations were performed in four patients due to one early recurrent disk herniation, one late recurrent disk herniation 8 weeks after surgery, and two new neurologic deficits on the day of the operation and assumed postoperative hemorrhage on MRI that could not, however, be confirmed on surgical inspection.

Follow-up

At follow-up, the median Macnab score was 2 (range: 1–4). We determined an overall satisfaction of 90%. A total of 95% of the formerly working patients were able to return to work. No lower back pain or reduced pain was observed in 80% of the patients, and 85% had no or reduced leg pain. A reduction or no further need for analgesic medication was observed in 95% of the patients. In the 11 prospectively analyzed patients, we observed a significantly lower ODI score ($p = 0.03$), SF-8 score ($p = 0.001$), VAS lower back pain score ($p = 0.03$), and VAS leg pain score ($p = 0.0008$) in comparison with the preoperative examination. ► **Table 2** summarizes the results of the prospective follow-up examinations.

MRI Volumetric Analysis

Volumetric analysis of the preoperative and postoperative MRI scans was performed in 10 patients; in one case we abandoned the preoperative volumetric scan, due to an emergency indication for surgery. The median preoperative disk herniation volume was 1.4 cm³ (range: 0.3–1.9 cm³) compared with a postoperative volume of 0.6 cm³ (range: 0.2–1.4 cm³). We found a statistically significant disk volume reduction of 57.1% ($p = 0.02$), as shown in ► **Fig. 8**.


This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

Table 2 Follow-up examination of prospectively analyzed patients ($n = 11$)^a

	Median	Range	<i>p</i> Value
ODI			
Preoperative	25	10–39	
At discharge	21	1–42	0.32
6-wk FU	12	0–40	0.06
12-mo FU	6.5	0–29	0.003
SF-8			
Preoperative	33	23–39	
At discharge	29	23–39	0.45
6-wk FU	25	8–43	0.1
12-mo FU	21.5	12–32	0.001
VAS lower back pain			
Preoperative	5	2–9	
At discharge	2	1–4	0.01
6-wk FU	3	1–10	0.05
12-mo FU	2	1–7	0.03
VAS leg pain			
Preoperative	6	2–10	
At discharge	2	1–10	0.01
6-wk FU	3	1–8	0.008
12-mo FU	2	1–5	0.0008

Abbreviations: FU, follow-up; ODI, Oswestry Disability Index, SF-8, Short Form-8 Health Survey, VAS, visual analog scale.

^aWe observed a significantly lower ODI score ($p = 0.03$), a lower SF-8 score ($p = 0.001$), and a lower VAS lower back pain score ($p = 0.03$) and VAS leg pain score ($p = 0.0008$) at the 12-month FU examination in comparison with the preoperative examination. Statistical analysis was performed with the Mann-Whitney *U* test.

**Fig. 8** Magnetic resonance imaging volumetric analysis. We observed a significant volume reduction of disk herniation volume in the postoperative volumetric analysis ($p = 0.02$).

Discussion

Advantages of Endoscopic Lumbar Discectomy

Minimal muscle trauma and blood loss, low rates of wound infections and CSF fistulas, as well as shorter hospital stay and recovery time, are potential advantages of transforaminal endoscopic discectomy compared with conventional techniques.^{11–14} Furthermore, microdiscectomy mostly requires muscle retraction, yellow ligament resection, and bone resection of the facet joint and/or the lamina that can lead to epidural fibrosis or segmental instability.^{11,13,14,19} These obvious potential advantages of the transforaminal endoscopic approach led us to introduce the TESSYS technique at our neurosurgical department. Our results also underline the advantages because no patient in our study population had a wound infection or CSF fistula, and the median blood loss was low. Furthermore, the median hospital stay of our study population was 4 days, which is low compared with 7 days that we found in a cohort of microdiscectomy patients we operated on during the same time interval as our study population (data not shown).

Clinical and Radiologic Outcome

Several studies reported that transforaminal endoscopic discectomy and microdiscectomy are both effective methods to treat a lumbar disk herniation, as also shown in ►Table 3.^{9,11,12,14,20–23} A large systematic review of 39 studies concluded that there are no differences between transforaminal endoscopic discectomy and microdiscectomy with regard to pain, overall improvement, patient satisfaction, recurrence rate, complications, and reoperations.⁹ However, the authors criticized the poor scientific evidence of the current literature. Our clinical results are comparable with other studies on endoscopic and microsurgical discectomy and showed a clear clinical improvement of Macnab, ODI, and VAS at 15 and 12 months after surgery, respectively. Furthermore, the overall satisfaction was 90%, 95% of the patients needed less or no analgesic medication, and 95% of the previously employed people were able to return to work, indicating a clinically effective procedure including patients with recurrent disk herniations.

We observed a statistically significant postoperative reduction of disk volume of 57.1% at the MRI volumetric analysis. Wang et al reported a residual mass of 93.6% on MRI 1 week after endoscopic discectomy.²⁴ After lumbar microdiscectomy, a rate of 80% residual epidural tissue on MRI scans on the third day after surgery was reported.²⁵ Annertz et al even described a nerve root involvement of 100% on MRI 5 days after surgery.²⁶ However, the comparability of these different studies is very limited because different methods for quantifying residual disk volume were used, and the interpretation of an early postoperative MRI can be very difficult. Residual nerve root compression is common, but the correlation with clinical symptoms is poor.^{24,25} However, our results show a relevant reduction of disk herniation volume and indicate that good clinical results can be achieved even without complete removal of the herniated disk.

Table 3 Outcome after endoscopic versus microdiskectomy: Literature review

Study	Surgical technique	Recurrent symptomatic disk herniation, %	Reoperation rate, %	Complication rate, %	Clinical outcome/Satisfaction
Endoscopic lumbar diskectomy					
Our data n = 44	Transforaminal endoscopic sequestrectomy	2.3	9.1	13.6	FU: 15 mo Macnab score: 90% excellent or good FU: 12 mo ODI reduction: 25 preop./6.5 postop. SF-8 reduction: 33 preop./21.5 postop. VAS leg pain reduction: 6 preop./2 postop. VAS back pain reduction: 6 preop./2 postop.
Ramsbacher et al ⁹ n = 39	Transforaminal endoscopic sequestrectomy	NS	10	5.1	FU: 6 wk VAS leg pain reduction: 6.7 preop./0.8 postop. VAS back pain reduction: 5.1 preop./1.3 postop.
Yeung et al ⁹ n = 307	Posterolateral endoscopic excision for lumbar disk herniation	0.7	4.6	3.9	FU: 19 mo Macnab score: 84% excellent or good
Eustacchio et al ⁹ n = 122	Endoscopic percutaneous transforaminal treatment	12	27	9	FU: 35 mo Macnab score: 45% excellent and 27% good
Morgenstern et al ⁹ n = 144	Endoscopic spine surgery	NS	5.6	9	FU: 24 mo Macnab score: 83% excellent or good
Schubert et al ⁹ n = 558	Transforaminal nucleotomy with foraminoplasty	3.6	3.6	0.7	FU: 12 mo Macnab score: 51% excellent and 43% good VAS leg pain reduction: 8.4 preop./1.0 postop. VAS back pain reduction: 8.6 preop./1.4 postop.
Ruetten et al ⁹ n = 517	Extreme-lateral transforaminal approach	6.9	6.9	0	FU: 12 mo VAS leg pain reduction: 7.1 preop./0.8 postop. VAS back pain reduction: 1.8 preop./1.6 postop. Functional status (ODI) reduction: 78 preop./20 postop.
Jang et al ⁹ n = 35	Transforaminal percutaneous endoscopic diskectomy	0	8.6	17	FU: 18 mo Macnab score 86% excellent or good
Tzaan ⁹ n = 134	Transforaminal percutaneous endoscopic lumbar diskectomy	0.7	4.5	6.0%	FU: 38 mo Macnab score: 28% excellent and 61% good
Choi et al ⁹ n = 41	Extraforaminal targeted fragmentectomy	5.1	7.7	5.1	FU: 34 mo VAS leg pain reduction: 8.6 preop./1.9 postop. Functional status (ODI) reduction: 66.3 preop./11.5 postop.

(Continued)

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

Table 3 (Continued)

Study	Surgical technique	Recurrent symptomatic disk herniation, %	Reoperation rate, %	Complication rate, %	Clinical outcome/Satisfaction
Conventional lumbar (micro)diskectomy					
Peul et al ²⁰ n = 141	Microscopic unilateral transforaminal approach	3.2	3.2	1.6	FU: 52 wk SF-36 Physical Function reduction: 33.9 preop./84.2 postop.
Weinstein et al ²¹ n = 545 (observational cohort)	Open diskectomy	7	NS	3	FU: 4 y ODI reduction: 49.3 preop./11.2 postop. SF-36 Physical Function reduction: 37.9 preop./44.6 postop. Work status: working: 84.4%
Hsu et al ¹⁴ n = 66	Open microdiskectomy	NS	6.1	1.5	FU: NS VAS reduction: 9 preop./1.3 postop ODI reduction: 32 preop./3.3 postop
Strömqvist et al ²² n = 12,840 (cohort "younger")	Lumbar diskectomy	NS	NS	5	FU: 12 mo VAS leg pain reduction: 66 preop./22 postop VAS back pain reduction: 46 preop./25 postop. ODI: 20
Gibson et al ¹¹ N = 70	Microscopic transforaminal approach	0	2.9	1.4	FU: 2 y VAS affected leg reduction: 5.8 preop./3.5 postop. VAS nonaffected leg reduction: 0.7 preop./0.8 postop. VAS back pain reduction: 4.6 preop./3.0 postop.
Liu et al ²³ n = 69	Microscopic transforaminal approach	0	0	7.2	FU: 2 y VAS leg pain reduction: 6.9 preop./1.4 postop. VAS back pain reduction: 5.8 preop./1.4 postop.

Abbreviations: FU, follow-up; NS, not specified, ODI: Oswestry Disability Index (range: 0–100), postop., postoperative; preop., preoperative; SF- 8: Short Form-8 Health Survey (range: 0–100), SF-36: Short Form-36 Health Survey (range: 0–100), VAS, visual analog scale (range: 0–10).

Transition Process and Learning Curve

The standard procedure for surgery of lumbar disk herniations at our department is a microdiskectomy using a transmuscular tubular or a subperiosteal specular approach. Residents are expected to be proficient in this procedure at the end of the German 6-year neurosurgical training program. In addition to these requirements, an increasing number of patients with uncomplicated lumbar disk herniations are now operated on in private practice, which leaves a lower overall number and more complex cases including revision surgery to university teaching hospitals like ours. All of these circumstances limit the caseload needed for the implementation of a new technique such as endoscopic diskectomy.

However, due to the various potential advantages, we decided to establish a spinal endoscopy program at our

department. One experienced spine surgeon (K.S.), who is practiced in performing microsurgical procedures, was assigned to implement the technique. After a period of observation and a cadaver course, we started with the first transforaminal endoscopic procedure supported by an experienced surgeon. Furthermore, during the following 13 operations (and during two more operations at later time points) surgery was supported by an experienced application specialist who helped improve the surgical technique and eliminate systematic errors.

We observed a median duration of surgery of 100 minutes with no significant difference between the first and the final 22 patients ($p = 0.5$), indicating a flat learning curve. Hsu and coworkers, who also described their transition process from microdiskectomy to endoscopic diskectomy (interlaminar and transforaminal route), found an operation time of 86.5

minutes in their first 57 patients that was almost twice as high (48.1 minutes) as in a parallel microdiscectomy group of patients who were operated on by the same surgeon.¹⁴ Martin-Láez et al found similar numbers during their transition from microsurgical discectomy (operation time: 66 minutes) to microendoscopic discectomy using the interlaminar approach (operation time: 100 minutes).²⁷ Challenging learning curves of endoscopic lumbar discectomy that include increased complication and reoperation rates, as well as recurrent disk herniations and the necessity of converting the procedure to microsurgery, were described by several authors.^{14,28–31}

Despite our limited caseload, we were able to implement the technique without any major complications. Minor complications occurred in six patients (13.6%) including a temporary new neurologic deficit in five patients and an early recurrent disk herniation in one patient. Temporary sensory and motor deficits are not uncommon after endoscopic discectomy, and the rate of dura perforations can increase, particularly while mastering the learning curve.^{14,27,29,32} Hsu et al found a nerve injury rate of 4.3% during their transition process; Singh et al reported new sensory deficits in 8.7% and new motor deficits in 17.4% of their patients. New deficits are found in patients with or without dura perforations, and the rates of CSF fistulas are generally low as they were in our series.¹⁴

Our complication rate is slightly higher than reported in most publications describing the transition and learning curve. However, the complication rate is naturally influenced by the number of patients in the respective series and the point of the learning curve that has been reached. Furthermore, the rates are difficult to compare due to the different endoscopic techniques and systems used. In our opinion, the highest risk for a dura and/or a neurologic injury exists during the placement of the approach rods or during reaming of the neuroforamen. In this phase, particularly if the instruments are placed too posteriorly or if the medial pedicle wall is not respected, contusion of the nerve roots can occur. Therefore, meticulous technique may help prevent neurologic deficits. Tips and tricks for beginners are offered later in this article. Operations under local anesthesia are an option to receive online patient feedback and might help prevent nerve injuries.³³ However, an awake patient places the surgeon under stress and can negatively affect the workflow, particularly during the learning curve.

In our series, a reoperation was indicated in four patients (9%) including one patient with an early recurrent disk herniation, one patient with a late recurrent disk herniation, and two patients with a new neurologic deficit and suspected postoperative hemorrhage on MRI. Several studies reported a reoperation rate in the range of 4.2 to 11% after endoscopic discectomy that can increase up to 21% during the learning curve as found by Tenenbaum et al. Thus our results are comparable with the literature.^{11,30,34–37} Cong et al found a recurrence rate of 5.04% after endoscopic discectomy in a meta-analysis, so our results with a rate of 2.3% are also comparable.¹³ A conversion to microdiscectomy was necessary in four (9%) of our patients. Similar conversion rates during the learning were reported by Joswig

et al (10%) and by Lee and Lee (7.8%).^{28,38} Reasons can be complications, reduced visibility, or challenging anatomy²⁸ as in our series. Based on these facts, we recommend informing every patient about a potential conversion to microdiscectomy until the surgeon feels comfortable with the procedure and the learning curve is complete.

Tips and Tricks for Beginners

In our opinion there are several aspects to consider during the transition process to transforaminal endoscopic discectomy:

- Adequate patient selection is an important step. We recommend starting transforaminal endoscopic surgery in patients with caudally migrated mediolateral disk herniations at the L4–L5 level or above. The trajectory to the disk herniation is easy and intuitive, and the iliac crest is not in the way.

If the surgeon feels more comfortable with the technique, disk herniations at the level L5–S1, intra- and extraforaminal herniations, and cranially migrated disk herniations can be approached.

The endoscopic technique should not be withheld from patients with recurrent disk herniations because the transforaminal approach allows the surgeon to mostly bypass the typical epidural fibrosis en route to the surgical target.

- A preoperative AP and lateral radiograph should be performed in patients with a herniation at the level L5–S1 to rule out a high iliac crest. The combination of a disk herniation at the L5–S1 disk level or a cranially migrated herniation in combination with a high iliac crest is not a case for beginners.
- We recommend an observation period and cadaver training before starting the first procedures. During the initial cases, discussion of the indication and supervision by an experienced spine surgeon is strongly advised. Also, during the first 10 to 20 cases and particularly if there are long time intervals between the operations, a follow-up visit and support of an application specialist might be helpful.
- Neurophysiologic monitoring is not helpful for a transforaminal endoscopic approach in our opinion.
- We identified three critical steps to avoid a nerve injury: (1) The small bent rod and the Seldinger wire should only just cross the midline on the AP radiograph. Slippage of these devices far to the contralateral side, which can easily happen in cases of a wide neuroforamen, places the contralateral nerve at risk. Alternatively, the small straight rod can be used and can be placed just medially to the medial pedicle wall. (2) The approach devices have to be placed in the anterior epidural space in strict vicinity to the posterior wall of the vertebral body to prevent a dura perforation. (3) The medial pedicle wall has to be meticulously respected during reaming of the neuroforamen.

Limitations

Our study had several limitations. The most important limitation is the retrospective study design with its well-

known limitations. Determination of the outcome for the retrospective cohort was only possible at discharge and by a questionnaire because most of the patients did not present to the routine follow-up examinations. Another limitation is the absence of a proper control group treated with microscopic surgery to compare both methods. However, this was not the aim of our study. Furthermore, patients were very carefully selected because we only included patients in which a transforaminal endoscopic approach seemed to be technically feasible. Especially in patients with a marked neuroforaminal stenosis, a high iliac crest, or cranially migrated disk herniations, we performed a microscopic discectomy procedure. In addition, MRI volumetric analysis was only performed in 10 patients. Nevertheless, we observed a significant reduction of disk volume herniation.

Conclusion

Implementation of the transforaminal endoscopic lumbar sequestrectomy technique in a university hospital setting is feasible and safe in selected patients with primary and recurrent disk herniations. However, several factors hinder the implementation process including the flat learning curve and the caseload that is significantly influenced by other competing team members and the residents' training requirements in microsurgery. As a result, only one surgeon can learn the technique at a time.

Conflict of Interest

Karsten Schöller reports personal fees from invited lectures at Medicon and Baxter outside the submitted work. The other authors have declared no conflicts of interest for this article.

References

- Cheng YP, Lee KW, Lin PY, et al. Full-endoscopic interlaminar removal of chronic lumbar epidural hematoma after spinal manipulation. *Surg Neurol Int* 2014;5:55
- Andrews DW, Lavyne MH. Retrospective analysis of microsurgical and standard lumbar discectomy. *Spine* 1990;15(04):329–335
- Brouwer PA, Brand R, van den Akker-van Marle ME, et al. Percutaneous laser disc decompression versus conventional microdiscectomy in sciatica: a randomized controlled trial. *Spine J* 2015;15(05):857–865
- Bokov A, Isrelov A, Skorodumov A, Aleynik A, Simonov A, Mlyavkykh S. An analysis of reasons for failed back surgery syndrome and partial results after different types of surgical lumbar nerve root decompression. *Pain Physician* 2011;14(06):545–557
- Fan G, Han R, Zhang H, He S, Chen Z. Worldwide research productivity in the field of minimally invasive spine surgery: a 20-year survey of publication activities. *Spine* 2017;42(22):1717–1722
- Jasper GP, Francisco GM, Telfeian AE. Endoscopic transforaminal discectomy for an extruded lumbar disc herniation. *Pain Physician* 2013;16(01):E31–E35
- Lühmann D, Burkhardt-Hammer T, Borowski C, Raspe H. Minimally invasive surgical procedures for the treatment of lumbar disc herniation. *GMS Health Technol Assess* 2005;1:Doc07
- Xin G, Shi-Sheng H, Hai-Long Z. Morphometric analysis of the YESS and TESSYS techniques of percutaneous transforaminal endoscopic lumbar discectomy. *Clin Anat* 2013;26(06):728–734
- Nellensteijn J, Ostelo R, Bartels R, Peul W, van Royen B, van Tulder M. Transforaminal endoscopic surgery for symptomatic lumbar disc herniations: a systematic review of the literature. *Eur Spine J* 2010;19(02):181–204
- Jasper GP, Francisco GM, Telfeian AE. Clinical success of transforaminal endoscopic discectomy with foraminotomy: a retrospective evaluation. *Clin Neurol Neurosurg* 2013;115(10):1961–1965
- Gibson JNA, Subramanian AS, Scott CEH. A randomised controlled trial of transforaminal endoscopic discectomy vs microdiscectomy. *Eur Spine J* 2017;26(03):847–856
- Pan Z, Ha Y, Yi S, Cao K. Efficacy of transforaminal endoscopic spine system (TESSYS) technique in treating lumbar disc herniation. *Med Sci Monit* 2016;22:530–539
- Cong L, Zhu Y, Tu G. A meta-analysis of endoscopic discectomy versus open discectomy for symptomatic lumbar disc herniation. *Eur Spine J* 2016;25(01):134–143
- Hsu HT, Chang SJ, Yang SS, Chai CL. Learning curve of full-endoscopic lumbar discectomy. *Eur Spine J* 2013;22(04):727–733
- Macnab I. Negative disc exploration. An analysis of the causes of nerve-root involvement in sixty-eight patients. *J Bone Joint Surg Am* 1971;53(05):891–903
- Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine* 2000;25(22):2940–2952; discussion 2952
- Price DD, McGrath PA, Rafii A, Buckingham B. The validation of visual analogue scales as ratio scale measures for chronic and experimental pain. *Pain* 1983;17(01):45–56
- Turner-Bowker DM, Bayliss MS, Ware JE Jr, Kosinski M. Usefulness of the SF-8 Health Survey for comparing the impact of migraine and other conditions. *Qual Life Res* 2003;12(08):1003–1012
- Ahn Y, Lee HY, Lee SH, Lee JH. Dural tears in percutaneous endoscopic lumbar discectomy. *Eur Spine J* 2011;20(01):58–64
- Peul WC, van Houwelingen HC, van den Hout WB, et al; Leiden-The Hague Spine Intervention Prognostic Study Group. Surgery versus prolonged conservative treatment for sciatica. *N Engl J Med* 2007;356(22):2245–2256
- Weinstein JN, Lurie JD, Tosteson TD, et al. Surgical versus nonoperative treatment for lumbar disc herniation: four-year results for the Spine Patient Outcomes Research Trial (SPORT). *Spine* 2008;33(25):2789–2800
- Strömquist B, Strömquist B, Jönsson B, Karlsson MK. The outcome of lumbar disc herniation surgery is worse in old adults than in young adults. *Acta Orthop* 2016;87(05):516–521
- Liu X, Yuan S, Tian Y, et al. Comparison of percutaneous endoscopic transforaminal discectomy, microendoscopic discectomy, and microdiscectomy for symptomatic lumbar disc herniation: minimum 2-year follow-up results. *J Neurosurg Spine* 2018;28(03):317–325
- Wang Y, Luo G, Wang J, Zhu M, Li C, Teng H. Early postoperative magnetic resonance imaging findings after percutaneous endoscopic lumbar discectomy and their correlations with clinical outcomes. *World Neurosurg* 2018;111:e241–e249
- Floris R, Spallone A, Aref TY, et al. Early postoperative MRI findings following surgery for herniated lumbar disc. *Acta Neurochir (Wien)* 1997;139(03):169–175
- Annertz M, Jönsson B, Strömquist B, Holtås S. Serial MRI in the early postoperative period after lumbar discectomy. *Neuroradiology* 1995;37(03):177–182
- Martín-Láez R, Martínez-Agüeros JA, Suárez-Fernández D, Montaña-Núñez F, Vázquez-Barquero A. Complications of endoscopic microdiscectomy using the EASYGO! system: is there any difference with conventional discectomy during the learning-curve period? *Acta Neurochir (Wien)* 2012;154(06):1023–1032
- Joswig H, Richter H, Haile SR, Hildebrandt G, Fournier JY. Introducing interlaminar full-endoscopic lumbar discectomy: a critical analysis of complications, recurrence rates, and outcome in view of two spinal surgeons' learning curves. *J Neurol Surg A Cent Eur Neurosurg* 2016;77(05):406–415

- 29 Yörükoğlu AG, Göker B, Tahta A, et al. Fully endoscopic interlaminar and transforaminal lumbar discectomy: analysis of 47 complications encountered in a series of 835 patients. *Neurocirugia (Astur)* 2017;28(05):235–241
- 30 Tenenbaum S, Arzi H, Herman A, et al. Percutaneous posterolateral transforaminal endoscopic discectomy: clinical outcome, complications, and learning curve evaluation. *Surg Technol Int* 2011;21:278–283
- 31 Morgenstern R, Morgenstern C, Yeung AT. The learning curve in foraminal endoscopic discectomy: experience needed to achieve a 90% success rate. *SAS J* 2007;1(03):100–107
- 32 Singh S, Sardhara JC, Khatri D, et al. Technical pearls and surgical outcome of early transitional period experience in minimally invasive lumbar discectomy: a prospective study. *J Craniovertebr Junction Spine* 2018;9(02):122–129
- 33 Sanusi T, Davis J, Nicassio N, Malik I. Endoscopic lumbar discectomy under local anesthesia may be an alternative to microdiscectomy: a single centre's experience using the far lateral approach. *Clin Neurol Neurosurg* 2015;139:324–327
- 34 Wilson DH, Harbaugh R. Lumbar discectomy: a comparative study of microsurgical and standard technique. In: Hardy RW, ed. *Lumbar Disc Disease*. New York, NY: Springer; 1982:147–156
- 35 Goald HJ. Microlumbar discectomy: follow-up of 477 patients. *J Microsurg* 1980;2(02):95–100
- 36 Yeung AT, Tsou PM. Posterolateral endoscopic excision for lumbar disc herniation: surgical technique, outcome, and complications in 307 consecutive cases. *Spine* 2002;27(07):722–731
- 37 Schaffer JL, Kambin P. Percutaneous posterolateral lumbar discectomy and decompression with a 6.9-millimeter cannula. Analysis of operative failures and complications. *J Bone Joint Surg Am* 1991;73(06):822–831
- 38 Lee DY, Lee SH. Learning curve for percutaneous endoscopic lumbar discectomy. *Neurol Med Chir (Tokyo)* 2008;48(09):383–388; discussion 388–389