


A. M. BERMAN, J. S. JOHNSON* (UNIVERSITY OF NORTH CAROLINA AT CHAPEL HILL, USA)


Copper-Catalyzed Electrophilic Amination of Diorganozinc Reagents

J. Am. Chem. Soc. **2004**, *126*, 5680–5681.

Electrophilic Amination of Diorganozinc Reagents Using *N*-Hydroxylamine Benzoates



Selected examples:



Significance: In 2004, Berman and Johnson developed the first electrophilic amination of *N*-hydroxylamine benzoates using diorganozinc reagents in the presence of $[\text{Cu}(\text{OTf})_2\text{C}_6\text{H}_6]$. This umpolung-type reaction allowed access to a broad range of highly functionalized secondary and tertiary amines under mild conditions.

Comment: The outstanding performance of *N*-hydroxylamine benzoates as electrophilic nitrogen sources has inspired various following publications, opening up new opportunities for amination reactions. Based on experimental studies, the authors excluded an oxidative addition/reductive elimination pathway and proposed a $\text{S}_{\text{N}}2$ -type pathway. For further insights into electrophilic aminations based on this pioneering work, see: *Synthesis* **2011**, *24*, 3954–3964.

SYNFACTS Contributors: Paul Knochel, Simon Graßl
Synfacts 2020, 16(06), 0687 Published online: 15.05.2020
DOI: 10.1055/s-0040-1706966; Reg-No.: P04820SF

© 2020, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Metals in Synthesis

Key words

electrophilic amination

organozinc reagents

copper catalysis

Synfact
Classic