

Oral Carcinoma

In Vitro Evaluation of Anticancer Activity (Efficacy) of a Novel Ayurvedic Polyherbal Formulation on Human Oral Carcinoma Cell Line SCC-40



Sandeep Charak

Sandeep Charak¹, Monika Sharma¹ Sharad M. Porte¹

¹P.G. Department of Agad Tantra, National Institute of Ayurveda, Jaipur, Rajasthan, India

South Asian J Cancer 2021;10:211–212.

We report the efficacy of our ayurvedic polyherbal formulation (V2S2) on human oral carcinoma cell line SCC-40. As you know cancer is a leading cause of death and disability and globally approximately one in six deaths is due to cancer. According to the latest World Health Organization's International Agency for Research in Cancer and the Global Cancer Observatory (GCO) report,¹ cancer burden has risen to 19.3 million new cases and 10.0 million deaths this year. Surgery, radiotherapy, chemotherapy, and nowadays immunotherapy are commonly available treatments for this disease and recent advanced research studies make these treatments so advance but the world is still struggling for the safe and effective treatment for this deadly disease. Each advanced therapy has many complications, such as chemotherapy has numerous health hazards. The present study indicates that a person dies due more to chemotherapy than cancer.¹ Thus the world is now looking toward alternative medicines like *Ayurveda* and Chinese medicine for the treatment of this disease. In *Ayurveda* the disease which can be correlated and run parallel to cancer is *Arbuda*. Herbs nowadays get so much importance in cancer therapies because of low toxicity and cost. In this study the anticancer activity of herbal extract V2S2 is studied and compared with control drug Adriamycin, which is an established chemotherapeutic drug. The experimental procedure was performed as per National Cancer Institute (NCI) guidelines and with SRB (sulforhodamine B) assay in TATA Memorial, ACTREC (Advanced Centre for Treatment, Research and Education in Cancer) Mumbai. The *Ayurvedic* drug code named V2S2 is a hydroalcoholic extract of 10 *Ayurvedic* herbs processed in a National Institute of Ayurveda pharmacy. The dried extract was then multimilled and sieved through #40 before packing in moisture-proof triple-layer aluminum pouches. The cell lines were grown in RPMI 1640 medium containing 10% fetal bovine serum and 2 mM L-glutamine. For the present

Address for correspondence Sandeep Charak, MD Ayurvedic Toxicology (Ayu.), P.G. Department of Agad Tantra, National Institute of Ayurveda, Jaipur 302002, Rajasthan, India (e-mail: dr.sandeep.charak@gmail.com).

screening experiment, cells were inoculated into 96-well microtiter plates at 100 μ L plating density.² Experimental drugs were initially solubilized in dimethyl sulfoxide at 100 mg/mL and diluted to 1 mg/mL using water and stored frozen prior to use. At the time of drug addition, an aliquot of frozen concentrate (1 mg/mL) was thawed and diluted to 100, 200, 400, and 800 μ g/mL with complete medium containing the test article. Aliquots of 10 μ L of these different drug dilutions were added to the appropriate microtiter wells already containing 90 μ L of medium, resulting in the required final drug concentrations, i.e., 10, 20, 40, and 80 μ g/mL. After compound addition, plates were incubated at standard conditions for 48 hours and assay was terminated by the addition of cold trichloroacetic acid (TCA). Cells were fixed in situ by the gentle addition of 50 μ L of cold 30% (w/v) TCA (final concentration, 10% TCA) and incubated for 60 minutes at 4°C. Percent growth was calculated on a plate-by-plate basis for test wells relative to control wells. Percent growth was expressed as (the ratio of average absorbance of the test well to the average absorbance of the control wells) \times 100. Using the six absorbance measurements [time zero (T_z), control growth (C), and test growth in the presence of drug at the four concentration levels (T_i)], the percentage growth was calculated at each of the drug concentration levels. Percentage growth inhibition was calculated as: $[T_i/C] \times 100\%$. The anticancer activity of study drug has been measured and compared with Adriamycin, measuring the percentage growth inhibition of human cell line by using a microplate reader. The drug concentration of both study and standard control drugs was used as per guidelines of NCI, such as 10, 20, 40, and 80 μ g/mL. This experiment has been repeated three times and then the average percentage of control growth was calculated and compared.

The average percentage control growth of V2S2 on human oral carcinoma cell line SCC-40 is 42.8 on 10 μ g/mL,


DOI <https://doi.org/10.1055/s-0041-1735335> ISSN 2278-330X

How to cite this article: Charak S, Sharma M, Porte S. M. In Vitro Evaluation of Anticancer Activity (Efficacy) of a Novel Ayurvedic Polyherbal Formulation on Human Oral Carcinoma Cell Line SCC-40. *South Asian J Cancer* 2021;10(3):211–212.

© 2021. MedIntel Services Pvt Ltd.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>).

Thieme Medical and Scientific Publishers Private Ltd A-12, Second Floor, Sector -2, NOIDA -201301, India



Drug concentrations
(µg/ml) calculated
from graph

SCC-40	LC50	TGI	GI50*
V2S2	NE	70.0	<10
ADR	NE	NE	<10

DEFINITIONS

LC50 = Concentration of drug causing 50% cell kill
GI50 = Concentration of drug causing 50% inhibition of cell growth
TGI = Concentration of drug causing total inhibition of cell growth
ADR = Adriamycin, Positive control compound
The residual compound with ACTREC will be retained for one month from the date of this report. Enquiries regarding report will not be entertained after this date.
NE Non- evaluable data. Experiment needs to be repeated using different set of drug concentrations.
Note: Erratic data can result due to less solubility of the compound.
GI50 value of $\leq 10^{-6}$ molar (i.e. 1 µmolar) or $\leq 10\mu\text{g/ml}$ is considered to demonstrate activity in case of pure compounds. For extracts, GI50 value $\leq 20\mu\text{g/ml}$ is considered to demonstrate activity
Yellow highlighted test values under GI50 column indicate activity.

Fig. 1 Effect of different concentrations of V2S2 and Adriamycin on human oral cancer cell line SCC-40.

64.1 on 20 µg/mL, 37.1 on 40 µg/mL, and -15.2 on 80 µg/mL. The GI50 is ≤ 10 and TGI (tumor growth inhibition) is 70.0. The average percentage control growth of Adriamycin on SCC-40 is -34.9 on 10 µg/mL, -42.4 on 20 µg/mL, -43.7 on 40 µg/mL, -26.1 on 80 µg/mL, and GI 50 is ≤ 10 . According to guidelines of NCI, a GI50 value of $\leq 10^{-6}$ molar (i.e., 1 µmolar) or $\leq 10\mu\text{g/mL}$ is considered to demonstrate activity in case of pure compounds. For extracts, GI50 value $\leq 20\mu\text{g/mL}$ is considered to demonstrate activity. In this experiment, the GI50 of Adriamycin was been found to be $\leq 10\mu\text{g/mL}$, which is a pure compound, while the GI50 of V2S2 also showed the same values while it is a hydroalcoholic extract. It means that the *Ayurvedic* coded drug equally activates on human oral cancer cell line SCC40 when compared with Adriamycin.

Thus V2S2 may be proved to be a potent anticancer drug in future without any toxicity or with minimum toxicity compared with conventional drugs.

Conflict of Interest

None.

References

- 1 Wallington M, Saxon EB, Bomb M, et al. 30-day mortality after systemic anticancer treatment for breast and lung cancer in England: a population-based, observational study. *Lancet Oncol* 2016;17(9):1203-1216
- 2 Vichai V, Kirtikara K. Sulforhodamine B colorimetric assay for cytotoxicity screening. *Nat Protoc* 2006;1(3):1112-1116