



# Elaboration and Validation of Flowchart for the Prevention and Treatment of Intestinal Peristomal Skin Complications

Imaculada Aparecida Cardoso<sup>1</sup> Geraldo Magela Salomé<sup>1</sup>   
 Adriana Rodrigues Dos Anjos Mendonça<sup>1</sup> Flávio Dutra Miranda<sup>1</sup> José Ronaldo Alves<sup>1</sup>

<sup>1</sup> Programa de Mestrado Profissional em Ciências Aplicadas à Saúde, Universidade do Vale do Sapucaí, São Paulo, SP, Brazil

Address for correspondence Geraldo Magela Salomé, RN, PhD, Sapucaí Valley University – UNIVÁS, São Paulo, SP 04330-020, Brazil (e-mail: salomereiki@yahoo.com.br).

J Coloproctol 2021;41(3):316–324.

## Abstract

**Objective** To build and validate a flowchart for the prevention and treatment of intestinal peristomal skin complications.

**Method** For the construction of the algorithms, a systematic review was carried out in health sciences databases comprising the last 10 years. The evaluation of the algorithms was carried out by 38 nurses. For the validation of the algorithms, the Delphi technique was used. The statistical analysis used was the content validity index and the Cronbach alpha coefficient. The questionnaire was sent by e-mail and in person after approval by the Ethics and Research Committee.

**Results** In the first evaluation of the algorithms, there was no agreement among the experts. However, after making the corrections suggested by the evaluators, the algorithms were resent, with a 100% consensus among the evaluators. The questions used to validate the algorithms contributed favourably to the internal consistency and content validation of the instrument, since the respective Cronbach alpha was 0.9062 and the global content validity index (g-CVI) was 0.91 in the first validation and 1.0 in the second validation.

**Conclusion** After an integrative literature review, the flowcharts were built and validated by a professional with experience in the area, showing 100% agreement among the experts in the second evaluation.

## Keywords

- ▶ stoma
- ▶ dermatitis
- ▶ algorithms
- ▶ mobile apps

## Introduction

In the 21<sup>st</sup> century, sociodemographic, economic, and epidemiological changes caused by accelerated urbanization, new consumption patterns, globalization of knowledge, scientific and technological advances have had a major national impact, with a decline in the birth rate, increased life

expectancy, and, consequently, of longevity associated in parallel with the increase of chronic noncommunicable diseases, a high number of traffic accidents, and an increase in urban violence, resulting in a growing increase in the number of people with stomas in Brazil. Approximately 1.4 million stoma-making surgical procedures are performed every year.<sup>1</sup>

received  
August 25, 2020  
accepted  
January 15, 2021

DOI <https://doi.org/10.1055/s-0041-1735460>.  
ISSN 2237-9363.

© 2021. Sociedade Brasileira de Coloproctologia. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (<https://creativecommons.org/licenses/by-nc-nd/4.0/>)

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

The purpose of this surgery is the diversion of the contents of the intestine (gases and faeces) to an external bag.<sup>2</sup> This procedure is carried out so that the elimination function is maintained, and it causes several changes, among which can be highlighted the elimination of gases, odour and faeces by the stoma located in the abdominal wall.

According to the 2010 census of the Brazilian Institute of Geography and Statistics, when the total population of Brazil was 190,732,694 people, it was estimated that there were ~ 190,000 ostomy patients in Brazil, and 120,000 stoma creation surgeries annually in the United States.<sup>3</sup>

Between 21 and 70% of ostomized patients develop some type of complication, although a significant percentage of these patients could live with the stoma without complications. These complications are related to the malfunction of the ostomy, to demarcation in an incorrect anatomical region, and to the difficulty of the patient in performing self-care.<sup>4</sup> Such complications can lead to dermatitis, bleeding, prolapse, necrosis, hernias, edema, waste leakage, peristomal hernia, stenosis, and retraction, among other adversities.<sup>5-11</sup>

Nurses have a great responsibility in the care of the ostomate, since they participate in all stages of care. In the preoperative period, it is recommended that the stoma location be demarcated by the stoma nurse, starting to teach the patient and family about the surgery and its consequences, as well as identifying other learning needs. In the postoperative period, self-care teaching is resumed in relation to the stoma, indication and replacement of equipment, and the patient is prepared for hospital discharge.<sup>9,12-15</sup>

Health professionals who provide assistance to ostomized patients should always seek new knowledge, as scientific updates and advances in technologies related to the care of intestinal peristomal skin are constantly evolving. It is not difficult to find professionals who work in the hospital and outpatient setting who have difficulties in carrying out the exchange of the bag, in hygienizing the intestinal peristomal skin, in identifying the ideal bag for each patient, and in identifying the type of stoma. In addition, the evaluation of a stoma can lead to different interpretations due to its diversity in terms of nature, shape and location, depending on each professional's own perception and technical knowledge, considering the subjective factors of the evaluation.<sup>16-19</sup>

Algorithms are technologies that guide decision-making in the face of clinical care issues, add scientific rationality, and serve as guides for self-care, prevention and treatment of peristomal skin complications. They provide information about the best prophylactic-therapeutic conduct to be adopted in each clinical evaluation made by the nursing and multiprofessional team, which confirms its proficiency and accuracy as a guiding instrument of care. The elaboration and structuring of the algorithm must be composed by the evaluation of the algorithm by a professional with knowledge in the area, care actions, and therapeutic proposal.<sup>17-20</sup>

Thus, it was essential to develop protocols in the form of algorithms that make the clinical management of ostomy patients more effective, as well as minimizing difficulties and deficiencies of the nurses in relation to clinical practice. With

this, it is expected to contribute to the dissemination of knowledge about the evaluation of intestinal peristomal skin, promoting the health of this population, as well as assisting professionals in the elaboration of preventive actions, therapeutic conduct, and health education actions.

The present study aimed to build and validate flowcharts for the prevention and treatment of intestinal peristomal skin complications.

## Methods

A methodological study on the development of a technological application.

For the elaboration of the algorithms, an integrative literature review was carried out in the following health sciences databases: Scientific Electronic Library Online (SciELO), Latin American and Caribbean Literature in Health Science (LILACS), and National Library of Medicine – USA (MEDLINE). For the search, the descriptors *stoma* and *dermatitis* were used.

For the selection of publications, the following inclusion criteria were adopted: only primary studies that had a direct connection with the theme; full text available and no time limitation proposed, as the intention was to compile all studies that met the established criteria. The following were excluded: book chapters, theses, dissertations, monographs, technical reports, reference works, and articles that, after reading the abstract, did not converge with the proposed study object.

The articles were classified according to the level of evidence in: 1, when the evidence came from a systematic review or meta-analysis of all randomized controlled clinical trials, relevant or derived from clinical guidelines based on systematic reviews of randomized controlled clinical trials; 2, if the evidence was derived from at least one well-designed randomized controlled clinical trial; 3, when the evidence was obtained from well-designed clinical trials without randomization; 4, evidence from cohort and well-designed case control studies; 5, evidence from a systematic review of descriptive and qualitative studies; 6, evidence derived from a single descriptive or qualitative study; 7, evidence originating from the opinion of authorities and/or the report of expert committees.<sup>21</sup>

After reading the abstracts, articles that described the following subjects were selected: evaluation of intestinal peristomal skin, care of intestinal peristomal skin, prevention of intestinal peristomal skin, and treatment of complications of intestinal peristomal skin. From this survey, the algorithms for the prevention and treatment of intestinal peristomal skin complications were developed in three stages.

### First stage: Construction of the Flowcharts for the Prevention of Intestinal Peristomal Skin Injuries

**First step** – In this phase, a standardization was elaborated to carry out anamnesis and physical examination.

**Second step** – At this stage, a plan was developed with preventive measures and actions for self-care, in case the patient does not present signs and symptoms of complications in the intestinal peristomal skin.

### **Second Stage: Elaboration of Flowcharts for the Treatment of Intestinal Peristomal Skin Complications**

**First step:** At this stage, the types of dermatitis were classified, exposing signs and symptoms present in each type of dermatitis and its causes based on the reviewed literature.

**Second step:** Standardization of therapeutic approaches to each dermatitis: at this stage, a plan of therapeutic approaches based on the reviewed literature for each type of dermatitis found was developed.

### **Third Stage: Validation of the Algorithms**

The present study was carried out at the Hospital das Clínicas Samuel Libânio, at the Universidade do Vale do Sapucaí, São Paulo, state of São Paulo, Brazil. The study population consisted of 38 nurses. The study was approved by the Research Ethics Committee of the Faculty of Medical Sciences Dr. José Antônio Garcia Coutinho of the Universidade do Vale do Sapucaí, under the consubstantiated opinion number 2.557.232.

The inclusion criteria of the evaluators were: to be a professional holding a certificate of an undergraduate nursing course, with at least 1 year of experience in treating patients with skin lesions. The exclusion criteria were professionals who agreed to participate in the research but did not answer neither/nor submitted the evaluation questionnaire within 15 days.

For the validation of the algorithms, the following documents were elaborated and sent to the research participants:

The invitation letter, which was composed of: initial personal presentation and clarifications on the research topic, with an objective definition regarding the ostomy, opinion of the Research Ethics Committee of the Faculty of Health Sciences Dr. José Antônio Garcia Coutinho, and explanations about the importance of the professional evaluator in the research. The step by step for the effective participation of the evaluators, as well as the period of 15 days for each evaluation round, counting from the day of delivery, to elaborate and forward the answers, were also included in the letter.

The informed consent form (ICF) made the content of the research clear to the evaluator, guaranteeing the confidentiality of personal information and their free decision whether or not to participate in it, in addition to the knowledge of the right to withdraw, at any time, their consent to participate in the research. In that term, in case of acceptance, name and profession of the participant were requested.

The professionals evaluated the following items of the algorithms: thematic content, graphic presentation, vocabulary, sequence, clarity and understanding of the information, description of the physical examination and anamnesis of the peristomal skin, description of the peristomal skin care, peristomal skin cleaning technique, and replacement of

pouching systems of one and two pieces, classification of dermatitis, its causes, signs, and symptoms, and appropriate therapeutic procedures, when evidenced.

The Likert scale was used in the questions to evaluate the algorithms, with the following answer options: “appropriate”, “totally appropriate”, “not appropriate”, “partially appropriate”, and “not applicable”. In all items, there was room so that the participants could insert their own opinions and suggestions.

The process of validating the algorithms went as follows:

1. Selection of professional evaluators: the e-mails or personal contacts of the research participants registered in the institutions mentioned above were verified.
2. Sending the instrument to the evaluators by e-mail in a simple way, as an attachment, or delivered by hand in printed form. The participants read and signed the ICF. Freely agreeing to participate in the survey, they provided their full name, profession, and individual taxpayer registration (CPF, in the Portuguese acronym). Those who received the instrument electronically, printed, signed, replied, scanned and sent it.
3. In the analysis of the data, the answers marked with classification 3 (appropriate) or 4 (totally appropriate) were considered valid. Responses rated 1 (inappropriate) or 2 (partially appropriate) were not excluded. The suggestions made by the experts were evaluated and revisions were made so that the items were considered validated, and these guidelines were indicated in previous studies on this evaluation method.<sup>23</sup> Questions that received ratings 1 (inappropriate) or 2 (partially appropriate) were sent to the participants in the second round of evaluation with suggestions made for a new round, reaching a consensus of 100% of approval among the evaluators; this type of procedure is known as the Delphi technique.

The Delphi technique is a method that obtains opinions from evaluators with specific knowledge in a certain area. The Delphi technique uses questionnaires whose contents are analysed and judged by specialists seeking a 100% consensus among the evaluators. Generally, there are two to three rounds or evaluation cycles, but there may be more.<sup>23</sup>

This is a result counting technique depending on the degree of expertise, without specifying the number of judges. This technique motivates the evaluators to think more about the subject, as they will be the creators of the theme in question. For this process, two groups were used: an executor group, which was composed by the researchers, whose function was to contact the respondents, to prepare the initial questionnaire, to analyse the data, and to elaborate the other questionnaires; and the group formed by the selected experts.<sup>24</sup>

The obtained data were electronically tabulated with the aid of Microsoft Excel 2010 (Microsoft Corporation, Redmond, WA, USA) and were subsequently analysed quantitatively using the same program.

The data were analysed and correlated to determine the Cronbach alpha coefficient, which aims to estimate the

reliability of the instrument. The higher the covariance or correlation between the items, that is, the closer the value obtained is to one, the greater the homogeneity of the items and the consistency with which they measure the same dimension or theoretical construct. A Cronbach's alpha coefficient  $> 0.7$  indicates a good internal consistency of the instrument.

To validate the content of the algorithms, the content validity index (CVI) was used, whose purpose was to measure the proportion or percentage of participants who agree on certain aspects of the instrument and its items. The CVI is a widely used method in the health field. The CVI uses a Likert scale with a certain number of points for concordance and representativeness. Answers may vary, for example, from relevant to unrepresentative, or from clear to unclear.<sup>25</sup> For the present study, the questionnaire contained 5 alternative answers with an answer for each question, as follows: 1 = Inappropriate (I); 2 = Partially Appropriate (PA); 3 = Appropriate (A); 4 = Totally Appropriate (TA). In all items, there was room so that the experts could insert their own opinions and suggestions.

For validation, a quantitative assessment was carried out item by item in the questionnaire. The CVI was calculated considering the number of responses "3" (Appropriate) or "4" (Totally Appropriate) for each item divided by the total number of responses. The value of the CVI for the validation of a questionnaire must be  $\geq 0.78$ , when there is the participation of  $\geq 6$  validation specialists.<sup>25</sup>

For the validation of the instrument as a whole, there is no consensus between the different formulas. In the present study, we chose to use the following calculation: the sum of all CVIs of each item calculated separately divided by the number of items that were considered in the questionnaire evaluation, with a minimum mandatory agreement  $> 0.90$ .<sup>25</sup>

## Results

► **Table 1** shows the evaluation of the algorithms using the Delphi technique. In the first evaluation, the experts evaluated each question of the algorithms between inappropriate and totally appropriate; however, in the second evaluation, the algorithms obtained the evaluation of appropriate and totally appropriate.

► **Table 2** shows that the mean Cronbach alpha was 0.9062. By relating the questions of the algorithms to  $\alpha$ , it varied from 0.8916 to 0.9100, characterizing that the questions contained in the algorithms have excellent internal reliability.

► **Table 3** makes it possible to observe that, in the first evaluation, the CVI of the questions varied between 0.89 and 0.97 and that the g-CVI was 0.91. After the corrections requested by the participants, the algorithms were reevaluated, with the CVI of all questions being 1.0 and the global content validity index (g-CVI) was 1.0. These results demonstrate that the experts agree regarding the content of the algorithms.

## Discussion

The descriptive protocols or in form of algorithms must be developed based on scientific evidence, as they have the purpose of guiding professionals, patients, companions, and caregivers in decision-making so that they provide care free of damages, adverse events, and with minimum possible risk.<sup>17,18,26</sup>

The present study built two algorithms (► **Figures 1 and 2**) that will serve as a clinical tool in decision-making in the assessment, prevention and therapeutic approaches related to intestinal peristomal skin complications.

The use of protocols in algorithm format in clinical practice contributes to the individualized registration of care and systematic assessment, enabling the continuity of preventive measures and of the treatment, and promoting the quality of care. The algorithms must be built on a scientific basis and after a review of the literature on the topic.<sup>27</sup> The systematic evaluation of the peristomal skin minimizes the healing time and allows analysis of the costs and benefits of the treatment used.<sup>17</sup>

The construction of an algorithm for the evaluation of injuries must be strongly based on the literature and on clinical evidence to provide technological, technical, clinical, administrative, and financial subsidies, always aiming at improving patient care and at the best results for the institution.<sup>17-19</sup>

The content of the algorithms developed in the present research was evaluated by nurses using the Delphi technique. In the first evaluation, most of the evaluators considered the content of the algorithms as inappropriate to appropriate. However, after making the corrections that were suggested by the evaluators, the algorithms were evaluated as appropriate to totally appropriate, with a Cronbach alpha of 0.9062. These findings characterize that the questionnaire used to validate the content of the algorithms has good internal consistency and excellent reliability, corroborating the findings of other authors.<sup>17-19,25</sup>

Several studies that validated the content of the protocols, booklets, apps, and algorithms using the Delphi technique report that the corrections suggested by the evaluators must be made on a scientific basis, as they contribute to a better understanding, effectiveness, and implantation of the material in the institution, allowing the professional to choose the best preventive measure and therapeutic approaches, resulting in safe assistance, without damage and with the least possible risk and reduction in the cost of treatment.<sup>17-19,28</sup>

Regarding the validation of the content of the algorithms, in the first evaluation, the g-CVI was 0.91 and, in the second evaluation, the g-CVI was 1.0. These results demonstrate agreement among the experts only in the second evaluation, and they coincide with the results found by other authors.<sup>17-19,28</sup>

Protocols, algorithms, booklets, manuals, flowcharts, and guidelines are considered important tools for coping with various problems in the care and management of health services. Studies validated by scientific evidence are based on guidelines of a technical, organizational, and political nature.

**Table 1** Evaluation of the content of the items of the algorithms for the prevention and treatment of peristomal skin complications using the Delphi technique. Pouso Alegre, MG, Brasil, 2019

Delphi Technique										
First evaluation										
Questions	Inappropriate		Partially appropriate		Appropriate		Totally appropriate		Total	
	n	%	n	%	n	%	n	%	N	%
Regarding Algorithm's design	00	00	03	07.90	16	42.10	19	50	38	100
Regarding Algorithm's content	00	00	03	07.90	14	36.80	21	55.30	38	100
Regarding Algorithm's sequence	00	00	04	10.50	10	26.30	24	63.20	38	100
Regarding Algorithm's vocabulary	00	00	03	07.90	11	28.90	24	63.2	38	100
Regarding the understanding of the Algorithm information	00	00	03	07.90	10	26.30	25	65.80	38	100
Regarding the description of the physical examination and anamnesis of the peristomal skin	00	00	03	07.90	15	39.50	20	52.60	38	100
Regarding the description of skin care around the stoma	00	00	05	13.20	08	21.00	25	65.80	38	100
Regarding the technique for cleaning peristomal skin	01	2.60	04	10.50	12	31.60	21	55.30	38	100
Regarding the description of the technique of changing an one-part pouching system	01	2.60	04	10.50	12	31.60	21	55.30	38	100
Regarding the description of the technique of changing a two-part pouching system	00	00	00	01	02.60	16	42.10	21	55.30	38
Regarding dermatitis classification	00	00	01	02.60	16	42.10	21	55.30	38	100
Regarding the description of the causes of dermatitis	00	00	02	05.30	14	36.80	22	57.90	38	100
Regarding the description of the signs and symptoms of dermatitis	01	2.6	03	07.90	10	26.30	24	63.20	38	100
Regarding the description of therapeutic approaches	00	00	004	10.50	18	47.40	16	42.10	38	100
Delphi Technique										
Second evaluation										
Questions	Inappropriate		Partially appropriate		Appropriate		Totally appropriate		Total	
	n	%	n	%	n	%	n	%	n	%
Regarding Algorithm's design	00	00	00	00	13	34.20	25	65.80	38	100
Regarding Algorithm's content	00	00	00	00	08	21.10	30	78.90	38	100
Regarding Algorithm's sequence	00	00	00	00	09	23.70	29	76.30	38	100
Regarding Algorithm's vocabulary	00	00	00	00	09	23.70	29	76.30	38	100
Regarding the understanding of the Algorithm information	00	00	00	00	08	21.10	30	78.90	38	100
Regarding the description of the physical examination and anamnesis of the peristomal skin	00	00	00	00	11	28.90	27	71.10	38	100
Regarding the description of skin care around the stoma	00	00	00	00	05	13.20	33	86.80	38	100
Regarding the technique for cleaning peristomal skin	00	00	00	00	08	21.10	30	78.90	38	100
Regarding the description of the technique of changing an one-part pouching system	00	00	00	00	10	26.30	28	73.70	38	100
Regarding the description of the technique of changing a two-part pouching system	00	00	00	00	10	26.30	28	73.70	38	100
Regarding dermatitis classification	00	00	00	00	09	23.70	29	76.30	38	100
Regarding the description of the causes of dermatitis	00	00	00	00	09	23.70	29	76.30	38	100
Regarding the description of the signs and symptoms of dermatitis	00	00	00	00	08	21.10	30	78.90	38	100
Regarding the description of therapeutic approaches	00	00	00	00	10	26.30	28	73.70	38	100

They also focus on the standardization of clinical, surgical, and preventive procedures.<sup>27,29-32</sup>

The algorithms developed in the present study provide relevant information about the best therapeutic and preven-

tive direction to be adopted, guiding health professionals more safely in the evaluation and decision-making process before the ostomate who needs the intervention to prevent and treat intestinal peristomal skin complications, which

**Table 2** Internal consistency of items related to the algorithms. Pouso Alegre, MG, Brasil, 2019

Questions presented in the algorithm validation questionnaire	Cronbach alpha
Regarding Algorithm's design	*0.9089
Regarding Algorithm's content	*0.9073
Regarding Algorithm's sequence	*0.8937
Regarding Algorithm's vocabulary	*0.9059
Regarding the understanding of the algorithm information	*0.9100
Regarding the description of the physical examination and anamnesis of the peristomal skin	*0.90140
Regarding the description of skin care around the stoma	*0.8996
Regarding the technique for cleaning peristomal skin	*0.8940
Regarding the description of the technique of changing an one-part pouching system	*0.8916
Regarding the description of the technique of changing a two-part pouching system	*0.8994
Regarding dermatitis classification	*0.8943
Regarding the description of the causes of dermatitis	*0.8933
Regarding the description of the signs and symptoms of dermatitis	*0.8953
Regarding the description of therapeutic approaches	*0.8977
Cronbach alpha coefficient	*0.9062

Cronbach alpha test. \* Significance level  $\alpha > 0.800$ .

**Table 3** Content validity index of the questions presented in the algorithms for the prevention and treatment of peristomal skin complications by the research participants. Pouso Alegre, MG, Brasil, 2019


Questions	Content Validity Index	
	CVI before correction	CVI after correction
Regarding Algorithm's design	*0.92	*1.0
Regarding Algorithm's content	*0.92	*1.0
Regarding Algorithm's sequence	*0.89	*1.0
Regarding Algorithm's vocabulary	*0.92	*1.0
Regarding the understanding of the algorithm information	*0.92	*1.0
Regarding the description of the physical examination and anamnesis of the peristomal skin	*0.92	*1.0
Regarding the description of skin care around the stoma	*0.87	*1.0
Regarding the technique for cleaning peristomal skin	*0.87	*1.0
Regarding the description of the technique of changing an one-part pouching system	*0.89	*1.0
Regarding the description of the technique of changing a two-part pouching system	*0.89	*1.0
Regarding dermatitis classification	*0.97	*1.0
Regarding the description of the causes of dermatitis	*0.95	*1.0
Regarding the description of the signs and symptoms of dermatitis	*0.89	*1.0
Regarding the description of therapeutic approaches	*0.92	*1.0
<b>Global Content Validity Index (g-CVI)</b>	<b>**0.91</b>	<b>**1.0</b>

Abbreviation: CVI, content validity index.

Content validity index. \*Significance g-CVI > 0.78.

Content validity index. \*\*Significance g-CVI > 0.90.





**Fig. 1** Algorithm for the Prevention of Peristomal Skin Complications. Pouso Alegre, MG, Brasil, 2019.




Fig. 2 Algorithm for the Treatment of Peristomal Skin Complications. Pouso Alegre, MG, Brasil, 2019.

results in a better quality of life for the patients. Algorithms are modern ways of information that contribute to future advances in the provision of work content, as well as protocols and palliative care.

Therefore, concluding, it is possible to affirm that after an integrative literature review, the algorithms were built and validated by professionals with experience in the area, showing 100% agreement among the experts in the second evaluation.

**Conflict of Interests**

The authors have no conflict of interests to declare.

**References**

1 Lenza NFB, Sonobe HM, Buetto LS, Santos MG, Lima MS. O ensino do autocuidado aos pacientes estomizados e seus familiares: uma revisão integrativa. Rev Bras Promoç Saúde 2013;26(01): 139-145



- 2 Santos AC, Dutra RAA, Salomé GM, Ferreira LM. Construction and internal reliability of an algorithm for choice cleaning and topical therapy on wounds. *J Nurs UFPE online* 2018;12(05): 680–85
- 3 Instituto Brasileiro de Geografia e Estatística. <https://www.ibge.gov.br/calendario-de-divulgacoes-novoportal.html>2010. [cited June 9<sup>th</sup>, 2019].
- 4 Soares CS, Santos I, Berardinelli LMM. Obesity as a social problem: identifying guidance needs of nursing for self-care. *Rev Enferm UFPE on line* 2010;4(01):18–27. Doi: 10.5205/reuol.520-5592-3-LE.0401201003
- 5 Salome GM, Almeida SA. Association of sociodemographic and clinical factors with the self-image and self-esteem of individuals with intestinal stoma. *J Coloproctol* 2014;4(03):159–66
- 6 Salomé GM, Almeida SA, Mendes B, Carvalho MRF, Massahud. Assessment of subjective well-being and quality of life in patients with intestinal stoma. *J Coloproctol* 2015;35(03):168–174. Doi: 10.1016/j.jcol.2015.03.002
- 7 Lima JA, Muniz KC, Salomé GM, Ferreira LM. Association of socio-demographic and clinical factors with self-image, self-esteem and locus of health control in patients with an intestinal stoma. *J Coloproctol* 2018;38(01):56–64. Doi: 10.1016/j.jcol.2017.11.003
- 8 Moreira CNO, Marques CB, Salomé GM, Cunha DR, Pinheiro FAM. Health locus of control, spirituality and hope for healing in individuals with intestinal stoma. *J Coloproct* 2016;36(04): 208–215. Doi: 10.1016/j.jcol.2016.04.013
- 9 Perissotto S, Breder JSC, Zulian LR, de Oliveira VX, da Silveira NI, Alexandre NMC. Ações de enfermagem para prevenção e tratamento de complicações em estomias intestinais: revisão integrativa. *Estima, Braz J Enterostomal Ther* 2019;17:e0519. Doi: 10.30886/estima.v17.638\_PT
- 10 Russell S. Exercise after ostomy surgery and peristomal hernia: a view from here. *J Wound Ostomy Continence Nurs* 2019;46(03): 215–218. Doi: 10.1097/WON.0000000000000521
- 11 Nunes MLG, Santos VLGC. Instrumentos de avaliação das complicações da pele peristoma: revisão integrativa. *Aquichan* 2018;18(02):477–491. Doi: 10.5294/aqui.2018.18.4.9
- 12 Somayaji R. Delving into skin soft tissue infections (SSTI) part I: an overview of infection. *WCET Journal* 2016;36(02):29–34. Doi: 10.3316/informit.095716017578970
- 13 Antonini M, Militello G, Manfreda S, Arena R, Veraldi S. A revised version of the SACS Scale for peristomal skin disorders classification. *WCET Journal*. 2016;36(03):22–29. Doi: 10.3316/INFORMIT.361794847146066
- 14 Salomé GM, Carvalho MRF, Massahud MR Junior, Mendes B. Profile of ostomy patients residing in Pouso Alegre city. *J Coloproctol* 2015;35(02):106–112. Doi: 10.1016/j.jcol.2015.02.002
- 15 Salomé GM, de Almeida SA, de Jesus Pereira MT, et al. The Impact of Venous Leg Ulcers on Body Image and Self-esteem. *Adv Skin Wound Care* 2016;29(07):316–321
- 16 Salomé GM, Santos LF, Cabeceira HS, Panza AMM, Paula MAB. Knowledge of undergraduate nursing course teachers on the prevention and care of peristomal skin. *J Coloproctol* 2014;34(04):224–230. Doi: 10.1016/j.jcol.2014.05.008
- 17 Carvalho MRF, Salomé GM, Ferreira LM. Construção e validação de algoritmo para tratamento da lesão por pressão. *Rev Enferm UFPE on line* 2017;11(Suppl 10):2533–2540
- 18 Cunha DRD, Salomé GM, Massahud MR, Mendes B, Ferreira LM. Development and validation of an algorithm for laser application in wound treatment. *Rev Lat Am Enfermagem* 2017;25:e2955
- 19 Cunha JB, Dutra RAA, Salomé GM. Elaboration of an algorithm for wound evaluation and treatment. *Estima, Braz J Enterostomal Ther* 2018;16:e2018. Doi: 10.30886/estima.v16524
- 20 van Rijswijk L, Beitz JM. Pressure ulcer prevention algorithm content validation: a mixed-methods, quantitative study. *Ostomy Wound Manage* 2015;61(04):48–57
- 21 Melnyk BM, Fineout-Overholt E. Evidence-based practice in nursing & healthcare: a guide to best practice. Philadelphia: Lippincott Williams & Wilkins; 2005
- 22 Grant JS, Davis LL. Selection and use of content experts for instrument development. *Res Nurs Health* 1997;20(03):269–274
- 23 Cassiani SH, Rodrigues LP. A técnica de Delphi e a técnica de grupo nominal como estratégias de coleta de dados das pesquisas em enfermagem. *Acta Paul Enferm* 1996;9(3):76–83
- 24 Brandão ES, Santos I, Lanzillotti RS. Validação de um instrumento para avaliação do cliente com afecções cutâneas. *Acta Paul Enferm* 2013;26(05):460–466. Doi: 10.1590/S0103-21002013000500009
- 25 Wynd CA, Schmidt B, Schaefer MA. Two quantitative approaches for estimating content validity. *West J Nurs Res* 2003;25(05):508–518
- 26 Catunda HLO, Bernardo EBR, Vasconcellos CTM, Moura ERF, Pinheiro AKB, Aquino OS. Methodological approach in nursing research for constructing and validating protocols. *Texto Contexto Enferm* 2017; 26(02):e00650016. Doi: 10.1590/0104-07072017000650016
- 27 Pott FS, Ribas JD, Silva OBM, et al. Algoritmo de prevenção e tratamento de úlcera por pressão. *Cogitare Enferm*. 2013;18(02): 23844. Doi: 10.5380/ce.v18i2.26085
- 28 Galvão EC F, Puschel VA, Galvão ECF, Puschel VAA. A Multimídia application in mobile platform for teaching the measurement of central venous pressure. *Rev Esc Enferm USP* 2012;46;(n.spe):107–115. Doi: 10.1590/s0080-62342012000700016
- 29 Gonçalves MB, Rabeb SAN, Nogueira PC. Revisão - Terapia tópica para ferida crônica: recomendações para prática baseada em evidências. *Estima [Internet]* 2015;12(1):42–49. Available from: <https://www.revistaestima.com.br/estima/article/view/337>
- 30 Johnson JJ, Paustrian C. Guideline for management of wounds in patients with lower-extremity venous disease [Internet]. Glenview: Wound Ostomy and Continence Nurses Society-WOCN; 2005. [cited jun 09, 2019]. Available from: <http://www.wocn.org/news/67120/Guideline-for-Management-of-Wounds-in-Patients-with-Lower-Extremity-Venous-Disease-Available-Now.htm>
- 31 Mendes B, Salomé GM, Pinheiro FAM, Massahud MR Júnior, da Cunha DM, Ferreira LM. Prevention and treatment of the trench foot: validation of a teaching manual for military personnel. *J Wound Care* 2018;27 (Sup10): S33–S38
- 32 Salomé GM, Cunha AL, Pereira AP, Miranda FD, Alves JR. Educational handbook for healthcare professionals: Preventing complications and treating peristomal skin. *J Coloproctol* 2019;39(04):332–338. Doi: 10.1016/j.jcol.2019.07.005