

R. B. WOODWARD*, F. E. BADER, H. BICKEL, A. J. FREY, R. W. KIERSTEAD (HARVARD UNIVERSITY, CAMBRIDGE, USA)

The Total Synthesis of Reserpine


J. Am. Chem. Soc. **1956**, 78, 2023–2025, DOI: 10.1021/ja01590a079.

R. B. WOODWARD*, F. E. BADER, H. BICKEL, A. J. FREY, R. W. KIERSTEAD (HARVARD UNIVERSITY, CAMBRIDGE, USA)

The Total Synthesis of Reserpine

Tetrahedron **1958**, 2, 1–57, DOI: 10.1016/0040-4020(58)88022-9.

Total Synthesis of (±)-Reserpine



Significance: (±)-Reserpine was isolated from *Rauwolfia serpentina*, a flower from the *Apocynaceae* family. Today it is used as a drug for the treatment of high blood pressure. The configuration was elucidated in 1955 by the Ciba group. In 1956, Woodward and co-workers disclosed the first total synthesis, which completed (±)-reserpine in 21 steps.

Comment: Lactone **D** is obtained through a Diels–Alder reaction of quinone **A** with diene **B** and subsequent reduction. Functional group interconversions lead to aldehyde **I** bearing five stereocenters. Reductive amination of **I** and indole derivative **J**, followed by lactamization forms **K**. Subsequent Bischler–Napieralski reaction completes the carbon skeleton of (±)-reserpine.

SYNFACTS Contributors: Erick M. Carreira, Willi M. Amberg
Synfacts 2022, 18(05), 0473 Published online: 20.04.2022
DOI: 10.1055/s-0041-1738025; Reg-No.: C021225F

© 2022, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Synthesis of Natural Products and Potential Drugs

Key words

(±)-reserpine

indole alkaloids

Meerwein–Ponndorf–Verley reduction

Jones oxidation

Bischler–Napieralski reaction

Steglich esterification

Synfact Classic

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.