

Natural Indole Alkaloids from Marine Fungi: Chemical Diversity and Biological Activities

Jiao Li¹ Chun-Lin Zhuang^{2,3,*}


¹ Clinical Medicine Scientific and Technical Innovation Center, Shanghai Tenth People's Hospital, Tongji University School of Medicine, Shanghai, People's Republic of China

² Department of Natural Product Chemistry, School of Pharmacy, The Second Military Medical University, Shanghai, People's Republic of China

³ Department of Medicinal Chemistry, School of Pharmacy, Ningxia Medical University, Yinchuan, People's Republic of China

Address for correspondence Chun-Lin Zhuang, PhD, School of Pharmacy, The Second Military Medical University, 325 Guohe Road, Shanghai 200433, People's Republic of China (e-mail: zclnathan@163.com).

Pharmaceut Fronts 2021;3:e139–e163.



received
August 31, 2021
accepted
October 14, 2021

DOI <https://doi.org/10.1055/s-0041-1740050>.
ISSN 2628-5088.

© 2021. The Author(s).
This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (<https://creativecommons.org/licenses/by/4.0/>)
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Abstract

The indole scaffold is one of the most important heterocyclic ring systems for pharmaceutical development, and serves as an active moiety in several clinical drugs. Fungi derived from marine origin are more liable to produce novel indole-containing natural products due to their extreme living environments. The indole alkaloids from marine fungi have drawn considerable attention for their unique chemical structures and significant biological activities. This review attempts to provide a summary of the structural diversity of marine fungal indole alkaloids including prenylated indoles, diketopiperazine indoles, bisindoles or trisindoles, quinazoline-containing indoles, indole-diterpenoids, and other indoles, as well as their known biological activities, mainly focusing on cytotoxic, kinase inhibitory, antiinflammatory, antimicrobial, anti-insecticidal, and brine shrimp lethal effects. A total of 306 indole alkaloids from marine fungi have been summarized, covering the references published from 1995 to early 2021, expecting to be beneficial for drug discovery in the future.

Keywords

- ▶ marine fungi
- ▶ indole alkaloids
- ▶ structural diversity
- ▶ biological activity

Introduction

The indole fragment is a valuable unit in a wide range of clinical drugs for treating various diseases, such as sunitinib (anticancer), nintedanib (anti-idiopathic pulmonary fibrosis), reserpine (antihypertension), indomethacin (antiinflammation), amedalin (antidepression), atevirdine (anti-human immunodeficiency virus), zafirlukast (asthma), etc. (▶ Fig. 1).¹⁻⁷ This ring system is one of the most important heterocycles for pharmaceutical development^{8,9} and widely distributed in bioactive heterocyclic natural products.¹⁰ The marine fungi are a rich underexploited source to produce novel indole-containing secondary metabolites for drug discovery, due to their extreme marine living conditions.¹¹⁻¹⁴ Thus, the indole alkaloids from marine fungi have drawn considerable attention for their unique chemical

structures and significant biological activities.^{13,15,16} In the light of the increasing attention paid on the marine fungal indoles, it is necessary to give a comprehensive summary on these indoles from the specific source. Herein, we reviewed the chemical diversity and biological properties of marine fungal indole alkaloids, expecting to provide clear evidence that these metabolites possess potential of application as lead compounds in the drug innovation and discovery.

Marine Fungal Indole Alkaloids

Prenylated Indole Alkaloids

Prenylated indole alkaloids are hybrid natural products with indole rings and isoprenoid fragments derived from tryptophan and prenyl diphosphates or their precursors, displaying a high structural diversity, especially the prenylated




Fig. 1 Representative-approved drugs containing an indole moiety for various diseases.




Fig. 2 Chemical structures of prenylated indole alkaloids 1–29 and 46–49.

tryptophan diketopiperazine skeleton (as shown in ►Figs. 2–10).¹⁷ These alkaloids are widely discovered from terrestrial and marine fungi, mainly focusing on the spectra of *Aspergillus* and *Penicillium*, with a wide spectrum of biological and pharmacological activities such as insecticidal, anti-parasitic, cytotoxic, and antimicrobial effects.^{17–19}

Notoamide/stephacidin-type alkaloids with a pyranindole ring are one of the most typical indole alkaloids. Notoamides A–D (1–4) are four new doubly prenylated indole alkaloids, first isolated from the marine fungus *Aspergillus* sp., which was derived from the common mussel *Mytilus edulis* (►Fig. 2). Notoamides A–C (1–3) with a dihydroxypyran-2-oxindole ring system exhibited moderate cytotoxic effect toward HeLa and L1210 cells.²⁰

Notoamide E (5) (►Fig. 2) was found in a marine-derived fungus *Aspergillus* sp., being considered as a key precursor of prenylated alkaloids in the biosynthesis process.^{21,22}

One new notoamide/stephacidin-type alkaloid, 21-hydroxystephacidin (6) (►Fig. 2), was harbored from the culture of a marine fungus *Aspergillus ostianus*.²³ Notoamides F–K (7–12) (►Fig. 2) were harbored from a marine-derived fungus strain *Aspergillus* sp. Notoamide I (10) exhibited weak cytotoxic effect toward HeLa cells (IC₅₀ = 21 μ g/mL).^{24,25}

Four notoamide/stephacidin-type analogues, named anti-podal (–)-versicolamide B (13) and notoamides L–N (14–16) (►Fig. 2), were produced by a marine-derived *Aspergillus* sp. Compound 14 is the first prenylated indole alkaloid presenting 25 carbons. Compound 15 is probably the precursor in the biosynthesis of the bicyclo[2.2.2]diazaoctane ring system.²⁶

Notoamides O–Q (17–19) (►Fig. 2) were isolated from a culture medium of marine-derived *Aspergillus* sp. Compound 17 contained a unique hemiacetal/hemiaminal ether moiety, which was unrepresented in these groups of prenylated indole alkaloids.^{27,28}

17-Epi-notoamides Q (20) and M (21) (►Fig. 2) were two new prenylated indole alkaloids, which were obtained from a marine-derived fungus *Aspergillus* sp.^{29,30}

Four new notoamide-type alkaloids, notoamides W–Z (22–25), as well as seven known analogues, notoamide F (7), notoamide G (8), 19-*epi*-notoamide R (26), notoamide I (10), stephacidin A (27), avrainvillamide (28), and a dimer of notoamide-type alkaloid stephacidin B (29), were discovered from a coral-associated fungus *Aspergillus ochraceus* LZDX-32–15 (►Fig. 2). Compounds 8, 25, and 26 exhibited potent inhibitory activity toward a panel of HCC (hepatocellular carcinoma) cell lines with IC₅₀ values in the range of 0.42 to




Fig. 3 Chemical structures of prenylated indole alkaloids 30–45 and 50.

3.39 $\mu\text{mol/L}$. Notoamide G (**8**) inhibited the viability of HepG2 and Huh-7 cells via apoptosis and autophagy way through a P38/JNK signaling pathway.³¹

Six new prenylated indole alkaloids with diketopiperazine ring, asperthrins A–F (**30–35**), were obtained from marine fungi *Aspergillus* sp. YJ191021 (**Fig. 3**). Asperthrin A (**30**) showed moderate inhibitory activities against three agricultural pathogenic microorganisms and significant antiinflammatory effect with IC_{50} value of $1.46 \pm 0.21 \mu\text{mol/L}$ in the model of human monocyte cell line (THP-1) induced by *Propionibacterium acnes*.³²

Paraherquamide J (**36**) (**Fig. 3**) is a new prenylated indole alkaloid, obtained from the fungus *Penicillium janthinellum* HK1-6, which was derived from mangrove rhizosphere soil. This alkaloid showed inactive effect in the assay of topoisomerase I (topo I) inhibitory, antibacterial, and lethality against brine shrimp *Artemia salina*.³³

Three new prenylated indole alkaloids waikikiamides A–C (**37–39**) (**Fig. 3**) with a complex diketopiperazine moiety were produced by the marine fungus *Aspergillus* sp. FM242. Compounds **37** and **38** contain an unprecedented indole alkaloid skeleton featuring with a hendecacyclic ring system. Compound **39** is the first unique heterodimer of two

notoamide derivatives joined by an N–O–C bridge. Compounds **37** and **39** showed significant antiproliferative activity against four cancer cell lines, HT1080, PC3, Jurkat, and A2780, with IC_{50} values in the range of 0.56 to 1.86 $\mu\text{mol/L}$.³⁴

Di-6-hydroxydeoxybrevianamide E (**40**) and dinotoamide J (**41**) (**Fig. 3**), two new homodimers, represent new examples of prenylated indole alkaloid, and were discovered from *Aspergillus austroafricanus* Y32-2. They exhibited proangiogenic effect in a zebrafish model of vascular injury induced by PTK787.³⁵

Three new alkaloids, spirotryprostatin G (**42**), and cyclotryprostatins F and G (**43** and **44**) (**Fig. 3**), were isolated from the marine-derived fungal strain *Penicillium brasilianum* HBU-136. Compound **42** exhibited potent cytotoxic effect against the HL-60 cell line ($IC_{50} = 6.0 \mu\text{mol/L}$). Compounds **43** and **44** showed cytotoxicity toward the MCF-7 cell line with IC_{50} values of 7.6 and 10.8 $\mu\text{mol/L}$, respectively.³⁶

17-Hydroxynotoamide D (**45**), 17-O-ethylnotoamide M (**46**), 10-O-acetylsclerotiamide (**47**), 10-O-ethylsclerotiamide (**48**), and 10-O-ethylnotoamide R (**49**) (**Figs. 1** and **3**) are five new prenylated indole alkaloids, being obtained from two marine-derived fungi *Aspergillus sulphureus* KMM 4640 and *Isaria felina* KMM 4639 by co-culture




Fig. 4 Chemical structures of prenylated indole alkaloids 51–70.

method. Compound **46** showed inhibitory effect against the human prostate cancer cells 22Rv1 at a concentration of 10 $\mu\text{mol/L}$.³⁷

Aspersiamides A–H (**50–57**) (**Fig. 3** and **4**), eight new linearly fused prenylated indole alkaloids with a rare pyrano[3,2-*f*]indole moiety, were found from the marine-derived fungus *Aspergillus versicolor*. Compound **56** showed potent inhibitory activity toward iNOS with an IC_{50} value of 5.39 $\mu\text{mol/L}$ in the antiinflammatory test.³⁸

Three new indole diketopiperazine alkaloids, asperochramides A–C (**58–60**) (**Fig. 4**), were obtained from a marine fungus *A. ochraceus*. Compound **58** exhibited significant antiinflammatory effects against the lipopolysaccharide (LPS)-stimulated RAW 264.7 cells.³⁹

Three new prenylated diketopiperazine indole alkaloids eurtiumins A–C (**61–63**) and one new prenylated indole alkaloid eurtiumin D (**64**) (**Fig. 4**) were produced by the marine-derived fungus *Eurotium* sp. SCSIO F452. Compounds **61** and **62** are a pair of diastereomers both with a hexahydropyrrolo[2,3-*b*]indole skeleton. Compound **63** exhibited significant radical scavenging effects toward DPPH with IC_{50} values of 13 $\mu\text{mol/L}$.⁴⁰

Taichunamide H (**65**) (**Fig. 4**), a new indole alkaloid with a fused-imine-containing pyrrole ring, was isolated from the fungus *A. versicolor*. The resonance at 190.4 ppm was assigned as the imine carbon in the molecule by using

X-ray diffraction, and the structure of taichunamide A was also revised. However, compound **65** showed no antifungal and cytotoxic activity.⁴¹

Mangrovamides D–G (**66–69**) (**Fig. 4**) are four new prenylated indole alkaloids from the mangrove sediment-derived fungus *Penicillium* sp. SCSIO041218 with no anti-allergic effect *in vitro* assay.⁴²

SF5280–415 (**70**) (**Fig. 4**), a new bispyrrolidinoindoline diketopiperazine alkaloid, and a known analogue SF5280–451 (**71**) (**Fig. 5**) were obtained from the marine-derived fungus *Aspergillus* sp. SF-5280. Compounds **70** and **71** displayed potent inhibitory effect toward PTP1B with IC_{50} values of 14.2 ± 0.7 and 12.9 ± 0.7 $\mu\text{mol/L}$, respectively.⁴³

Two new prenylated indole derivatives brevicompanine B (**72**) and verrucofortine (**73**) (**Fig. 5**) were found from a marine fungus *Penicillium* sp. NH-SL, and compound **73** showed potent cytotoxicity toward Hepa 1c1c7 cells.⁴⁴

Four new indole diketopiperazine alkaloids, *N*-(40-hydroxyprenyl)-cyclo(alanyltryptophyl) (**74**), isovariocolin I (**75**), 30-hydroxyechinulin (**76**), and 29-hydroxyechinulin (**77**) (**Fig. 5**), were obtained from the marine-derived fungus *Eurotium cristatum* EN-220. Compound **75** exhibited lethal effect against brine shrimp with the LD_{50} value of 19.4 $\mu\text{g/mL}$.⁴⁵

Two new prenylated indole derivatives, named penicimutamides D and E (**78** and **79**) (**Fig. 5**), were produced by the




Fig. 5 Chemical structures of prenylated indole alkaloids 71–89.

mutant fungus strain of *Penicillium purpurogenum* G59 derived from marine through the stimulation of diethyl sulfate. These two derivatives displayed weak suppressive effect toward cancer cell lines.⁴⁶

4,3-Hydroxysperadine A (**80**) (►Fig. 5), a new cyclopianzonic acid congener, was isolated from a marine sponge-associated fungus *Aspergillus oryzae* HMP-F28 by a bioassay-guided separation.⁴⁷

A known alkaloid, melegrin (**81**) (►Fig. 5), was rediscovered from a marine-derived fungus *Emericella dentata* Nq45. Its absolute structure was determined by the single-crystal X-ray diffraction method. This compound exhibited potent cytotoxic activity toward KB-3-1 cell line (the human cervix carcinoma) and KB-V1, a multidrug resistant subclone of KB-3-1, with the IC₅₀ values of 3.07 and 6.07 μmol/L, respectively. It also showed potent antibacterial effect against *Staphylococcus aureus* (minimum inhibitory concentration [MIC] = 0.25 mg/mL).⁴⁸

Misszrtine A (**82**) (►Fig. 5) is a novel prenylated indole alkaloid possessing a *N*-isopentenyl fragment, which was the first example of tryptophan methyl ester in this kind of alkaloids. This molecule, obtained from a marine sponge-derived fungus *Aspergillus* sp. SCSIO XWS03F03, displayed a

significant antagonistic effect against HL60 and LNCaP cell lines, with the IC₅₀ values of 3.1 and 4.9 μmol/L, respectively.⁴⁹

Cycloexpansamines A (**83**) and B (**84**) (►Fig. 5), two novel prenylated alkaloids with a spiroindolinone moiety, were produced by a marine fungus strain *Penicillium* sp. SF-5292.⁵⁰

Penioxamide A (**85**) (►Fig. 5), a new prenylated indole congener with a piperidine moiety and a unique antirelative configuration of the bicyclo[2.2.2]diazaoctane ring system, was afforded from the culture medium of fungus *Penicillium oxalicum* EN-201. Compound **85** exhibited pronounced lethality effect against brine shrimp (LD₅₀ = 5.6 μmol/L).⁵¹

Two diketopiperazine indole alkaloids, named fumitremorgin C (**86**) and 12,13-dihydroxy-fumitremorgin C (**87**) (►Fig. 5), were obtained from the culture of a fungus *Aspergillus* sp. BRF 030, displaying cytotoxic activity toward the HCT-116 cell line (IC₅₀ = 15.17 and 4.53 μmol/L, respectively).⁵²

A new diketopiperazine indole derivative, amaumine (**88**) (►Fig. 5), was isolated from the marine fungus *Auxarthron reticulatum* derived from marine sponge. This compound was considered to be a remarkable lead molecule for




Fig. 6 Chemical structures of prenylated indole alkaloids 90–105.

the research of selective antagonists of GPR18, for its potent suppressive effect toward GPR18 with the IC₅₀ value of 3.74 μmol/L.⁵³

Spirotryprostatin K (**89**) (→Fig. 5) is a new diketopiperazine alkaloid, obtained from an extract of the marine fungus *Aspergillus fumigatus*.⁵⁴

A new prenylated natural alkaloid takakiamide (**90**) (→Fig. 6) was obtained from the culture medium of the algicolous-derived fungus *Neosartorya takakii* KUFC 7898. However, this compound showed no antibacterial effect and no quorum sensing inhibitory activity.⁵⁵

Rubrumazines A–C (**91–93**) (→Fig. 6) are three new isochininulin-type indole diketopiperazine alkaloids bearing an oxygenated prenyl ether segment and produced by a mangrove-derived fungus *Eurotium rubrum* MA-150. Compounds **92** exhibited remarkable lethality toward brine shrimp with the LD₅₀ values of 2.43 μmol/L.⁵⁶

Rubrumlines A–O (**94–108**) as well as its known analogue neoechinulin B (**109**) are a series of indole diketopiperazine alkaloids obtained from the extract of the culture of the marine fungus *E. rubrum* (→Fig. 6). Neoechinulin B (**109**) showed significant inhibitory activity toward H1N1 virus in MDCK cells, and a class of influenza virus strains comprising

amantadine- and oseltamivir-resistant ones, which were isolated from clinical samples.⁵⁷

Dihydrocarneamide A (**110**) and iso-notoamide B (**111**) (→Fig. 7), two new prenylated indole analogues with the rare fused dimethyl-dihydropyran ring in the indole moiety by C-5 prenylation, are produced by the culture of the marine fungus *Paecilomyces variotii* EN-291. However, these two alkaloids represent weak cytotoxic effect toward the NCI-H460 cell line.⁵⁸

Cladosporin A (**112**) and cladosporin B (**113**) (→Fig. 7), two new sulfur-containing diketopiperazine indole derivatives, are harbored from a culture of fungus *Cladosporium* sp. derived from marine. These two compounds show moderate cytotoxicity against HepG2 cell line (IC₅₀ = 21 and 48 μg/mL).⁵⁹

Penipalines A and B (**114** and **115**), two new β-carbolines, and penipaline C (**116**), one new indole carbaldehyde congener, were afforded from the culture of the deep-sea-sediment fungus *Penicillium paneum* SD-44 (→Fig. 7). Compounds **115** and **116** exhibited significant cytotoxicity on A-549 and HCT-116 cell lines.⁶⁰

One new diketomorpholine derivative shornephine A (**117**) and a new prenylated indole 15b-β-methoxy-5-N-




Fig. 7 Chemical structures of prenylated indole alkaloids 110–135.

acetyladremin (**118**) were yielded from a marine sediment-derived *Aspergillus* sp. CMB-M081F (► **Fig. 7**). Compound **117** exhibited significant inhibitory activity against drug efflux mediated by P-glycoprotein in human multidrug-resistant colon cancer cells at the concentration of $20 \mu\text{mol/L}$.⁶¹

Versicamides A–H (**119–126**) (► **Fig. 7**), eight new prenylated alkaloids, were produced by the marine fungus *A. versicolor* HDN08–60. Compound **126** presented moderate suppressive effect on HL-60 cells ($IC_{50} = 8.7 \mu\text{mol/L}$), and it showed selective inhibitory function to PTK by further research of target screening.⁶²

Speradine F (**127**), a new alkaloid with a rare hexacyclic oxindole ring system, and two new alkaloids bearing a tetracyclic oxindole moiety, speradines G (**128**) and H (**129**), were obtained from a strain of the marine fungus *A. oryzae* (► **Fig. 7**). Unfortunately, these compounds displayed weak cytotoxic function toward the HeLa, HL-60, and K562 cell lines.⁶³

Three new indole diketopiperazine peroxides, 24-hydroxyverruculogen (**130**), 26-hydroxyverruculogen (**131**), and 13-O-prenyl-26-hydroxyverruculogen (**132**), were produced by the fungus *Penicillium brefeldianum* SD-273 isolated from marine sediment (► **Fig. 7**). Compound **132** exhibited remarkable brine shrimp lethal effect toward *A. salina* ($LD_{50} = 9.44 \mu\text{mol/L}$).⁶⁴

A prenylated indole, fumigaclavine C (**133**) (► **Fig. 7**), was obtained from a marine-derived fungus *A. fumigatus*. This compound induced apoptosis in MCF-7 cells via PI3/Akt and NF- κ B signaling, resulting in activating the mitochondrial cell death pathway.⁶⁵

Neoechinulins A (**134**) and B (**135**) (► **Fig. 7**) are two diketopiperazine indole alkaloids which were discovered and identified from the marine fungus *Eurotium* sp. SF-5989. Compounds **134** exhibited antiinflammatory activity by inhibiting the NF- κ B and p38 MAPK pathways in LPS-stimulated RAW264.7 macrophages.⁶⁶




Fig. 8 Chemical structures of prenylated indole alkaloids 136–161.

Brocaeloid C (**136**) (►Fig. 8), a new indole alkaloid possessing a C-2 reversed prenylated segment, was harbored from the cultures of *Penicillium brocae* MA-192, a marine fungus isolated from the fresh leaves of the mangrove *Avicennia marina*.⁶⁷

6-*Epi*-stephacidin A (**137**), *N*-hydroxy-6-*epi*-stephacidin A (**138**), and 6-*epi*-avrainvillamide (**139**) (►Fig. 8), prenylated alkaloids bearing a unique anti-bi cyclo-[2.2.2]diazaoctane core structure, were obtained from the cultures of marine fungus *Aspergillus taichungensis*. (+)-Versicolamides B and C (**140** and **141**) with a spiro-center, as well as six derivatives (**142**–**147**), were harbored as conversion products of compound **138** through a photo-induced reaction (►Fig. 8). Compounds **138** and **139** displayed remarkable cytotoxic effect toward two cell lines, with IC_{50} values of 3.02 and 1.92 $\mu\text{mol/L}$ against A549 cells, 4.45 and 1.88 $\mu\text{mol/L}$ against HL-60 cells, respectively.⁶⁸

An indole alkaloid, named neoechinulin A (**148**) (►Fig. 8), which was produced by a marine fungus *Microsporium* sp., exhibited inhibitory effect toward the microglia activation induced by amyloid- β oligomer, and the protection of

inflammation-mediated toxicity in PC-12 cells, suggesting its potential to be developed as a protective agent for neuro-inflammation associated with Alzheimer's disease. Compound **148** also showed an effect of inducing apoptosis in HeLa cells in another biotest.^{69,70}

Carneamides A–C (**149**–**151**) (►Fig. 8) are three prenylated indole alkaloids that are afforded from the fungus *Aspergillus carneus* KMM 4638 derived from marine environment. However, these compounds showed no *in vitro* antimicrobial and cytotoxic effects.⁷¹

Cristatumins A–D (**152**–**155**) (►Fig. 8), four new indole analogues, were obtained and identified from the culture of the marine alga endophytic fungus *E. cristatum* EN-220. Compound **152** produced antibacterial effect both toward *Escherichia coli* and *S. aureus*.⁷²

Waikialoid A (**156**) (►Fig. 8), a new dimer of prenylated indole derivative, is produced by a strain of marine fungus *Aspergillus* sp. This molecule exhibits potent inhibitory activity against the biofilm formation of *Candida albicans* (IC_{50} = 1.4 $\mu\text{mol/L}$). And waikialoid A (**156**) displayed the potential to be developed as a promising lead for the biofilm




Fig. 9 Chemical structures of prenylated indole alkaloids 162–183.




Fig. 10 Chemical structures of prenylated indole alkaloids 184–195.

inhibitors in combination with antibiotics for this metabolite, and showed no cytotoxicity toward fungi or human cells (200 $\mu\text{mol/L}$).⁷³

Cyclotryprostatin E (**157**) (\blacktriangleright Fig. 8), a new indole diketopiperazine alkaloid, was discovered from the strain of marine fungus *Aspergillus sydowii* SCSIO 00305, which was separated from a healthy tissue of gorgonian coral *Verrucella umbraculum*.⁷⁴

Tryptoquivalines P and Q (**158** and **159**) (\blacktriangleright Fig. 8) are two new indole alkaloids which were produced by a marine-derived fungal strain *Neosartorya* sp. HN-M-3.⁷⁵

Two new prenylated indole alkaloids, identified as 2-(3,3-dimethylprop-1-ene)-costaclavine (**160**) and 2-(3,3-dimethylprop-1-ene)-epicostaclavine (**161**) (\blacktriangleright Fig. 8), had been obtained from the culture of marine fungal strain *A. fumigatus*. Compounds **160** and **161** exhibited weak cytotoxicity against P388 cell lines.⁷⁶

Two novel chlorinated prenylated indole alkaloids, (–)-spiromalbramide (**162**) and (+)-isomalbrancheamide B (**163**), and two new brominated derivatives, (+)-malbrancheamide C (**164**) and (+)-isomalbrancheamide C (**165**) (\blacktriangleright Fig. 9), were discovered from an invertebrate-derived fungal strain *Malbranchea graminicola* assisted by a direct analysis in real time mass spectrometry technique.⁷⁷

A prenylated indole alkaloid, neoechinulin A (**166**) (\blacktriangleright Fig. 9), was produced by a marine fungus, and exhibited a protected effect toward PC12 cells against the cytotoxicity of 1-methyl-4-phenylpyridinium (MPP⁺) and rotenone, two neurotoxins inducing Parkinsonian.⁷⁸

7-O-Methylvariecolortide A (**167**) (\blacktriangleright Fig. 9), a new alkaloid with spirocyclic diketopiperazine ring, was produced by the cultures of fungal strain *E. rubrum* derived from the stems of mangrove *Hibiscus tiliaceus*.⁷⁹

Compound (**168**), spirotryprostatins C–E (**169–171**), fumitremorgin B derivatives **172** and **173**, and 13-oxoverruculogen (**174**) (\blacktriangleright Fig. 9) were seven new prenylated indole diketopiperazine alkaloids, which were obtained and identified from the marine fungus *A. fumigatus* isolated from holothurian. Compounds **171–173** exhibited better selectivity toward MOLT-4, HL-60, and A549 than toward other compounds.⁸⁰

Three new prenylated indole alkaloids with an oxaspiro [4.4]lactam core ring system, 6-methoxyspirotryprostatin B (**175**), 18-oxotryprostatin A (**176**), and 14-hydroxyterezine D (**177**), were discovered from the cultures of a marine-derived fungus *A. sydowii* PFW1-13 (\blacktriangleright Fig. 9). Compounds **175–177** showed significant cytotoxic effect toward the A549 cell line (IC₅₀ = 8.29, 1.28, and 7.31 $\mu\text{mol/L}$, respectively). Compound **175** also exhibited weak inhibitory function to HL-60 cells (IC₅₀ = 9.71 $\mu\text{mol/L}$). Compounds **176** and **177** produced potent antimicrobial effect toward *Bacillus subtilis*, *E. coli*, and *Micrococcus lysodeikticus* (MICs = 14.97, 3.74, and 7.49 $\mu\text{mol/L}$ for **176**; MICs = 5.33, 10.65, and 10.65 $\mu\text{mol/L}$ for **177**).⁸¹

Shearinines D – F (**178–180**) (\blacktriangleright Fig. 9), three new prenylated indole alkaloids, were yielded from the culture of marine fungal strain *P. janthinellum* Biourge. Compounds **178** and **179** could induce apoptosis in HL-60 cells, and **179**

also exhibited inhibitory activity against EGF-induced malignant transformation in JB6 P+ Cl 41 cells.⁸²

Two new pentacyclic indolinone alkaloids, named citrinadins A and B (**181** and **182**), as well as **181**'s known derivative compound **183** (\blacktriangleright Fig. 9), were obtained from the marine red alga-derived fungal strain of *Penicillium citrinum*. Compound **181** exhibited growth-inhibitory effect against L1210 and KB cells (IC₅₀ = 6.2 and 10 $\mu\text{g/mL}$); compound **182** displayed modest cytotoxic activity toward L1210 cells (IC₅₀ = 10 $\mu\text{g/mL}$).^{83,84} Lep F (**184**) and Lep C (**185**) (\blacktriangleright Fig. 10), two prenylated bisindole alkaloids produced by marine fungal strain of *Leptoshaeria* species, exhibited potent growth suppressive effect toward RPMI8402 and 293 tumor cell lines. And these two compounds also induced apoptosis through suppressing the survival pathway by inactivation of Akt/protein kinase B. What's more, they were remarkable topoisomerase catalytic inhibitors. Compound **185** targets topo I both *in vitro* and *in vivo* and **184** targets both topo I and II *in vitro*.^{85,86}

A class of isoechinulin-type indole alkaloids, dihydroxyisoechinulin A (**186**), golmaenone (**187**), neoechinulin A (**188**), *L*-alanyl-*L*-tryptophan anhydride (**189**), and echinulin (**190**) (\blacktriangleright Fig. 10), bearing isoprenic chains in the indole ring, were obtained from the cultures of marine fungal strain *Aspergillus* sp. Compounds **186** and **188** displayed potent radical scavenging effect toward 1,1-diphenyl-2-picrylhydrazyl (DPPH) (IC₅₀ = 20, 20, and 24 $\mu\text{mol/L}$, respectively), similar to ascorbic acid (positive control, IC₅₀ = 20 $\mu\text{mol/L}$). They also displayed significant ultraviolet (UA)-A protective activity (ED₅₀ = 130, 90, and 170 $\mu\text{mol/L}$, respectively), more potent than oxybenzone (a currently used sunscreen agent, ED₅₀ = 350 $\mu\text{mol/L}$).^{87,88}

A new chiral dipyrrolbenzoquinone alkaloid, terreusinone (**191**), was found from the marine algicolous fungus *Aspergillus terreus*. Another new analogue terreusinol (**192**) was produced by biotransformation of terreusinone (**191**) in the co-culture of terreusinone and *Streptomyces* sp. (\blacktriangleright Fig. 10). These two compounds showed potent UV-A protecting activity (ED₅₀ = 70 $\mu\text{g/mL}$ for **191**, and 150 $\mu\text{mol/L}$ for **192**), which were more active than positive control oxybenzone (ED₅₀ = 350 $\mu\text{mol/L}$).^{89,90}

Indolyl alkaloids with a prenylated chain, oxaline (**193**) (\blacktriangleright Fig. 10), was isolated from the extract of the culture of an unidentified fungal strain derived from the marine red alga *Gracilaria verrucosa*.⁹¹

Tryprostatins A and B (**194** and **195**) (\blacktriangleright Fig. 10), two new prenylated indole alkaloids, were isolated from a marine fungal strain *A. fumigatus* BM939, which was collected from a sea sediment sample. These two metabolites showed mammalian cell-cycle inhibitory activity.⁹²

Diketopiperazine Indole Alkaloids

The diketopiperazine indole alkaloids without a prenylated fragment were included in this section (\blacktriangleright Fig. 11). The 2,5-diketopiperazine ring is usually a cyclodipeptide that is condensed by two amino acids. In the diketopiperazine indoles, the condensed six-membered ring is formed by tryptophan and another amino acid.¹⁸




Fig. 11 Chemical structures of diketopiperazine indole alkaloids 196–210.

Haenamindole (**196**) (**►Fig. 11**) is a diketopiperazine alkaloid with benzyl-hydroxypiperazindione and phenylpyrimidoindole fragments, discovered from the marine-derived fungus *Penicillium* sp. KCB12F005. However, this compound displayed no significant cytotoxic and antimicrobial activity.⁹³

Three pairs of enantiomers (\pm)-acrozines A–C ((\pm)-**197** to (\pm)-**199**) (**►Fig. 11**) with a novel *N*-methoxy diketopiperazine ring system, were affogred from the marine green alga-derived fungus *Acrostalagmus luteoalbus* TK-43. (+)-Acrozines A ((+)-**197**) displayed better inhibitory activity toward acetylcholinesterase ($IC_{50} = 2.3 \mu\text{mol/L}$) than that of (–)-acrozines A ((–)-**197**) and (\pm)-**197**.⁹⁴

A new indole diketopiperazine alkaloid, raistrickindole A (**200**) (**►Fig. 11**), bearing a unique pyrazino[1',2':2,3]-[1,2]oxazino[6,5-b]indole tetraheterocyclic core ring, was produced by the marine fungal strain of *Penicillium raistrickii* IMB17-034. Compound **200** exhibited inhibitory effects toward the hepatitis C virus.⁹⁵

Dichotomocej D (**201**), a new aliphatic amide, and dichocerazines A and B (**202** and **203**) (**►Fig. 11**), two diketopiperazines, were isolated from the culture of a marine fungal strain of *Dichotomomyces cejpui* F31-1.⁹⁶

A new indole diketopiperazine alkaloid, asperochramide D (**204**) (**►Fig. 11**), was affogred from the culture extract of marine-derived fungus *A. ochraceus*. Compound **204** represents a rare example of indole diketopiperazines possessing a 3-hydroxyl-2-indolone ring system.³⁹

A pair of bridged irregularly epimonothiodiketopiperazine diastereomers, pseudellones A and B (**205** and **206**) (**►Fig. 11**), containing a unique 3-indolylglycine and alanine segments, and a new alkaloid pseudellone C (**207**), bearing an

unusual nucleus, were obtained from the culture medium of marine fungus *Pseudallescheria ellipsoidea* F42-3.⁹⁷

An indole diketopiperazine alkaloid, gliotoxin (**208**) (**►Fig. 11**), discovered from the marine-derived fungus *Aspergillus* sp., produced apoptosis via the mitochondrial pathway in HeLa and SW1353 cells, resulting in an apoptotic type of cell death.⁹⁸

Luteoalbusins A and B (**209** and **210**) (**►Fig. 11**), two new indole diketopiperazine alkaloids, were yielded from the deep-sea sediment-derived fungus *A. luteoalbus* SCSIO F457. Compounds **209** and **210** exhibited potent cytotoxic activity toward SF-268, MCF-7, NCI-H460, and HepG-2 cell lines.⁹⁹

Quinazoline-Containing Indole Alkaloids

Aspertoryadins A–G (**211–217**) are seven new quinazoline-containing indole alkaloids (**►Fig. 12**), produced by the mollusk-derived marine fungus *Aspergillus* sp. HNMF114. Aspertoryadin A (**211**) bears a unique aminosulfonyl group in the molecule, an exceedingly rare moiety in nature. Aspertoryadins F and G (**216** and **217**) were found to have quorum-sensing inhibitory effect to *Chromobacterium violaceum* CV026, both with MIC values of 32 $\mu\text{g/well}$.¹⁰⁰

Chaetominine (CHA) (**218**) (**►Fig. 12**), a quinazolinone alkaloid produced by marine crab-derived fungus *A. fumigatus* CY018, showed potent growth-inhibitory activity toward K562 and SW1116 cell lines.¹⁰¹

Neofiscalin A (**219**) and fiscalin C (**220**) (**►Fig. 12**) were two quinazolinone alkaloids obtained from *Neosartorya siamensis* KUFA 0017, a marine sponge-associated fungus. They exhibited potential for the development as new leads of anti-Gram-positive bacterial infectious agents especially in multidrug-resistant strains.¹⁰²




Fig. 12 Chemical structures of quinazoline-containing indole alkaloids 211–222.

Tryptoquivalines *R* and *S* (**221** and **222**) (\blacktriangleright Fig. 12) are two new indole quinazolinone alkaloids harbored from the organic extract of a marine fungal strain *Neosartorya* sp. HN-M-3.¹⁰³

Bisindoles

Two new bisindole alkaloids, fusariumindoles A and B (**223** and **224**) (\blacktriangleright Fig. 13), were yielded from the marine fungus *Fusarium* sp. L1 stimulated by *L*-tryptophan supplementation.¹⁰⁴

Asterriquinone F (**225**) (\blacktriangleright Fig. 13), a new bisindole quinone alkaloid, was obtained from the culture of *A. terreus* LM.1.5.¹⁰⁵

(\pm)-Fusaspoid A (**226a/226b**) (\blacktriangleright Fig. 13), obtained as a pair of new bisindole alkaloid enantiomers, were produced by the marine fungal strain of *Fusarium* sp. XBB-9. Compounds **226a/226b** were inactive in the cytotoxic assay in HCT-15 and RKO cell lines.¹⁰⁶

Chaetoindolone A (**227**) and chaetoindolone C (**228**) (\blacktriangleright Fig. 13), two new indole alkaloids, were produced by




Fig. 13 Chemical structures of bisindole alkaloids 223–234.

the strain of *Chaetomium globosum* 1C51 through biotransformation, a fungus collected from a marine fish sample. Compound **227** was confirmed to suppress the growth of the rice-pathogenic bacteria *Xanthomonas oryzae* pv. *oryzae* (*xoo*).¹⁰⁷

2,2-Bis(6-bromo-3-indolyl)ethylamine (**229**) (►Fig. 13), a bisindole alkaloid derived from a strain of marine fungus, was confirmed to have the potential to be developed as an antibiofilm lead compound for its greatest antimicrobial and biofilm formation inhibitory activity.¹⁰⁸

A new bisindole alkaloid indolepyrazine A (**230**) (►Fig. 13) containing an unrepresented indole-pyrazine-oxindole skeleton was obtained from the marine fungal strain of *Acinetobacter* sp. ZZ1275. Compound **230** exhibited significant antimicrobial effects toward methicillin-resistant strains *E. coli*, *S. aureus*, and *C. albicans* (MIC values, 10, 12, and 12 µg/mL, respectively).¹⁰⁹

Pseudobindoles A and B (**231** and **232**) (►Fig. 13), two new bisindole alkaloids, were yielded from the cultures of marine fungus *Pseudallescheria boydii* F44-1 through adding amino acids to the culture medium.¹¹⁰

Varioloids C and D (**233** and **234**) (►Fig. 13), two indolyl-6,10*b*-dihydro-5*a*H-[1]benzofuro[2,3-*b*]indole alkaloids, were isolated from the strain of *P. variotii* EN-291, a marine alga-derived fungus. Both compounds **233** and **234** displayed cytotoxic activity toward HepG2, HCT116, and A549 cell lines (IC₅₀ = 2.6–8.2 µg/mL).^{111,112}

Indoloditerpenes

Indoloditerpenes are a series of structurally diverse meroterpenoids featuring with an indole ring connected with a cyclic diterpene backbone, distributed widely both in terrestrial and marine fungi, exhibiting great potential of drug research as lead compounds for their potent insecticidal, antiviral, cytotoxic, and antimicrobial effects.^{17,104,113}

Fusaindoterpenes A and B (**235** and **236**) (►Fig. 14), two new indoloditerpenes or their derivatives, were afforded from a marine-derived fungal strain of *Fusarium* sp. L1 by adding *L*-tryptophan in culture supplementation. Compound **235** possesses a unique 6/9/6/6/5 heterocyclic ring system. And compound **236** exhibited significant effect toward Zika virus (EC₅₀ = 7.5 µmol/L).¹⁰⁴

Compound **237** (►Fig. 14), a new indoloditerpene, was obtained from a culture of marine fungal strain of *A. versicolor* ZZ761. This compound displayed antimicrobial effects against *E. coli* and *C. albicans* (MIC = 20.6 and 22.8 µmol/L, respectively).¹¹³

Anthcolorin G and H (**238** and **239**) (►Fig. 14), two new oxindoloditerpene epimers, were yielded from the cultures of *A. versicolor*, a mangrove endophytic fungus. Compound **239** produced weak cytotoxicity toward HeLa cells.¹¹⁴

Penicindopene A (**240**) (►Fig. 14), a new indoloditerpene possessing a rare 3-hydroxyl-2-indolone fragment, was obtained from the strain of *Penicillium* sp. YPCMAC1, a deep-sea-derived fungus. Compound **240** showed moderate




Fig. 14 Chemical structures of indoloditerpenes 235–249.

cytotoxic activity against A549 and HeLa cell lines ($IC_{50} = 15.2$ and $20.5 \mu\text{mol/L}$).¹¹⁵

Emindole SB β -mannoside (**241**) and 27-*O*-methylasporzyn C (**242**) (\blacktriangleright Fig. 14), two new indoloditerpenes, were produced by a marine-derived strain of *D. cejpii*. Compound **241** was confirmed to be a CB2 antagonist, and compound **242** was found to be the first indole derivative possessing selective GPR18 inhibitory activity. These two indole derivatives may be investigated as lead molecules for the research of GPR18- and CB receptor-blocking drugs.¹¹⁶

Asporzyns A–C (**243–245**) (\blacktriangleright Fig. 14), three new indoloditerpene derivatives, were found from the cultures of *A. oryzae*, an endophytic fungus derived from the marine red alga *Heterosiphonia japonica*. Compound **245** showed potent inhibitory effect toward *E. coli*.¹¹⁷

Epipaxilline (**246**) and penerpene J (**247**) (\blacktriangleright Fig. 14), two new indoloditerpenes, were obtained from the organic extract of the cultures of marine fungus *Penicillium* sp. KFD28. They displayed inhibitory effects toward PTP1B ($IC_{50} = 31.5$ and $9.5 \mu\text{mol/L}$) and compound **247** also presented suppressive effects against TCPTP ($IC_{50} = 14.7 \mu\text{mol/L}$).¹¹⁸

Penerpenes A–D (**248–251**) (\blacktriangleright Figs. 14 and 15), four unique indoleterpenoids, were obtained and identified from the marine fungal strain of *Penicillium* sp. KFD28. Compounds **248** and **249** exhibited significant inhibitory effect against protein tyrosine phosphatases (PTP1B and TCPTP).¹¹⁹

Penerpenes E–I (**252–256**) (\blacktriangleright Fig. 15), five new indoleterpenoids, were obtained from *Penicillium* sp. KFD28, a marine fungus isolated from a bivalve mollusk *Meretrix lusoria*. Compound **252** possesses a rare 6/5/5/6/6/5/5 heptacyclic moiety. Compound **253** is a new carbon skeleton of indole-diterpenoid derived from paxilline. Compound **254** contains a unique 6/5/5/6/6/7 hexacyclic skeleton bearing a 1,3-dioxepane ring. Compounds **252**, **253**, and **256** displayed inhibitory effects toward protein tyrosine phosphatase 1B (PTP1B) ($IC_{50} = 14$, 27 , and $23 \mu\text{mol/L}$, respectively).¹²⁰

Penijanthenes C and D (**257** and **258**) (\blacktriangleright Fig. 15), two new indole-diterpenoid derivatives, were obtained from the cultures of a marine-derived fungal strain *P. janthinellum*. These two compounds exhibited potent antivibrio effect toward three pathogenic *Vibrio* spp. ($MIC = 3.1$ – $50.0 \mu\text{mol/L}$).¹²¹




Fig. 15 Chemical structures of indoloditerpenes 250–272.




Fig. 16 Chemical structures of cytochalasans 273–287.

Asperindoles A–D (**259–262**) (**Fig. 15**), four new indole-diterpene alkaloids, were produced by the cultures of a marine ascidian-derived fungal strain *Aspergillus* sp. Asperindoles C and D (**261** and **262**) bear an unusual 2-hydroxyisobutyric acid (2-HIBA) moiety, which is very rare in natural products. Compound **259** showed cytotoxic effect toward PC-3 and 22Rv1 cells that are resistant to hormone therapy, as well as human prostate cancer cells that are sensitive to hormone therapy, and apoptosis-induced activity in the above-mentioned cells.¹²²

Rhizovarin A–F (**263–268**) (**Fig. 15**), six new indole-diterpenes, were discovered by a genome-mining method from the strain of *Mucor irregularis* QEN-189, a marine fungus derived from the mangrove plant *Rhizophora stylosa*. Among these molecules, compounds **263–265** contain the most complex and unique structure of the indole-diterpenes, with a rare acetal linked to a hemiketal or a ketal forming a novel 4,6,6,8,5,6,6,6-fused indole-diterpene skeleton. Compounds **263** and **264** displayed inhibitory effect toward A-549 and HL-60 cancer cell lines.¹²³

19-Hydroxyphenitrem A (**269**) and 19-hydroxyphenitrem E (**270**) (**Fig. 15**), two new chlorinated indole-diterpenoids, were yielded from the marine strain of *Aspergillus nidulans* EN-330, which was isolated from red alga *Polysiphonia scopulorum* var. *villum*. Compounds **269** and **270** showed cytotoxicity toward brine shrimp ($LD_{50} = 3.2$ and $4.6 \mu\text{mol/L}$). In addition, compound **269** displayed moderate antimicrobial function against four pathogens *Edwardsiella tarda*, *Vibrio anguillarum*, *E. coli*, and *S. aureus*.¹²⁴

Compounds **271** and **272** (**Fig. 15**), two new indole-diterpenoids, were harbored from the cultures of the marine-derived fungal strain of *Aspergillus flavus* OUCMDZ-

2205. Compound **271** presented antibacterial effect toward *S. aureus* ($MIC = 20.5 \mu\text{mol/L}$). And both compounds could arrest the cell cycle in the S phase at $10 \mu\text{mol/L}$ in A549 cell lines. Compound **271** exhibited PKC- β inhibitory activity ($IC_{50} = 15.6 \mu\text{mol/L}$).¹²⁵

Cytochalasans

Chaetoglobosin-510 (**273**), -540 (**274**), and -542 (**275**), three cytochalasin-type alkaloids (**Fig. 16**), were obtained from the cultures of the marine fungal strain of *Phomopsis asparagi*. Chaetoglobosin-542 (**275**) produced antimicrofilament effect and cytotoxic activity against leukemia cancer and murine colon cell lines.¹²⁶

Cytoglobosins A–G (**276–282**) (**Fig. 16**), seven new cytochalasin-type alkaloids, were obtained from the cultures of marine green alga-derived fungus *C. globosum* QEN-14. Compounds **278** and **279** exhibited cytotoxicity toward the A549 tumor cell line.¹²⁷

A new cytochalasans derivative, 6-O-methyl-chaetoglobosin Q (**283**), along with several known analogues, was produced by a coral-associated fungus *C. globosum* C2F17. The known congeners chaetoglobosin E (**284**) and Fex (**285**) (**Fig. 16**), exhibited significant antiproliferative activity toward a series of cancer cell lines ($IC_{50} = 1.4 - 9.2 \mu\text{mol/L}$).¹²⁸

Cytoglobosins H and I (**286** and **287**) (**Fig. 16**), two new cytochalasans derivatives, as well as several known congeners, were obtained from a deep-sea sediment-derived marine fungal strain of *C. globosum*. The known compound chaetoglobosin E (**284**) displayed potent growth-inhibitory effect against LNCaP and B16F10 cell lines ($IC_{50} = 0.62$ and $2.78 \mu\text{mol/L}$, respectively). Besides, compound **284** suppressed the growth of LNCaP cells via inducing apoptosis.¹²⁹




Fig. 17 Chemical structures of other indole alkaloids 288–306.

Other Indole Alkaloids

Plectosphaeric acids A–C (288–290) (►Fig. 17), three new plectosphaeric acid derivatives, were obtained from the laboratory cultures of the fungus *Plectosphaerella cucumerina* collected from marine sediments. They were evaluated and presented significant inhibitory activity against indoleamine-2,3-dioxygenase (IDO), with IC₅₀ values of 2 μmol/L.¹³⁰

Chaetindolone B (291), chaetindolone D (292), 19-*O*-demethylchaetogline A (293), 20-*O*-demethylchaetogline F (294), and chaetogline A (295) (►Fig. 17) are five new indole alkaloids obtained from the marine fungus *C. globosum* 1C51, through biotransformation induced by 1-methyl-*L*-tryptophan (1-MT) supplemented in the culture medium. Chaetogline A (295) showed fungicidal effect toward a pathogenic fungus *Sclerotinia sclerotiorum*, the cause of rape sclerotinia rot, indicating the potential agrochemical significance of indole alkaloids.¹⁰⁷

Two new indole alkaloids, Fusariumindoles C (296) and (±)-isoalternatine A (297) (►Fig. 17), were produced by the marine fungus *Fusarium* sp. L1, induced by the *L*-tryptophan added in the culture condition. However, they exhibited no significant effect against Zika virus.¹⁰⁴

A known marine indole alkaloid derivative isolated from a fungal strain of *Neosartorya pseudofischeri*, isochaetominine C (298) (►Fig. 17), displayed potent cytotoxic activity toward the Sf9 cells.⁵²

5,6-Dihydroxyindole-2-carboxylic acid (DHICA) (299) was yielded from the cultures of marine fungus *A. nidulans* (►Fig. 17). The simple indole alkaloid 299 showed remarkable UV-B protecting effect both *in vivo* and *in vitro* assays, presenting the potential as a sun-protective agent added to sunscreen cream.¹³¹

Two simple indole alkaloids, tryptamine (300) and indole-3-carbaldehyde (301) (►Fig. 17), were expressed by the marine-derived fungus *Penicillium* species. Compound 301 showed modest antimicrobial activity.¹³²

1-(4-Hydroxybenzoyl)indole-3-carbaldehyde (302) (►Fig. 17), a new indole alkaloid with an aldehyde group, was isolated from a strain of marine fungus *Engyodontium album* IVB1b. It was inactive both in cytotoxic and antimicrobial tests.¹³³

Indolepyrazine B (303) (►Fig. 17), a new indole alkaloid, was harbored from the culture medium of marine-derived fungus *Acinetobacter* sp. ZZ1275. It displayed antimicrobial effect against *E. coli*, *S. aureus*, and *C. albicans*, three methicillin-resistant pathogenic strains, with MIC values of 8, 12, and 14 μg/mL, respectively.¹⁰⁹

A new indole carboxylic acid, nigrospin A (304) (►Fig. 17), was obtained from the cultures of the marine fungal strain *Nigrospora oryzae* SCSGAF 0111.¹³⁴

Compound 305 (►Fig. 17) was a new tryptoquivaline derivative isolated from the marine alga-derived fungus *N.*

takakii KUFC 7898. In the antimicrobial biotests, it exhibited no significant activity.⁵⁵

Fumiquinazoline K (**306**) (► Fig. 17), a new indole alkaloid, had been obtained from a strain of marine fungus *A. fumigatus* KMM 4631, which was derived from the soft coral *Simularia* sp.¹³⁵

Conclusion

In this review, we have investigated and summarized comprehensively the chemical diversity and biological activity of marine fungal indole alkaloids from 1995 to early 2021, covering a total of 306 indole derivatives. The chemical types of these marine fungal indole alkaloids can be mainly classified to prenylated indoles, diketopiperazine indoles, bisindoles, quinazoline-containing indoles, indole-diterpenoids, and others. As shown in ► Fig. 18, the prenylated indoles represent the predominant marine fungal alkaloids (63.5%) exhibiting high structural diversity, especially the prenylated tryptophan diketopiperazine skeleton. As for the sources, the species of *Aspergillus* (41.2%) and *Penicillium* (19.2%) are the two main producing strains of marine fungal indoles. As shown in ► Table 1, the natural indole metabolites from marine fungi displayed excellent cytotoxic, antimicrobial, sun-protective, antiinflammatory, antiviral, neuropro-




Fig. 18 Percentage of structural types of marine fungal indole alkaloids.

TECTIVE, kinase inhibitory, crop-protective, and brine shrimp lethal activities. They presented great potential of research as new lead structures for the development of new drugs, especially GPR18-selective antagonist **88**, biofilm inhibitor **156**, anti-multidrug-resistant bacterial molecules **219** and **220**, and GPR18- and CB antagonists **241** and **242**.

Table 1 The indole alkaloids from marine fungus covering from 1995 to the early 2021

Compounds	Sources	Bioactivities	Ref.
Notoamides A–D (1–4)	<i>Aspergillus</i> sp.	Cytotoxicity (1–3)	20
Notoamide E (5)	<i>Aspergillus</i> sp.		21,22
21-Hydroxystephacidin (6)	<i>Aspergillus ostianus</i>		23
Notoamides F–K (7–12)	<i>Aspergillus</i> sp.	Cytotoxicity (10)	24,25
(–)-Versicolamide B (13) and notoamides L–N (14–16)	<i>Aspergillus</i> sp.		26
Notoamides O–Q (17–19)	<i>Aspergillus</i> sp.		27,28
17-Epi-notoamides Q (20) and M (21)	<i>Aspergillus</i> sp.		29,30
Notoamides W–Z (22–25), 19-epi-notoamide R (26), stephacidin A (27), avrainvillamide (28), stephacidin B (29)	<i>Aspergillus ochraceus</i>	Cytotoxicity (8, 25, 26)	31
Asperthrins A–F (30–35)	<i>Aspergillus</i> sp. YJ191021	Cytotoxicity (30)	32
Paraherquamide J (36)	<i>Penicillium janthinellum</i> HK1–6	Inactive	33
Waikikiamides A–C (37–39)	<i>Aspergillus</i> sp. FM242	Cytotoxicity (37, 39)	34
Di-6-hydroxydeoxybrevianamide E (40) and dinotoamide J (41)	<i>Aspergillus austroafricanus</i> Y32–2	Proangiogenic effect (40 and 41)	35
Spirotryprostatin G (42), cyclotryprostatins F and G (43 and 44)	<i>Penicillium brasilianum</i> HBU-136	Cytotoxicity (42–44)	36
17-Hydroxynotoamide D (45), 17-O-ethylnotoamide M (46), 10-O-acetylsclerotiamide (47), 10-O-ethylsclerotiamide (48), 10-O-ethylnotoamide R (49)	<i>Aspergillus sulphureus</i> KMM 4640 and <i>Isaria felina</i> KMM 4639	Cytotoxicity (46)	37
Asperversiamides A–H (50–57)	<i>Aspergillus versicolor</i>	Antiinflammatory (56)	38

Table 1 (Continued)

Compounds	Sources	Bioactivities	Ref.
Asperochramides A–C (58–60)	<i>Aspergillus ochraceus</i>	Antiinflammatory (58)	39
Eurotiumins A–D (61–64)	<i>Eurotium</i> sp. SCSIO F452	Radical scavenging (63)	40
Taichunamide H (65)	<i>Aspergillus versicolor</i>	No activity	41
Mangrovamides D–G (66–69)	<i>Penicillium</i> sp. SCSIO041218	No antiallergic effect	42
SF5280–415 (70), SF5280–451 (71)	<i>Aspergillus</i> sp. SF-5280	PTP1B inhibition	43
Brevicompanine B (72) and verrucofortine (73)	<i>Penicillium</i> sp. NH-SL	Cytotoxicity (73)	44
<i>N</i> -(40-hydroxyprenyl)-cyclo(alanyltryptophyl) (74), isovaricolorin I (75), 30-hydroxyechinulin (76), 29-hydroxyechinulin (77)	<i>Eurotium cristatum</i> EN-220	Brine shrimp lethality (75)	45
Penicimutamides D and E (78 and 79)	<i>Penicillium purpurogenum</i> G59	Cytotoxicity	46
4,3-Hydroxysperadine A (80)	<i>Aspergillus oryzae</i> HMP-F28		47
Meleagrins (81)	<i>Emericella dentata</i> Nq45	Cytotoxicity, antibacterial	48
Misszrtine A (82)	<i>Aspergillus</i> sp. SCSIO XWS03F03	Cytotoxicity	49
Cycloexpansamines A (83) and B (84)	<i>Penicillium</i> sp. SF-5292		50
Penioxamide A (85)	<i>Penicillium oxalicum</i> EN-201	Brine shrimp lethality	51
Fumitremorgin C (86), 12,13-dihydroxy-fumitremorgin C (87)	<i>Aspergillus</i> sp. BRF 030	Cytotoxicity	52
Amauromine (88)	<i>Auxarthron reticulatum</i>	Antagonists of GPR18	53
Spirotryprostatin K (89)	<i>Aspergillus fumigatus</i>		54
Takakiamide (90)	<i>Neosartorya takakii</i> KUFC 7898	No antibacterial effect	55
Rubrumazines A–C (91–93)	<i>Eurotium rubrum</i> MA-150	Brine shrimp lethality (92)	56
Rubrumlines A–O (94–108), neoechinulin B (109)	<i>Eurotium rubrum</i>	Anti-influenza virus (109)	57
Dihydrocarneamide A (110) and isonotoamide B (111)	<i>Paecilomyces variotii</i> EN-291	Cytotoxicity	58
Cladosporin A (112), cladosporin B (113)	<i>Cladosporium</i> sp.	Cytotoxicity	59
Penipalines A–C (114–116)	<i>Penicillium paneum</i> SD-44	Cytotoxicity	60
Shornephine A (117) and 15b- β -methoxy-5- <i>N</i> -acetyladreemin (118)	<i>Aspergillus</i> sp. CMB-M081F	Inhibitory against drug efflux	61
Versicamides A–H (119 – 126)	<i>Aspergillus versicolor</i> HDN08–60	Cytotoxicity	62
Speradines F–H (127 – 129)	<i>Aspergillus oryzae</i>	Cytotoxicity	63
24-Hydroxyverruculogen (130), 26-hydroxyverruculogen (131), and 13- <i>O</i> -prenyl-26-hydroxyverruculogen (132)	<i>Penicillium brefeldianum</i> SD-273	Brine shrimp lethality (132)	64
Fumigaclavine C (133)	<i>Aspergillus fumigatus</i>	Apoptosis	65
Neoechinulins A (134) and B (135)	<i>Eurotium</i> sp. SF-5989	Antiinflammatory (134)	66
Brocaeloid C (136)	<i>Penicillium brocae</i> MA-192		67
6- <i>Epi</i> -stephacidin A (137), <i>N</i> -hydroxy-6- <i>epi</i> -stephacidin A (138), 6- <i>epi</i> -avrainvillamide (139), (+)-versicolamides B and C (140 and 141), compounds 142–147	<i>Aspergillus taichungensis</i>	Cytotoxicity (138 and 139)	68
Neoechinulin A (148)	<i>Microsporium</i> sp.	Apoptosis, neuroinflammatory modulation	69,70

(Continued)

Table 1 (Continued)

Compounds	Sources	Bioactivities	Ref.
Carneamides A–C (149–151)	<i>Aspergillus carneus</i> KMM 4638	No effects	71
Cristatumins A–D (152–155)	<i>Eurotium cristatum</i> EN-220	Antibacterial effect (152)	72
Waikialoid A (156)	<i>Aspergillus</i> sp.	Biofilm inhibitors	73
Cyclotryprostatin E (157)	<i>Aspergillus sydowii</i> SCSIO 00305		74
Tryptoquivalines P and Q (158 and 159)	<i>Neosartorya</i> sp. HN-M-3		75
2-(3,3-Dimethylprop-1-ene)-costaclavine (160) and 2-(3,3-dimethylprop-1-ene)-epicostaclavine (161)	<i>Aspergillus fumigatus</i>	Cytotoxicity	76
(–)-Spiromalbramide (162), (+)-isomalbrancheamide B (163), (+)-malbrancheamide C (164), (+)-isomalbrancheamide C (165)	<i>Malbranchea graminicola</i>		77
Neoechinulin A (166)		Neuroprotection	78
7-O-Methylvariecolortide A (167)	<i>Eurotium rubrum</i>		79
Compound (168), spirotryprostatins C–E (169–171), fumitremorgin B derivatives 172 and 173, and 13-oxoverruculogen (174)	<i>Aspergillus fumigatus</i>	Cytotoxicity (171–173)	80
6-Methoxyspirotryprostatin B (175), 18-oxotryprostatin A (176), and 14-hydroxyterezine D (177)	<i>Aspergillus sydowii</i> PFW1–13	Cytotoxicity (175–177) Antimicrobial effect (176 and 177)	81
Shearinines D–F (178–180)	<i>Penicillium janthinellum</i> Biourge	Antimalignant transformation	82
Citrinadins A, B (181 and 182) and derivative 183	<i>Penicillium citrinum</i>	Cytotoxicity	83,84
Lep F (184) and Lep C (185)	<i>Leptoshaeria</i> sp.	Topo inhibition cytotoxicity	85,86
Dihydroxyisoechinulin A (186), golmaenone (187), neoechinulin A (188), L-alanyl-L-tryptophan anhydride (189), and echinulin (190)	<i>Aspergillus</i> sp.	Ultraviolet-A protective activity radical scavenging effect	87,88
Terreusinone (191), terreusinol (192)	<i>Aspergillus terreus</i> (191) <i>Streptomyces</i> sp. (192)	UV-A protective activity	89,90
Oxaline (193)	Unidentified fungal strain		91
Tryprostatins A and B (194 and 195)	<i>Aspergillus fumigatus</i> BM939	Cell cycle inhibitory activity	92
Haenamindole (196)	<i>Penicillium</i> sp. KCB12F005	No cytotoxic and antimicrobial activity	93
(±)-Acrozines A–C ((±)-197–(±)-199)	<i>Acrostalagmus luteoalbus</i> TK-43	Acetylcholinesterase inhibition ((+)-197)	94
Raistrickindole A (200)	<i>Penicillium raistrickii</i> IMB17–034	Anti-hepatitis C virus	95
Dichotomocej D (201), dichocerazines A and B (202 and 203)	<i>Dichotomomyces cejpji</i> F31–1		96
Asperochramides D (204)	<i>Aspergillus ochraceus</i>		39
Pseudellones A–C (205–207)	<i>Pseudallescheria ellipsoidea</i>		97
Gliotoxin (208)	<i>Aspergillus</i> sp.	Apoptosis	96
Luteoalbusins A and B (209 and 210)	<i>Acrostalagmus luteoalbus</i> SCSIO F457	Cytotoxicity	99
Aspertoryadins A–G (211–217)	<i>Aspergillus</i> sp. HNMF114	Antibacterial activity (216 and 217)	100

Table 1 (Continued)

Compounds	Sources	Bioactivities	Ref.
Chaetominine (CHA) (218)	<i>Aspergillus fumigatus</i> CY018	Cytotoxicity	101
Neofiscalin A (219) and fiscalin C (220)	<i>Neosartorya siamensis</i> KUFA 0017	Antibacterial activity	102
Tryptoquivalines R and S (221 and 222)	<i>Neosartorya</i> sp. HN-M-3		103
Fusariumindoles A and B (223 and 224)	<i>Fusarium</i> sp. L1		104
Asterriquinone F (225)	<i>Aspergillus terreus</i> LM.1.5		105
(±)-Fusaspoid A (226a/226b)	<i>Fusarium</i> sp. XBB-9	Inactive	106
Chaetoinolone A (227) and chaetoinolone C (228)	<i>Chaetomium globosum</i> 1C51	Antibacterial activity	107
2,2-Bis(6-bromo-3-indolyl) ethylamine (229)		Antibiofilm formation	108
Indolepyrazines A (230)	<i>Acinetobacter</i> sp. ZZ1275	Antimicrobial effect	109
Pseudoindoles A and B (231 and 232)	<i>Pseudallescheria boydii</i> F44-1		110
Varioloids C and D (233 and 234)	<i>Paecilomyces variotii</i> EN-291	Cytotoxicity	111,112
Fusaindoterpenes A and B (235 and 236)	<i>Fusarium</i> sp. L1	Anti-Zika virus	104
Compound 237	<i>Aspergillus versicolor</i> ZZ761	Antimicrobial effect	113
Anthcolorin G and H (238 and 239)	<i>Aspergillus versicolor</i>	Cytotoxicity	114
Penicindopene A (240)	<i>Penicillium</i> sp. YPCMAC1	Cytotoxicity	115
Emindole SB β -mannoside (241) and 27-O-methylasporozin C (242)	<i>Dichotomomyces cejpaii</i>	CB2 antagonist (241) GPR18 antagonist (242)	116
Asporozins A–C (243–245)	<i>Aspergillus oryzae</i>	Antimicrobial effect (245)	117
Epipaxilline (246) and penerpene J (247)	<i>Penicillium</i> sp. KFD28	PTP1B inhibition (246 and 247) TCPTP inhibition (247)	118
Penerpenes A–D (248–251)	<i>Penicillium</i> sp. KFD28	PTP1B and TCPTP inhibition (248 and 249)	119
Penerpenes E–I (252–256)	<i>Penicillium</i> sp. KFD28	PTP1B inhibition (252, 253, and 256)	120
Penijanthines C and D (257 and 258)	<i>Penicillium janthinellum</i>	Antivibrio effect	121
Asperindoles A–D (259–262)	<i>Aspergillus</i> sp.	Cytotoxicity and apoptosis (259)	122
Rhizovarins A–F (263–268)	<i>Mucor irregularis</i> QEN-189	Cytotoxicity (263 and 264)	123
19-Hydroxypenitrem A (269) and 19-hydroxypenitrem E (270)	<i>Aspergillus nidulans</i> EN-330	Brine shrimp lethality (269 and 270) Antimicrobial activity (269)	124
Compounds 271 and 272	<i>Aspergillus flavus</i> OUCMDZ-2205	Antibacterial effect (271) PKC β inhibitory activity (271)	125
Chaetoglobosin-510 (273), -540 (274), and -542 (275)	<i>Phomopsis asparagi</i>	Antimicrofilament effect (275) Cytotoxicity (275)	126
Cytoglobosins A–G (276–282)	<i>Chaetomium globosum</i> QEN-14	Cytotoxicity (278 and 279)	127
6-O-Methyl-chaetoglobosin Q (283), chaetoglobosins E (284) and Fex (285)	<i>Chaetomium globosum</i> C2F17	Cytotoxicity (284 and 285)	128
Cytoglobosins H and I (286 and 287), chaetoglobosin E (284)	<i>Chaetomium globosum</i>	Cytotoxicity and apoptosis (284)	129
Plectosphaeroic acids A–C (288–290)	<i>Plectosphaerella cucumerina</i>	IDO inhibition	130
	<i>Chaetomium globosum</i> 1C51	Fungicidal effect (295)	107

(Continued)

Table 1 (Continued)

Compounds	Sources	Bioactivities	Ref.
Chaetoinolone B (291), chaetoinolone D (292), 19-O-demethylchaetogline A (293), 20-O-demethylchaetogline F (294), and chaetogline A (295)			
Fusariumindoles C (296) and (\pm)-isoalternatine A (297)	<i>Fusarium</i> sp. L1	Inactive against Zika virus	104
Isochaetominine C (298)	<i>Neosartorya pseudofischeri</i>	Cytotoxicity	52
5,6-Dihydroxyindole-2-carboxylic acid (DHICA) (299)	<i>Aspergillus nidulans</i>	UVB protecting effect	131
Tryptamine (230) and indole-3-carbaldehyde (231)	<i>Penicillium</i> sp.	Antimicrobial activity (231)	132
1-(4-Hydroxybenzoyl)indole-3-carbaldehyde (302)	<i>Engyodontium album</i> IVB1b		133
Indolepyrazines B (303)	<i>Acinetobacter</i> sp. ZZ1275	Antimicrobial effect	109
Nigrospin A (304)	<i>Nigrospora oryzae</i> SCSGAF 0111		134
Compound 305	<i>Neosartorya takakii</i> KUFC 7898		55
Fumiquinazoline K (306)	<i>Aspergillus fumigatus</i> KMM 4631		135

Funding/Acknowledgments

This work was funded by grants from the National Natural Science Foundation of China (Grant No. 82022065, 81872791, 82073696, and U20A20136), the Key Project of Science and Technology of Shanghai (Grant No. 21S11900800), the Sanhang Program of Second Military Medical University, and the Key Research and Development Program of Ningxia (Grant No. 2019BFG02017).

Conflict of Interest

The authors declare no conflict of interest.

References

- Roth GJ, Binder R, Colbatzky F, et al. Nintedanib: from discovery to the clinic. *J Med Chem* 2015;58(03):1053–1063
- Zhou S, Huang G. The synthesis and biological activity of marine alkaloid derivatives and analogues. *Rsc Adv* 2020;10(53):31909–31935
- Singh TP, Singh OM. Recent progress in biological activities of indole and indole alkaloids. *Mini Rev Med Chem* 2018;18(01):9–25
- Cong H, Zhao X, Castle BT, et al. An indole-chalcone inhibits multidrug-resistant cancer cell growth by targeting microtubules. *Mol Pharm* 2018;15(09):3892–3900
- Zhuang C, Zhang W, Sheng C, Zhang W, Xing C, Miao Z. Chalcone: a privileged structure in medicinal chemistry. *Chem Rev* 2017;117(12):7762–7810
- Ma H, Wu Y, Zhang W, Zhang H, Miao Z, Zhuang C. Radio-sensitization of human pancreatic cancer by piperlongumine analogues. *Chin Chem Lett* 2021;32(03):1197–1201
- Kumar D, Sharma S, Kalra S, Singh G, Monga V, Kumar B. Medicinal perspective of indole derivatives: recent developments and structure-activity relationship studies. *Curr Drug Targets* 2020;21(09):864–891
- Abraham W-R. Fumitremorgins and relatives—from tremorgenic compounds to valuable anti-cancer drugs. *Curr Med Chem* 2018;25(02):123–140
- Ishikura M, Yamada K, Abe T. Simple indole alkaloids and those with a nonrearranged monoterpene unit. *Nat Prod Rep* 2010;27(11):1630–1680
- Rodrigues T, Reker D, Schneider P, Schneider G. Counting on natural products for drug design. *Nat Chem* 2016;8(06):531–541
- Carroll AR, Copp BR, Davis RA, Keyzers RA, Prinsep MR. Marine natural products. *Nat Prod Rep* 2021;38(02):362–413
- Stien D. Marine microbial diversity as a source of bioactive natural products. *Mar Drugs* 2020;18(04):215
- Zhang P, Wei Q, Yuan X, Xu K. Newly reported alkaloids produced by marine-derived *Penicillium* species (covering 2014–2018). *Bioorg Chem* 2020;99:103840
- Wang C, Tang S, Cao S. Antimicrobial compounds from marine fungi. *Phytochem Rev* 2021;20(01):85–117
- Xu K, Yuan XL, Li C, Li AX. Recent discovery of heterocyclic alkaloids from marine-derived *Aspergillus* species. *Mar Drugs* 2020;18(01):54
- Willems T, De Mol ML, De Bruycker A, De Maeseneire SL, Soetaert WK. Alkaloids from marine fungi: promising antimicrobials. *Antibiotics (Basel)* 2020;9(06):340
- Li SM. Prenylated indole derivatives from fungi: structure diversity, biological activities, biosynthesis and chemoenzymatic synthesis. *Nat Prod Rep* 2010;27(01):57–78
- Almeida MC, Resende DISP, da Costa PM, Pinto MMM, Sousa E. Tryptophan derived natural marine alkaloids and synthetic derivatives as promising antimicrobial agents. *Eur J Med Chem* 2021;209:112945
- Klas KR, Kato H, Frisvad JC, et al. Structural and stereochemical diversity in prenylated indole alkaloids containing the bicyclo [2.2.2]diazaoctane ring system from marine and terrestrial fungi. *Nat Prod Rep* 2018;35(06):532–558
- Kato H, Yoshida T, Tokue T, et al. Notoamides A–D: prenylated indole alkaloids isolated from a marine-derived fungus, *Aspergillus* sp. *Angew Chem Int Ed Engl* 2007;46(13):2254–2256
- Tsukamoto S, Kato H, Greshock TJ, Hirota H, Ohta T, Williams RM. Isolation of notoamide E, a key precursor in the biosynthesis of prenylated indole alkaloids in a marine-derived fungus, *Aspergillus* sp. *J Am Chem Soc* 2009;131(11):3834–3835

- 22 Tsukamoto S, Kato H, Greshock TJ, Hirota H, Ohta T, Williams RM. Isolation of notoamide E, a key precursor in the biosynthesis of prenylated indole alkaloids in a marine-derived fungus, *Aspergillus* sp. (vol 131, pg 3834, 2009). *J Am Chem Soc* 2013;135(29):10878
- 23 Kito K, Ookura R, Kusumi T, Namikoshi M, Ooi T. X-Ray structures of two stephacidins, heptacyclic alkaloids from the marine-derived fungus *Aspergillus Ostianus*. *Heterocycles* 2009;78(08):2101–2106
- 24 Tsukamoto S, Kato H, Samizo M, et al. Notoamides F-K, prenylated indole alkaloids isolated from a marine-derived *Aspergillus* sp. *J Nat Prod* 2008;71(12):2064–2067
- 25 Tsukamoto S, Kato H, Samizo M, et al. Correction to Notoamides F-K, prenylated indole alkaloids isolated from a marine-derived *Aspergillus* sp. *J Nat Prod* 2013;76(06):1233
- 26 Tsukamoto S, Kawabata T, Kato H, et al. Isolation of antipodal (-)-versicolamide B and notoamides L-N from a marine-derived *Aspergillus* sp. *Org Lett* 2009;11(06):1297–1300
- 27 Tsukamoto S, Umaoka H, Yoshikawa K, Ikeda T, Hirota H. Notoamide O, a structurally unprecedented prenylated indole alkaloid, and notoamides P-R from a marine-derived fungus, *Aspergillus* sp. *J Nat Prod* 2010;73(08):1438–1440
- 28 Tsukamoto S, Umaoka H, Yoshikawa K, Ikeda T, Hirota H. Notoamide O, a structurally unprecedented prenylated indole alkaloid, and notoamides p-r from a marine-derived fungus, *Aspergillus* sp. (vol 73, pg 1438, 2010). *J Nat Prod* 2013;76(06):1232
- 29 Chen M, Shao CL, Fu XM, et al. Bioactive indole alkaloids and phenyl ether derivatives from a marine-derived *Aspergillus* sp. *Fungus*. *J Nat Prod* 2013;76(04):547–553
- 30 Chen M, Shao CL, Fu XM, et al. Correction to bioactive indole alkaloids and phenyl ether derivatives from a marine-derived *Aspergillus* sp. *Fungus*. *J Nat Prod* 2013;76(06):1229
- 31 Hu L, Zhang T, Liu D, et al. Notoamide-type alkaloid induced apoptosis and autophagy via a P38/JNK signaling pathway in hepatocellular carcinoma cells. *Rsc Adv* 2019;9(34):19855–19868
- 32 Yang J, Gong L, Guo M, et al. Bioactive indole diketopiperazine alkaloids from the marine endophytic fungus *Aspergillus* sp. YJ191021. *Mar Drugs* 2021;19(03):157
- 33 Zheng YY, Shen NX, Liang ZY, Shen L, Chen M, Wang C-Y. Paraherquamide J, a new prenylated indole alkaloid from the marine-derived fungus *Penicillium janthinellum* HK1-6. *Nat Prod Res* 2020;34(03):378–384
- 34 Wang F, Sarotti AM, Jiang G, et al. Waikikiamides A-C: complex diketopiperazine dimer and diketopiperazine-polyketide hybrids from a Hawaiian marine fungal strain *Aspergillus* sp. FM242. *Org Lett* 2020;22(11):4408–4412
- 35 Li P, Zhang M, Li H, et al. New prenylated indole homodimeric and pteridine alkaloids from the marine-derived fungus *Aspergillus austroafricanus* Y32-2. *Mar Drugs* 2021;19(02):98
- 36 Zhang YH, Geng C, Zhang XW, et al. Discovery of bioactive indole-diketopiperazines from the marine-derived fungus *Penicillium brasilianum* aided by genomic information. *Mar Drugs* 2019;17(09):514
- 37 Afiyatullof SS, Zhuravleva OI, Antonov AS, et al. Prenylated indole alkaloids from co-culture of marine-derived fungi *Aspergillus sulphureus* and *Isaria felina*. *J Antibiot (Tokyo)* 2018;71(10):846–853
- 38 Li H, Sun W, Deng M, et al. Asperversiamides, linearly fused prenylated indole alkaloids from the marine-derived fungus *Aspergillus versicolor*. *J Org Chem* 2018;83(15):8483–8492
- 39 Wen H, Liu X, Zhang Q, et al. Three new indole diketopiperazine alkaloids from *Aspergillus ochraceus*. *Chem Biodivers* 2018;15(04):e1700550
- 40 Zhong WM, Wang JF, Shi XF, et al. Eurotiumins A-E, five new alkaloids from the marine-derived fungus *Eurotium* sp. SCSIO F452. *Mar Drugs* 2018;16(04):136
- 41 Li F, Zhang Z, Zhang G, et al. Determination of taichunamide H and structural revision of taichunamide A. *Org Lett* 2018;20(04):1138–1141
- 42 Yang B, Tao H, Lin X, et al. Prenylated indole alkaloids and chromone derivatives from the fungus *Penicillium* sp. SCSIO041218. *Tetrahedron* 2018;74(01):77–82
- 43 Cho KH, Sohn JH, Oh H. Isolation and structure determination of a new diketopiperazine dimer from marine-derived fungus *Aspergillus* sp. SF-5280. *Nat Prod Res* 2018;32(02):214–221
- 44 Ding H, Ding W, Ma Z. Mass spectrometric characteristics of prenylated indole derivatives from marine-derived *Penicillium* sp. NH-SL. *Mar Drugs* 2017;15(03):86
- 45 Du FY, Li X, Li XM, Zhu LW, Wang BG. Indolediketopiperazine alkaloids from *Eurotium cristatum* EN-220, an endophytic fungus isolated from the marine alga *Sargassum thunbergii*. *Mar Drugs* 2017;15(02):24
- 46 Wu CJ, Li CW, Gao H, Huang XJ, Cui CB. Penicimutamides D-E: two new prenylated indole alkaloids from a mutant of the marine-derived *Penicillium purpurogenum* G59. *RSC Advances* 2017;7(40):24718–24722
- 47 Cao T, Ling J, Liu Y, et al. Characterization and abolishment of the cyclopiazonic acids produced by *Aspergillus oryzae* HMP-F28. *Biosci Biotechnol Biochem* 2018;82(10):1832–1839
- 48 Hamed A, Abdel-Razek AS, Araby M, et al. Meleagrins from marine fungus *Emericella dentata* Nq45: crystal structure and diverse biological activity studies. *Nat Prod Res* 2021;35(21):3830–3838
- 49 Zhou R, Liao X, Li H, et al. Isolation and synthesis of misszrtine A: a novel indole alkaloid from marine sponge-associated *Aspergillus* sp. SCSIO XWS03F03. *Front Chem* 2018;6:212
- 50 Lee C, Sohn JH, Jang JH, et al. Cycloexpansamines A and B: spiroindolinone alkaloids from a marine isolate of *Penicillium* sp. (SF-5292). *J Antibiot (Tokyo)* 2015;68(11):715–718
- 51 Zhang P, Li XM, Liu H, Li X, Wang BG. Two new alkaloids from *Penicillium oxalicum* EN-201, an endophytic fungus derived from the marine mangrove plant *Rhizophora stylosa*. *Phytochem Lett* 2015;13:160–164
- 52 Lan WJ, Fu SJ, Xu MY, et al. Five new cytotoxic metabolites from the marine fungus *Neosartorya pseudofischeri*. *Mar Drugs* 2016;14(01):18
- 53 Nazir M, Harms H, Loefl, et al. GPR18 inhibiting amauromine and the novel triterpene glycoside auxarthronoside from the sponge-derived fungus *Auxarthron reticulatum*. *Planta Med* 2015;81(12–13):1141–1145
- 54 Shi YS, Zhang Y, Chen XZ, Zhang N, Liu YB. Metabolites produced by the endophytic fungus *Aspergillus fumigatus* from the stem of *Erythrophloeum fordii* Oliv. *Molecules* 2015;20(06):10793–10799
- 55 Zin WW, Buttachon S, Buaruang J, et al. A new meroditerpene and a new tryptoquivaline analog from the algicolous fungus *Neosartorya takakii* KUFC 7898. *Mar Drugs* 2015;13(06):3776–3790
- 56 Meng LH, Du FY, Li XM, Pedpradab P, Xu GM, Wang BG. Rubrumazines A–C, indolediketopiperazines of the isoehinulin class from *Eurotium rubrum* MA-150, a fungus obtained from marine mangrove-derived rhizospheric soil. *J Nat Prod* 2015;78(04):909–913
- 57 Chen X, Si L, Liu D, et al. Neoechinulin B and its analogues as potential entry inhibitors of influenza viruses, targeting viral hemagglutinin. *Eur J Med Chem* 2015;93:182–195
- 58 Zhang P, Li XM, Wang JN, Li X, Wang BG. Prenylated indole alkaloids from the marine-derived fungus *Paecilomyces variotii*. *Chin Chem Lett* 2015;26(03):313–316
- 59 Gu B, Zhang Y, Ding L, et al. Preparative separation of sulfur-containing diketopiperazines from marine fungus *Cladosporium* sp. using high-speed counter-current chromatography in step-wise elution mode. *Mar Drugs* 2015;13(01):354–365

- 60 Li CS, Li XM, An CY, Wang BG. Prenylated indole alkaloid derivatives from marine sediment-derived fungus *Penicillium paneum* SD-44. *Helv Chim Acta* 2014;97(10):1440–1444
- 61 Khalil ZG, Huang XC, Raju R, Piggott AM, Capon RJ. Shornephine A: structure, chemical stability, and P-glycoprotein inhibitory properties of a rare diketomorpholine from an Australian marine-derived *Aspergillus* sp. *J Org Chem* 2014;79(18):8700–8705
- 62 Peng J, Gao H, Li J, et al. Prenylated indole diketopiperazines from the marine-derived fungus *Aspergillus versicolor*. *J Org Chem* 2014;79(17):7895–7904
- 63 Hu X, Xia QW, Zhao YY, et al. Speradines F-H, three new oxindole alkaloids from the marine-derived fungus *Aspergillus oryzae*. *Chem Pharm Bull (Tokyo)* 2014;62(09):942–946
- 64 An CY, Li XM, Li CS, Xu GM, Wang BG. Prenylated indole-diketopiperazine peroxides and related homologues from the marine sediment-derived fungus *Penicillium brefeldianum* SD-273. *Mar Drugs* 2014;12(02):746–756
- 65 Li YX, Himaya SWA, Dewapriya P, Zhang C, Kim SK. Fumigaclavine C from a marine-derived fungus *Aspergillus fumigatus* induces apoptosis in MCF-7 breast cancer cells. *Mar Drugs* 2013;11(12):5063–5086
- 66 Kim KS, Cui X, Lee DS, et al. Anti-inflammatory effect of neoechinulin A from the marine fungus *Eurotium* sp. SF-5989 through the suppression of NF- κ B and p38 MAPK pathways in lipopolysaccharide-stimulated RAW264.7 macrophages. *Molecules* 2013;18(11):13245–13259
- 67 Zhang P, Meng LH, Mandi A, et al. Brocaeloids A-C, 4-oxoquinoline and indole alkaloids with C-2 reversed prenylation from the mangrove-derived endophytic fungus *Penicillium brocae*. *Eur J Org Chem* 2014;2014(19):4029–4036
- 68 Cai S, Luan Y, Kong X, Zhu T, Gu Q, Li D. Isolation and photoinduced conversion of 6-epi-stephacidins from *Aspergillus tai-chungensis*. *Org Lett* 2013;15(09):2168–2171
- 69 Dewapriya P, Li YX, Himaya SWA, Pangestuti R, Kim SK. Neoechinulin A suppresses amyloid- β oligomer-induced microglia activation and thereby protects PC-12 cells from inflammation-mediated toxicity. *Neurotoxicology* 2013;35:30–40
- 70 Wijesekera I, Li YX, Vo TS, Ta QV, Ngo DH, Kim SK. Induction of apoptosis in human cervical carcinoma HeLa cells by neoechinulin A from marine-derived fungus *Microsporium* sp. *Process Biochem* 2013;48(01):68–72
- 71 Zhuravleva OI, Afiyatullov SSH, Denisenko VA, et al. Secondary metabolites from a marine-derived fungus *Aspergillus carneus* Blochwitz. *Phytochemistry* 2012;80:123–131
- 72 Du FY, Li XM, Li CS, Shang Z, Wang BG. Cristatamins A-D, new indole alkaloids from the marine-derived endophytic fungus *Eurotium cristatum* EN-220. *Bioorg Med Chem Lett* 2012;22(14):4650–4653
- 73 Wang X, You J, King JB, Powell DR, Cichewicz RH. Waikialoid A suppresses hyphal morphogenesis and inhibits biofilm development in pathogenic *Candida albicans*. *J Nat Prod* 2012;75(04):707–715
- 74 He F, Sun YL, Liu KS, et al. Indole alkaloids from marine-derived fungus *Aspergillus sydowii* SCSIO 00305. *J Antibiot (Tokyo)* 2012;65(02):109–111
- 75 Sun FY, Chen G, Bai J, Li W, Pei YH. Two new alkaloids from a marine-derived fungus *Neosartorya* sp. HN-M-3. *J Asian Nat Prod Res* 2012;14(12):1109–1115
- 76 Zhang D, Satake M, Fukuzawa S, et al. Two new indole alkaloids, 2-(3,3-dimethylprop-1-ene)-costaclavine and 2-(3,3-dimethylprop-1-ene)-epicostaclavine, from the marine-derived fungus *Aspergillus fumigatus*. *J Nat Med* 2012;66(01):222–226
- 77 Watts KR, Loveridge ST, Tenney K, Media J, Valeriote FA, Crews P. Utilizing DART mass spectrometry to pinpoint halogenated metabolites from a marine invertebrate-derived fungus. *J Org Chem* 2011;76(15):6201–6208
- 78 Akashi S, Kimura T, Takeuchi T, et al. Neoechinulin A impedes the progression of rotenone-induced cytotoxicity in PC12 cells. *Biol Pharm Bull* 2011;34(02):243–248
- 79 Li DL, Li XM, Proksch P, Wang BG. 7-O-Methylvariecolortide A, a new spirocyclic diketopiperazine alkaloid from a marine mangrove derived endophytic fungus, *Eurotium rubrum*. *Nat Prod Commun* 2010;5(10):1583–1586
- 80 Wang F, Fang Y, Zhu T, et al. Seven new prenylated indole diketopiperazine alkaloids from holothurian-derived fungus *Aspergillus fumigatus*. *Tetrahedron* 2008;64(34):7986–7991
- 81 Zhang M, Wang WL, Fang YC, Zhu TJ, Gu QQ, Zhu WM. Cytotoxic alkaloids and antibiotic nordammarane triterpenoids from the marine-derived fungus *Aspergillus sydowii*. *J Nat Prod* 2008;71(06):985–989
- 82 Smetanina OF, Kalinovsky AI, Khudyakova YV, et al. Indole alkaloids produced by a marine fungus isolate of *Penicillium janthinellum* Biourge. *J Nat Prod* 2007;70(06):906–909
- 83 Mugishima T, Tsuda M, Kasai Y, et al. Absolute stereochemistry of citrinadins A and B from marine-derived fungus. *J Org Chem* 2005;70(23):9430–9435
- 84 Tsuda M, Kasai Y, Komatsu K, et al. Citrinadin A, a novel pentacyclic alkaloid from marine-derived fungus *Penicillium citrinum*. *Org Lett* 2004;6(18):3087–3089
- 85 Yanagihara M, Sasaki-Takahashi N, Sugahara T, et al. Leptosins isolated from marine fungus *Leptoshaeria* species inhibit DNA topoisomerases I and/or II and induce apoptosis by inactivation of Akt/protein kinase B. *Cancer Sci* 2005;96(11):816–824
- 86 Utsugi T, Aoyagi K, Asao T, et al. Antitumor activity of a novel quinoline derivative, TAS-103, with inhibitory effects on topoisomerases I and II. *Jpn J Cancer Res* 1997;88(10):992–1002
- 87 Li Y, Li X, Kang JS, Choi HD, Son BW. New radical scavenging and ultraviolet-A protecting prenylated dioxopiperazine alkaloid related to isoechinulin A from a marine isolate of the fungus *Aspergillus*. *J Antibiot (Tokyo)* 2004;57(05):337–340
- 88 Li Y, Li X, Kim SK, et al. Golmaenone, a new diketopiperazine alkaloid from the marine-derived fungus *Aspergillus* sp. *Chem Pharm Bull (Tokyo)* 2004;52(03):375–376
- 89 Lee SM, Li XF, Jiang H, et al. Terreusinone, a novel UV-A protecting dipyrroloquinone from the marine algicolous fungus *Aspergillus terreus*. *Tetrahedron Lett* 2003;44(42):7707–7710
- 90 Li X, Lee SM, Choi HD, Kang JS, Son BW. Microbial transformation of terreusinone, an ultraviolet-A (UV-A) protecting dipyrroloquinone, by *Streptomyces* sp. *Chem Pharm Bull (Tokyo)* 2003;51(12):1458–1459
- 91 Li Y, Li XF, Kim DS, Choi HD, Son BW. Indolyl alkaloid derivatives, Nb-acetyltryptamine and oxaline from a marine-derived fungus. *Arch Pharm Res* 2003;26(01):21–23
- 92 Cui CB, Makeya H, Okada G, et al. Tryprostatins A and B, novel mammalian cell cycle inhibitors produced by *Aspergillus fumigatus*. *J Antibiot (Tokyo)* 1995;48(11):1382–1384
- 93 Kim JW, Ko SK, Son S, et al. Haenamindole, an unusual diketopiperazine derivative from a marine-derived *Penicillium* sp. KCB12F005. *Bioorg Med Chem Lett* 2015;25(22):5398–5401
- 94 Cao J, Li XM, Meng LH, et al. Isolation and characterization of three pairs of indole-diketopiperazine enantiomers containing infrequent N-methoxy substitution from the marine algal-derived endophytic fungus *Acrostalagmus luteoalbus* TK-43. *Bioorg Chem* 2019;90:103030
- 95 Li J, Hu Y, Hao X, et al. Raistrickindole A, an anti-HCV oxazinoindole alkaloid from *Penicillium raistrickii* IMB17-034. *J Nat Prod* 2019;82(05):1391–1395
- 96 Chen YX, Xu MY, Li HJ, et al. Diverse secondary metabolites from the marine-derived fungus *Dichotomomyces ceipii* F31-1. *Mar Drugs* 2017;15(11):339
- 97 Liu W, Li HJ, Xu MY, et al. Pseudellones A–C, three alkaloids from the marine-derived fungus *Pseudallescheria ellipsoidea* F42-3. *Org Lett* 2015;17(21):5156–5159

- 98 Nguyen VT, Lee JS, Qian ZJ, et al. Gliotoxin isolated from marine fungus *Aspergillus* sp. induces apoptosis of human cervical cancer and chondrosarcoma cells. *Mar Drugs* 2013;12(01):69–87
- 99 Wang FZ, Huang Z, Shi XF, et al. Cytotoxic indole diketopiperazines from the deep sea-derived fungus *Acrostalagmus luteoalbus* SCSIO F457. *Bioorg Med Chem Lett* 2012;22(23):7265–7267
- 100 Kong FD, Zhang SL, Zhou SQ, et al. Quinazoline-containing indole alkaloids from the marine-derived fungus *Aspergillus* sp. HNMF114. *J Nat Prod* 2019;82(12):3456–3463
- 101 Liu C, Wei XC, An FL, Lu YH. Ammonium acetate supplement strategy for enhancement of chaetominine production in liquid culture of marine-derived *Aspergillus fumigatus* CY018. *J Microbiol Biotechnol* 2019;29(04):587–595
- 102 Bessa LJ, Buttachon S, Dethoup T, et al. Neofiscalin A and fiscalin C are potential novel indole alkaloid alternatives for the treatment of multidrug-resistant Gram-positive bacterial infections. *FEMS Microbiol Lett* 2016;363(15):fnw150
- 103 Xu N, Cao Y, Wang L, Chen G, Pei YH. New alkaloids from a marine-derived fungus *Neosartorya* sp. HN-M-3. *J Asian Nat Prod Res* 2013;15(07):731–736
- 104 Guo YW, Liu XJ, Yuan J, et al. L-Tryptophan induces a marine-derived *Fusarium* sp. to produce indole alkaloids with activity against the Zika virus. *J Nat Prod* 2020;83(11):3372–3380
- 105 Girich EV, Yurchenko AN, Smetanina OF, et al. Neuroprotective metabolites from vietnamese marine derived fungi of *Aspergillus* and *Penicillium* Genera. *Mar Drugs* 2020;18(12):608
- 106 Shaker S, Fan RZ, Li HJ, Lan WJ. A pair of novel bisindole alkaloid enantiomers from marine fungus *Fusarium* sp. XBB-9. *Nat Prod Res* 2021;35(09):1497–1503
- 107 Yan W, Zhao SS, Ye YH, et al. Generation of indoles with agrochemical significance through biotransformation by *Chaetomium globosum*. *J Nat Prod* 2019;82(08):2132–2137
- 108 Campana R, Favi G, Baffone W, Lucarini S. Marine alkaloid 2,2-bis(6-bromo-3-indolyl) ethylamine and its synthetic derivatives inhibit microbial biofilms formation and disaggregate developed biofilms. *Microorganisms* 2019;7(02):28
- 109 Anjum K, Kaleem S, Yi W, Zheng G, Lian X, Zhang Z. Novel antimicrobial indolepyrazines A and B from the marine-associated *Acinetobacter* sp. ZZ1275. *Mar Drugs* 2019;17(02):89
- 110 Yuan MX, Qiu Y, Ran YQ, et al. Exploration of indole alkaloids from marine fungus *Pseudallescheria boydii* F44-1 using an amino acid-directed strategy. *Mar Drugs* 2019;17(02):77
- 111 Zhang P, Li XM, Mao XX, Mándi A, Kurtán T, Wang BG. Varioloid A, a new indolyl-6,10b-dihydro-5aH-[1]benzofuro[2,3-b]indole derivative from the marine alga-derived endophytic fungus *Paecilomyces variotii* EN-291. *Beilstein J Org Chem* 2016;12:2012–2018
- 112 Zhang P, Li XM, Mao XX, Mándi A, Kurtán T, Wang BG. Correction: varioloid A, a new indolyl-6,10b-dihydro-5aH-[1]benzofuro[2,3-b]indole derivative from the marine alga-derived endophytic fungus *Paecilomyces variotii* EN-291. *Beilstein J Org Chem* 2018;14:2394–2395
- 113 Zhang D, Yi W, Ge H, Zhang Z, Wu B. A new antimicrobial indoloditerpene from a marine-sourced fungus *aspergillus versicolor* ZZ761. *Nat Prod Res* 2021;35(18):3114–3119
- 114 Elsbaey M, Tanaka C, Miyamoto T. New secondary metabolites from the mangrove endophytic fungus *Aspergillus versicolor*. *Phytochem Lett* 2019;32:70–76
- 115 Liu L, Xu W, Li S, et al. Penicindopene A, a new indole diterpene from the deep-sea fungus *Penicillium* sp. YPCMAC1. *Nat Prod Res* 2019;33(20):2988–2994
- 116 Harms H, Rempel V, Kehraus S, et al. Indoloditerpenes from a marine-derived fungal strain of *Dichotomomyces cejpui* with antagonistic activity at GPR18 and cannabinoid receptors. *J Nat Prod* 2014;77(03):673–677
- 117 Qiao MF, Ji NY, Liu XH, Li K, Zhu QM, Xue QZ. Indoloditerpenes from an algicolous isolate of *Aspergillus oryzae*. *Bioorg Med Chem Lett* 2010;20(19):5677–5680
- 118 Chen MY, Xie QY, Kong FD, et al. Two new indole-diterpenoids from the marine-derived fungus *Penicillium* sp. KFD28. *J Asian Nat Prod Res* 2021;23(11):1030–1036
- 119 Kong FD, Fan P, Zhou LM, et al. Peneperenes A–D, four indole terpenoids with potent protein tyrosine phosphatase inhibitory activity from the marine-derived fungus *Penicillium* sp. KFD28. *Org Lett* 2019;21(12):4864–4867
- 120 Zhou LM, Kong FD, Fan P, et al. Indole-diterpenoids with protein tyrosine phosphatase inhibitory activities from the marine-derived fungus *Penicillium* sp. KFD28. *J Nat Prod* 2019;82(09):2638–2644
- 121 Guo XC, Xu LL, Yang RY, et al. Anti-Vibrio indole-diterpenoids and C-25 epimeric steroids from the marine-derived fungus *Penicillium janthinellum*. *Front Chem* 2019;7:80
- 122 Ivanets EV, Yurchenko AN, Smetanina OF, et al. Asperindoles A–D and a p-terphenyl derivative from the ascidian-derived fungus *Aspergillus* sp. KMM 4676. *Mar Drugs* 2018;16(07):232
- 123 Gao SS, Li XM, Williams K, Proksch P, Ji NY, Wang BG. Rhizovarin A–F, indole-diterpenes from the mangrove-derived endophytic fungus *Mucor irregularis* QEN-189. *J Nat Prod* 2016;79(08):2066–2074
- 124 Zhang P, Li XM, Li X, Wang BG. New indole-diterpenoids from the algal-associated fungus *Aspergillus nidulans*. *Phytochem Lett* 2015;12:182–185
- 125 Sun K, Li Y, Guo L, Wang Y, Liu P, Zhu W. Indole diterpenoids and isocoumarin from the fungus, *Aspergillus flavus*, isolated from the prawn, *Penaeus vannamei*. *Mar Drugs* 2014;12(07):3970–3981
- 126 Christian OE, Compton J, Christian KR, Mooberry SL, Valeriote FA, Crews P. Using jasplakinolide to turn on pathways that enable the isolation of new chaetoglobosins from *Phomopsis asparagi*. *J Nat Prod* 2005;68(11):1592–1597
- 127 Cui CM, Li XM, Li CS, Proksch P, Wang BG. Cytochalasins A–G, cytochalasans from a marine-derived endophytic fungus, *Chaetomium globosum* QEN-14. *J Nat Prod* 2010;73(04):729–733
- 128 Luo XW, Gao CH, Lu HM, et al. HPLC-DAD-guided isolation of diversified chaetoglobosins from the coral-associated fungus *Chaetomium globosum* C2F17. *Molecules* 2020;25(05):1237
- 129 Zhang Z, Min X, Huang J, et al. Cytochalasins H and I, New antiproliferative cytochalasans from deep-sea-derived fungus *Chaetomium globosum*. *Mar Drugs* 2016;14(12):233
- 130 Carr G, Tay W, Bottrill H, Andersen SK, Mauk AG, Andersen RJ. Plectosphaeric acids A, B, and C, indoleamine 2,3-dioxygenase inhibitors produced in culture by a marine isolate of the fungus *Plectosphaerella cucumerina*. *Org Lett* 2009;11(14):2996–2999
- 131 Shanuja SK, Iswarya S, Gnanamani A. Marine fungal DHICA as a UVB protectant: assessment under in vitro and in vivo conditions. *J Photochem Photobiol B* 2018;179:139–148
- 132 Mourshid SS, Badr JM, Risinger AL, Mooberry SL, Youssef DT. Penicilloitins A and B, new antimicrobial fatty acid esters from a marine endophytic *Penicillium* species. *Z Natforsch C J Biosci* 2016;71(11–12):387–392
- 133 Meng LH, Chen HQ, Form I, Konuklugil B, Proksch P, Wang BG. New chromone, isocoumarin, and indole alkaloid derivatives from three sponge-derived fungal strains. *Nat Prod Commun* 2016;11(09):1293–1296
- 134 Dong JJ, Bao J, Zhang XY, Xu XY, Nong XH, Qi SH. Alkaloids and citrinins from marine-derived fungus *Nigrospora oryzae* SCSGAF 0111. *Tetrahedron Lett* 2014;55(16):2749–2753
- 135 Afyatullov SSH, Zhuravleva OI, Antonov AS, et al. New metabolites from the marine-derived fungus *Aspergillus fumigatus*. *Nat Prod Commun* 2012;7(04):497–500