

Rate of Revision Hip Arthroscopy for Pediatric Femoroacetabular Impingement: A Systematic Review

Edward S. Mojica, BS¹ Teren Yedikian, BS¹ Brittany DeClouette, MD¹ Eoghan T. Hurley, MB, BCh¹
Aaron Gipsman, MD¹ Pablo Castañeda, MD¹ Thomas Youm, MD¹

¹Department of Orthopaedic Surgery, Division of Sports Medicine, NYU Langone Health, New York, New York

Address for correspondence Edward S. Mojica, BS, 333 E38th, NYU Langone Health, New York, NY 10015
(e-mail: edward.mojica@nyulangone.org).

J Hip Surg 2022;6:35–42.

Abstract

Femoroacetabular impingement (FAI) can cause pain, dysfunction, and early arthritic progression in young patients. The purpose of this study was to systematically review the evidence in literature to determine patient-reported outcomes and failure rates as defined by the need for revision surgery, following hip arthroscopy for pediatric patients with FAI. The literature search was performed based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. Clinical studies evaluating the outcomes following primary hip arthroscopy for pediatric patients with FAI were included. Clinical outcomes evaluated included revisions, complications, functional outcome scores (modified Hip Harris Score [mHHS], Non-Arthritis Hip Score, and Visual Analogue Score), and return to play. Statistical analysis was performed using GraphPad Prism version 7. This study is a level IV systematic review. Overall, 20 clinical studies with 1,136 patients (1,223 hips) were included in this review, with an average age of 16.3 years. Overall, 8.6% patients experienced revision surgery. The mHHS was the most widely used metric, present in 17 of the 20 studies. The mHHS was reported as excellent (> 90) in six of these studies and good (80–89) in 11. The weighted mean of the post-operative mHHS found across reporting studies was 84.3, from a baseline score of 58.1. The overall return to play rate was 91%. This study reports excellent post-hip arthroscopy clinical outcomes for FAI and labral tears in the pediatric population. However, revision rates for this surgical procedure are higher than previously documented.

Keywords

- ▶ FAI
- ▶ pediatric
- ▶ systematic review
- ▶ revision

Femoroacetabular impingement (FAI) can cause pain, dysfunction, and early arthritic progression in young patients. FAI refers to the morphological abnormality of the bone on the acetabular rim (pincer lesion) and the femoral neck (cam lesion) or a combination of the two.^{1–4} Labral tears are common in FAI due to repetitive abnormal contact between the bone and the chondrolabral junction, occurring in up to 55% of patients with mechanical symptoms.⁵ FAI can affect pediatric patients, with youth sports being a

risk factor in its development due to excessive stress on the growing hip.^{6,7}

Hip arthroscopy, while technically challenging, is increasingly being performed to treat FAI and correct the pathological morphology.^{8–10} However, the evidence supporting the use of hip arthroscopy for FAI in pediatric patients is limited, with prophylactic surgery in the asymptomatic hip being contraindicated, even in the context of predisposing anatomy.¹¹ Additionally, many patients in the pediatric

received
July 7, 2021
accepted after revision
December 17, 2021
published online
February 4, 2022

© 2022, Thieme. All rights reserved.
Thieme Medical Publishers, Inc.,
333 Seventh Avenue, 18th Floor,
New York, NY 10001, USA

DOI <https://doi.org/10.1055/s-0041-1742280>
ISSN 2472-8446.

population can be managed non-operatively quite successfully.¹² Despite this, there is a bias not only in the clinic but also in the literature toward the surgical management of FAI, with further bias toward older demographics. The majority of the literature consists of retrospective case series and studies focusing primarily on surgical management of adult patients.

Chen et al¹³ performed a systematic review looking at adolescent return to play following hip arthroscopy for FAI. They found 10 studies that included 618 patients, demonstrated a return to play rate of 84.9%, and had a 3.1% revision rate. However, multiple other studies that did not report the rate of return to play were excluded in that review. This must be considered when evaluating overall outcomes and failure rates for this population. Additionally, Migliorini and Maffulli¹⁴ found among 10 studies and 406 pediatric patients that there was a 4.7% revision rate in their systematic review; however, this study failed to include many of the available studies in the literature.

The purpose of this study was to systematically review the current published evidence to determine patient-reported outcomes and failure rates following hip arthroscopy for pediatric patients with FAI. Our hypothesis was that there would be excellent patient-reported outcomes with a low failure rate.

Methods

Search Strategy and Study Selection

Two independent reviewers conducted the literature search based on the Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. The MEDLINE, EMBASE, and Cochrane Library databases were queried in March 2021 using the following search terms: (femoroacetabular or FAI or hip or coxa or acetabulofemoral joint) and (impingement) and (repair or refixation or preservation or reattachment or debridement or resection). The independent authors reviewed articles from the search for the relevance of this particular study, with the lead author deciding upon inclusion in cases of disagreement. Upon reviewing the abstract, papers that were deemed relevant to the research question were then thoroughly reviewed. The reference lists of articles considered suitable for this study were then manually screened for additional works that did not arise in the primary search.

Eligibility Criteria

The inclusion criteria for this analysis were as follows: (1) study analyzing outcomes of patients undergoing hip arthroscopy for FAI who are less than 18 years of age, (2) published in a peer-reviewed journal, (3) published in English, (4) full text assessment of studies available, and (5) having a minimum follow-up time of 6 months. Study designs including randomized controlled trials, prospective cohort studies, retrospective cohort studies, and case-control studies were considered for inclusion. The exclusion criteria included (1) review studies, (2) cadaver studies, (3) biomechanical studies, and (4) conference abstract only.

Data Extraction/Analysis

Information was pulled from the selected studies using a data sheet predetermined by the research team. Data included study design, level of evidence (LOE), the methodological quality of evidence (MQOE), patient population and demographics, and outcomes, including patient-reported and surgical measures. The modified Coleman scale was used to compute the MQOE for each study.¹⁵ The modified Coleman score is scaled from 0 to 100 points, with an increasing score denoting a study that better avoids chance, biases, and confounding factors. The Coleman score components are modeled after the Consolidated Standards of Reporting Trials statement, with modifications to allow for the analysis of various study types.

The reasons for revision were also collected across the various studies analyzed. As the reasons were reported with different verbiage depending on the author of study, several factors were selected by the authors as agreed upon after the review of the literature. These factors included: FAI recurrence, capsulolabral adhesions, overuse, continuous pain, re-injury, snapping hip, heterotopic ossification, and inability to return to the same level of activity. A frequently encountered term was “recurrence of FAI symptoms,” which was decided by the authors to be synonymous with continuous pain.

Statistical Analysis

Statistical analysis was performed using GraphPad Prism version 7. Qualitative analysis was performed for each study, and quantitative analysis was performed across all groups.

Results

Literature Search

The initial literature search resulted in 2,607 total studies. Once duplicates were removed and the articles were screened for inclusion and exclusion criteria, 1,384 studies were included, and full texts were assessed for eligibility. Overall, 20 clinical studies with 1,136 patients (1,223 hips) were included in this review (► Fig. 1).^{13,16–35}

Study Characteristics and Patient Demographics

Overall, 20 studies (LOE II: 2, LOE III: 5, and LOE IV: 14) reported the outcomes of hip arthroscopy for FAI in the pediatric population. The mean modified Coleman's was 56.1. The mean age of the studied population was 16 years (12–18 years). Patients undergoing FAI surgery had a mean α angle of 61.0 ± 10.4 and a mean lateral center edge angle (LCEA) of 30.3 ± 8.5 . The study characteristics and patient demographics are shown in ► Table 1.

Patient-Reported Outcomes

Overall, weighted means of patient follow-up demonstrated a range from 11.5 to 47.3 months, with a weighted average follow-up of 38.2 months. In the studies analyzed, modified Hip Harris Score (mHHS), Non-Arthritis Hip Score (NAHS), and Visual Analogue Score (VAS) were the most commonly reported.




Fig. 1 PRISMA study selection flow diagram.

The mHHS was the most widely used metric, present in 17 of the 20 studies. The mHHS was reported as excellent (> 90) in six of these studies and good (80–89) in 11. The weighted mean of the postoperative mHHS found across reporting studies was found to be 84.3 from a baseline score of 58.2. In all of these reporting studies, there was a significant increase in pre-operative values, meeting minimal clinically important difference (MCID) and patient acceptable symptomatic state for the values reported in the literature.³⁶ Although the average patient studied reached the threshold of change in mHHS that qualifies for MCID at final follow-up, Beck et al, Cvetanovich et al, and Larson et al further quantify the rate at which the studied patient population achieved MCID at 88.4, 84.0, and 81.2% of their population meeting the threshold, respectively.

NAHS was also reported in five studies, with three studies reporting excellent outcomes (> 90) and two studies reporting good outcomes (80–89). Baseline score for NAHS through weighted means was found to be 71, rising to 95 upon final

follow-up. VAS was reported in six studies, with all studies demonstrating a significant decrease in pain post-operatively. VAS was found to have a weighted mean baseline score of 5.9, recovering to 1.8 upon final follow-up. These results are further illustrated in ►Table 2.

Return to Sport

Return to sport among adolescents undergoing hip arthroscopy was investigated in six of the analyzed studies. Upon final follow-up, the return to play rate was determined to be 91%. Additionally, Willimon et al studied the amount of time it took for the pediatric population to return to school, finding a mean length of 10.8 days (standard deviation 5.1). These results are further illustrated in ►Table 3.

Complications and Revisions

The most common complication reported after FAI surgery was a neurological deficit. Transient numbness and nerve paresthesia most frequently occurred along the perineal and

Table 1 Study characteristics

Author	LOE	Coleman	Patients (n)	Hips (n)	LCEA (degrees)	Alpha angle (degrees)	Age (y)	Follow-up (mo)
Arashi et al	3	61	33	36	32	63	16.7 (12–17)	40 and 60
Ashberg et al	2	75	157	157	29.2	58.3	16.6 (13.2–17.9)	38.5 (24–89.9)
Beck et al	4	59	85	85	30.1 ± 6	63.6 ± 12.2	17.60	60
Byrd et al	4	72	104	116		59.9	16.00	38 (24–120)
Chandrasekaran et al	4	61	90	90	31.6	59.5	16.30	30.6 (24.1–60)
Cvetanovich et al	4	62	37	37	32.2 ± 4.8	59.4 ± 7.5	17 (12.8–18.3)	28.3
Degen et al	3	43	34	38	33 ± 5.1	65.2 ± 7.1	15.9 (10.8–18.0)	36.1
Fabricant et al	4	54	21	27		64 ± 16	17.60	18 (12–30)
Larson et al	4	52	28	37	30.7	60.9	15.90	39.8 (12.1–86.5)
Litrenta et al	4	64	69	81	28.5 ± 18.4	57.7 ± 11.6	15.9 (14–18)	45.2
McConkey et al	2	69	24	36			15.70	24
Menge et al	4	75	60	70	38 ± 5	67 ± 14	16.00	120
Mohan et al	4	37	50	57	31	54	17 (13–23)	34 (24–77)
Newman et al	3	51	42	42	33 ± 9	68 ± 14	16 (10–19)	43
Nwachukwu et al	4	37	47	47	31.2 ± 6.5	57 ± 12.4	16.5 (11–18)	12
Philippon et al	4	66	60	65	36	64	15.00	36 (24–60)
Richard et al	2	49	58	58			15.53 (10.4–20.5)	6
Tran et al	4	57	34	41			15.70	14 (12–24)
Willimon et al	4	26	17	17			15.30	
Youngman et al	3	48	86	86	31.6	61.7	16.50	37 (12–121)

Abbreviations: LCEA, lateral center edge angle; LOE, level of evidence.

Table 2 Patient-reported outcomes

Author	Pre-op mHHS	Post-op mHHS	p-Value	MCID for mHHS	Pre-op NAHS	Post-op NAHS	p-Value	Pre-op VAS	Post-op VAS	p-Value
Arashi et al		98				97				
Ashberg et al	64.8	84.1	$p < 0.001$,		65.9	99.4	$p < 0.001$,	5.80	1.9	0.001
Beck et al	58.9	85.1	$p < 0.001$,	88.40%			$p < 0.001$,	6.79	2.44	0.001
Byrd et al	69	94	$p < 0.001$,							
Chandrasekaran et al	64.5	89.6	$p < 0.001$,		65.3	89.9	$p < 0.001$,	6.07	2.16	0.001
Cvetanovich et al	58.1	86.9	$p < 0.001$,	84.00%						
Degen et al	63.8	86	$p < 0.001$,							
Fabricant et al	67	88	$p < 0.001$,							
Larson et al	66.8	94.5	$p < 0.001$,	81.20%				5.90	1.2	0.001
Litrenta et al	64.6	88.1	$p < 0.001$,		66.8	89.8	$p < 0.001$,	5.50	1.6	0.001
McConkey et al										
Menge et al	56	88	< 0.0001							
Mohan et al	64	85	0.04							
Newman et al	57.5	84.2								

Table 2 (Continued)

Author	Pre-op mHHS	Post-op mHHS	p-Value	MCID for mHHS	Pre-op NAHS	Post-op NAHS	p-Value	Pre-op VAS	Post-op VAS	p-Value
Nwachukwu et al	61.6	90	< 0.0001							
Philippon et al	57	91	0.005							
Richard et al										
Tran et al	77.39	94.15	< 0.0005		76.34	93.18	< 0.0005			
Willimon et al										
Youngman et al	62.8	86.5	< 0.0001							

Abbreviations: MHHS, modified Hip Harris Score; NAHS, Non-Arthritis Hip score; post-op, postoperative; pre-op, preoperative; VAS, Visual Analogue Score. Note: Bold values highlight statistical significance.

Table 3 Return to play of selected studies

Author	Athletes (n)	RTP total—return to play (%)
Cvetanovich et al	29	29 (100)
Larson et al	28	26 (93)
Litrenta et al	69	58 (84)
McConkey et al	24	24 (100)
Mohan et al	50	46 (92)
Tran et al	32	29 (91)

lateral femoral cutaneous nerves. Litrenta et al described four cases of temporary unspecified numbness occurring in 81 of the operated upon hips. McConkey et al and Byrd et al reported two transient left femoral cutaneous and two transient peri-

neal nerve paresthesias, respectively. There were few infections, with Litrenta et al reporting two minor infections. There were a total of 82 revisions in the literature. Therefore, 8.6% of the 959 hips in those studies investigating revision rates experienced revision surgery, at a mean time of 23.7 months. Revision procedures were qualified as labral repair, capsular release/plication, lysis of adhesions, and removal of loose bodies most commonly, although they were scarcely specified upon in the studies analyzed. The revisions and complications are further reported in ► **Table 4**.

The most common reasons for revision surgery were continuous pain and re-injury of the hip, with incidences of 14 (17%) and 17 (21%), respectively. Additionally, 5.0% of those that underwent revision surgery for the recurrence of FAI morphology, such as cam regrowth. These revisions are further illustrated in ► **Table 5**.

Table 4 Complications and revisions

Author	Complications	Revision	Time until revision (mean[range])
Arashi et al	0	6	19.5
Ashberg et al		20	
Beck et al	Two transient nerve paresthesia along perineal nerve	2	
Byrd et al	0	4	17.8 (6.9–41.5)
Chandrasekaran et al	One case of pudendal neuropraxia; one wound dehiscence	5	13.4 (8.2–22.1)
Cvetanovich et al	0		
Degen et al	0	2	
Litrenta et al	Two transient nerve paresthesia along lateral cutaneous nerve	6	37.3 (7.9–74.2)
McConkey et al		0	
Menge et al		7	
Mohan et al		3	
Newman et al		6	
Philippon et al		8	26 (10–46)
Tran et al	0		
Youngman et al	0	10	

Note: Time until revision given in months.

Table 5 Reasons for revisions

Reason for revision	n (%)
FAI recurrence	4 (5)
Capsulolabral adhesions	9 (11)
Overuse	3 (4)
Continuous pain	14 (17)
Re-injury	17 (21)
Snapping hip	4 (5)
Heterotopic ossification	3 (4)
Inability to return to the same level	3 (4)
Instability	3 (4)
Unknown	22

Discussion

The most important finding in this review is a revision rate of 8.6% for hip arthroscopy of the pediatric patient, which is higher than previous systematic reviews.^{13,37} However, while revision rates were noted to be higher, the pediatric patient demonstrated good, if not excellent, outcomes, as reported by mHHS and NAHS. These outcomes were shown to be significantly improved throughout all 17 studies that looked at baseline and post-operative scores for mHHS and in the five studies that reported NAHS. Similarly, six studies found both a significant decrease in VAS scores and a high return-to-play rate among the athletes that were observed. Nonetheless, it is important to note that while reported outcomes favor arthroscopy, the technical demands of this procedure present various theoretical risks.

Hip arthroscopy in the pediatric population is known to be a particularly challenging procedure, mainly due to anatomical variation in the skeletally immature patient.^{2,8,37} A significant decrease in the neck-shaft angle and anteversion must be kept in mind when positioning the patient and care must be taken to respect the open physes. Additionally, due to the immaturity of the acetabulum, theoretical risks of slipped capital femoral epiphysis (SCFE), avascular necrosis, and infection do exist. However, a systematic review by de Sa et al reported no violation of the physes resulting in premature closures, SCFE, or other theoretical and feared complications of arthroscopy of the pediatric hip. This postulates the idea that these risks might solely be theoretical. Furthermore, of the 435 hips studied, only 13 (3.0%) went on to require revision surgery, primarily for lysis of adhesions. Outcomes demonstrated across the review also favored the use of arthroscopy, with significant improvements in patient-reported outcomes among the six studies analyzed.

The findings of this paper are in accordance with the previous literature and continue to affirm good clinical outcomes. Similar to the review by de Sa et al, Chen et al¹³ conducted a systematic review focusing on return to sport and also found high post-operative outcome scores. Additionally, of 194 hips analyzed, a return-to-sport rate of 84.9%

was reported upon final follow-up, demonstrating tangible results in addition to reported good outcomes. Despite good scores, there seems to be a difference in percentages of the pediatric population achieving MCID as compared with the adult population. Overall, 97% of the adult population was found to reach MCID for mHHS in a systematic review by Levy et al,³⁸ leaving the rates in the 80s found by this study to be relatively low. This is perhaps due the adolescent population having less severe disease, and therefore, it being more difficult to achieve a significant increase from baseline than their older peers. However, Chen et al reported a revision rate of 3.1%, drawn from six separate studies, which is in line with the findings from de Sa et al. Additionally, Migliorini and Maffulli¹⁴ found among 10 studies and 406 patients that there was a 4.7% revision rate in their systematic review; however, this study failed to include many of the available studies in the literature. The revision rate presented in our paper (8.6%) is higher than rates previously documented, resulting from data including a larger number of hips studied, and thus is more indicative of the population and literature as a whole.

As documented by Litrenta et al and Chandrasekaran et al,^{20,25} the primary reasons for revision were recurrence of pain and re-injury. Pediatric populations who undergo surgery are less likely to rest their hip and can do subsequent damage to the vulnerable hip. This was commented on by Ashberg et al¹⁷ who discussed the pediatric populations proclivity toward revision surgery may be due to early return to unrestricted activity. Byrd et al¹⁹ made a similar observation, noting that athleticism in the pediatric population may correlate with the incidence of FAI, in general, reporting that 96% of those patients in their adolescent group were athletes compared with 61% of adults. While adolescents tend to be more active than adults, Menge et al compounded on this rationale stating that of the seven revision surgeries observed all were female and athletes, theorizing that increased athleticism in the pediatric population may lend itself to the need for revision surgery due to re-tear of the labrum or aggravation of the joint. They also documented that hypermobility as reported by Beighton score (≥ 4) was associated with revision surgery ($p = 0.045$), warranting further studies.²⁷ Increased return to unrestricted activity may contribute to this high revision rate reported, especially, in the context of 4% of adults going back for secondary surgery.³⁹

Interestingly, while re-injury and overuse seem to be the primary cause of revisions throughout the investigated literature, Arashi et al.¹⁶ documented four of their six revisions due to cam regrowth in the male population. Although a rare cause of the need for revision in this review, it has been theorized that cam regrowth is a theoretical risk in the skeletally immature patients, due to bony growth potential inherent in this younger population, and was indeed observed in 5% of the studied population.^{6,7} At risk for this pathology are those with initial cam morphology, seen in pediatric patients with borderline α angles of 60.1 ± 10.4 , as well as more commonly in males.^{3,7,16} These at-risk populations become important when determining ideal patient selection, as outcomes seem to correlate.

There are various risk factors that predispose to poorer outcomes, such as dysplasia, older age, and abnormal femoral version. However, the most significant risk factor is the need for revision surgery.^{3,9,29,40,41} Newman et al²⁹ demonstrated in a retrospective cohort study that patients undergoing revision hip arthroscopy for labral tears and FAI had significantly lower improvement and post-operative mHHS scores. These patients also demonstrated poorer sports function (as evaluated by Hip Outcome Score, sports) during the final post-operative visit. This suggests that while pediatric hip arthroscopy indicated for labral tears has excellent clinical outcomes, the need for revision surgery can easily jeopardize the operation's success and is typically triggered with early return to unrestricted activity. Furthermore, as this is a pediatric population, studies with longer-term follow-up are needed to ascertain the true failure rate over time, specifically evaluating if these patients suffer from premature arthritic hip degeneration.

Limitations

This study has several limitations and biases, largely due, in part, to the variable studies reviewed themselves. First, the age range was 12 to 18 years and some patients may be skeletally mature, which may impact the procedure outcomes and it was not possible to sub-stratify by this. While this study revolved around revision rate, certain studies did not report revision rates due to exclusion criteria and others still did not report why the revision was performed. Additionally, studies were included that had less than 1 full year of follow-up. The heterogeneity of the studies was also not evaluated because of the lack of comparative groups. Furthermore, apart from mHHS, there was a lack of consistency in the outcomes used making it difficult to compare across the different studies.

Conclusion

Our study established that clinical outcomes following hip arthroscopy for FAI and labral tears in the pediatric population are excellent. However, revision rates for this surgical procedure might be higher than previously documented. While hip arthroscopy overall is safe and effective in this population, the potential need for revision surgery should be discussed pre-operatively.

Conflict of Interest

None declared.

References

- Cetinkaya S, Tokar B, Ozden VE, Dikmen G, Taser O. Arthroscopic labral repair versus labral debridement in patients with femoroacetabular impingement: a minimum 2.5 year follow-up study. *Hip Int* 2016;26(01):20–24
- Espinosa N, Rothenfluh DA, Beck M, Ganz R, Leunig M. Treatment of femoro-acetabular impingement: preliminary results of labral refixation. *J Bone Joint Surg Am* 2006;88(05):925–935
- Fukui K, Briggs KK, Trindade CA, Philippon MJ. Outcomes after labral repair in patients with femoroacetabular impingement and borderline dysplasia. *Arthroscopy* 2015;31(12):2371–2379
- Jayakumar P, Ramachandran M, Youm T, Achan P. Arthroscopy of the hip for paediatric and adolescent disorders: current concepts. *J Bone Joint Surg Br* 2012;94(03):290–296
- McCarthy JC, Noble PC, Schuck MR, Wright J, Lee J. The Otto E. Aufranc Award: the role of labral lesions to development of early degenerative hip disease. *Clin Orthop Relat Res* 2001;(393):25–37
- Akel I, Songür M, Karahan S, Yilmaz G, Demirkıran HG, Tümer Y. Acetabular index values in healthy Turkish children between 6 months and 8 years of age: a cross-sectional radiological study. *Acta Orthop Traumatol Turc* 2013;47(01):38–42
- Kuo FC, Kuo SJ, Ko JY. Overgrowth of the femoral neck after hip fractures in children. *J Orthop Surg Res* 2016;11(01):50
- Awad MAH, Bajwa AK, Slaunwhite E, Logan KJ, Wong IH. Indications for hip arthroscopy in pediatric patients: a systematic review. *J Hip Preserv Surg* 2019;6(04):304–315
- Shapira J, Kyin C, Go C, et al. Indications and outcomes of secondary hip procedures after failed hip arthroscopy: a systematic review. *Arthroscopy* 2020;36(07):1992–2007
- Nasser R, Domb B. Hip arthroscopy for femoroacetabular impingement. *EFORT Open Rev* 2018;3(04):121–129
- Wylie JD, Kim Y-J. The natural history of femoroacetabular impingement. *J Pediatr Orthop* 2019;39(6, Supplement 1 Suppl 1):S28–S32
- Pennock AT, Bomar JD, Johnson KP, Randich K, Upasani VV. Nonoperative management of femoroacetabular impingement: a prospective study. *Am J Sports Med* 2018;46(14):3415–3422
- Chen SL, Maldonado DR, Go CC, Kyin C, Lall AC, Domb BG. Outcomes of hip arthroscopic surgery in adolescents with a sub-analysis on return to sport: a systematic review. *Am J Sports Med* 2020;48(06):1526–1534
- Migliorini F, Maffulli N. Arthroscopic management of femoroacetabular impingement in adolescents: a systematic review. *Am J Sports Med* 2021;49(13):3708–3715
- Jakobsen RB, Engebretsen L, Slauterbeck JR. An analysis of the quality of cartilage repair studies. *J Bone Joint Surg Am* 2005;87(10):2232–2239
- Arashi T, Murata Y, Utsunomiya H, et al. Higher risk of cam regrowth in adolescents undergoing arthroscopic femoroacetabular impingement correction: a retrospective comparison of 33 adolescent and 74 adults. *Acta Orthop* 2019;90(06):547–553
- Ashberg L, Walsh JP, Yuen LC, Perets I, Chaharbakhshi EO, Domb BG. Outcomes of hip arthroscopy in adolescents: a comparison of acute versus chronic presentation. Two-year minimum follow-up. *J Pediatr Orthop* 2018;38(02):e50–e56
- Beck EC, Nwachuckwu BU, Jan K, Nho SJ. Hip arthroscopy for femoroacetabular impingement syndrome in adolescents provides clinically significant outcome benefit at minimum 5-year follow-up. *Arthroscopy* 2021;37(05):1467–1473.e2. Doi: 10.1016/j.arthro.2020.12.188. Epub: 2020 Dec 24. PMID: 33359159
- Byrd JW, Jones KS, Gwathmey FW. Arthroscopic management of femoroacetabular impingement in adolescents. *Arthroscopy* 2016;32(09):1800–1806
- Chandrasekaran S, Darwish N, Chaharbakhshi EO, Lodhia P, Suarez-Ahedo C, Domb BG. Arthroscopic treatment of labral tears of the hip in adolescents: patterns of clinical presentation, intra-articular derangements, radiological associations and minimum 2-year outcomes. *Arthroscopy* 2017;33(07):1341–1351
- Cvetanovich GL, Weber AE, Kuhns BD, et al. Clinically meaningful improvements after hip arthroscopy for femoroacetabular impingement in adolescent and young adult patients regardless of gender. *J Pediatr Orthop* 2018;38(09):465–470
- Degen RM, Mayer SW, Fields KG, Coleman SH, Kelly BT, Nawabi DH. Functional outcomes and cam recurrence after arthroscopic treatment of femoroacetabular impingement in adolescents. *Arthroscopy* 2017;33(07):1361–1369

- 23 Fabricant PD, Heyworth BE, Kelly BT. Hip arthroscopy improves symptoms associated with FAI in selected adolescent athletes. *Clin Orthop Relat Res* 2012;470(01):261–269
- 24 Larson CM, Giveans MR, Stone RM. Arthroscopic debridement versus refixation of the acetabular labrum associated with femoroacetabular impingement: mean 3.5-year follow-up. *Am J Sports Med* 2012;40(05):1015–1021
- 25 Litrenta J, Mu BH, Ortiz-Declet V, et al. Response to hip arthroscopy successfully treats femoroacetabular impingement in adolescent athletes. *J Pediatr Orthop* 2021;41(01):e98–e99
- 26 McConkey MO, Chadayammuri V, Garabekyan T, Mayer SW, Kraeutler MJ, Mei-Dan O. Simultaneous bilateral hip arthroscopy in adolescent athletes with symptomatic femoroacetabular impingement. *J Pediatr Orthop* 2019;39(04):193–197
- 27 Menge TJ, Briggs KK, Dornan GJ, McNamara SC, Philippon MJ. Survivorship and outcomes 10 years following hip arthroscopy for femoroacetabular impingement: labral debridement compared with labral repair. *J Bone Joint Surg Am* 2017;99(12):997–1004
- 28 Mohan R, Johnson NR, Hevesi M, Gibbs CM, Levy BA, Krych AJ. Return to sport and clinical outcomes after hip arthroscopic labral repair in young amateur athletes: minimum 2-year follow-up. *Arthroscopy* 2017;33(09):1679–1684
- 29 Newman JT, Briggs KK, McNamara SC, Philippon MJ. Outcomes after revision hip arthroscopic surgery in adolescent patients compared with a matched cohort undergoing primary arthroscopic surgery. *Am J Sports Med* 2016;44(12):3063–3069
- 30 Nwachukwu BU, Chang B, Kahlenberg CA, et al. Arthroscopic treatment of femoroacetabular impingement in adolescents provides clinically significant outcome improvement. *Arthroscopy* 2017;33(10):1812–1818
- 31 Philippon MJ, Yen YM, Briggs KK, Koppersmith DA, Maxwell RB. Early outcomes after hip arthroscopy for femoroacetabular impingement in the athletic adolescent patient: a preliminary report. *J Pediatr Orthop* 2008;28(07):705–710
- 32 Richard HM, Cerza SP, De La Rocha A, Podeszwa DA. Preoperative mental health status is a significant predictor of postoperative outcomes in adolescents treated with hip preservation surgery. *J Child Orthop* 2020;14(04):259–265
- 33 Tran P, Pritchard M, O'Donnell J. Outcome of arthroscopic treatment for cam type femoroacetabular impingement in adolescents. *ANZ J Surg* 2013;83(05):382–386
- 34 Willimon SC, Johnson MM, Herzog MM, Busch MT. Time to return to school after 10 common orthopaedic surgeries among children and adolescents. *J Pediatr Orthop* 2019;39(06):322–327
- 35 Youngman TR, Wagner KJ III, Montanez B, et al. The association of α angle on disease severity in adolescent femoroacetabular impingement. *J Pediatr Orthop* 2021;41(02):88–92
- 36 Chahal J, Thiel GSV, Mather RC, Lee S, Salata MJ, Nho SJ. The minimal clinically important difference (MCID) and patient acceptable symptomatic state (PASS) for the modified harris hip score and hip outcome score among patients undergoing surgical treatment for femoroacetabular impingement. *Orthop J Sports Med* 2014;2(2 Suppl):2325967114S00105. Doi: 10.1177/2325967114S00105
- 37 de Sa D, Cargnelli S, Catapano M, et al. Femoroacetabular impingement in skeletally immature patients: a systematic review examining indications, outcomes, and complications of open and arthroscopic treatment. *Arthroscopy* 2015;31(02):373–384
- 38 Levy DM, Kuhns BD, Chahal J, Philippon MJ, Kelly BT, Nho SJ. Hip arthroscopy outcomes with respect to patient acceptable symptomatic state and minimal clinically important difference. *Arthroscopy* 2016;32(09):1877–1886
- 39 West CR, Bedard NA, Duchman KR, Westermann RW, Callaghan JJ. Rates and risk factors for revision hip arthroscopy. *Iowa Orthop J* 2019;39(01):95–99
- 40 Nepple JJ, Carlisle JC, Nunley RM, Clohisy JC. Clinical and radiographic predictors of intra-articular hip disease in arthroscopy. *Am J Sports Med* 2011;39(02):296–303
- 41 Nicholls AS, Kiran A, Pollard TC, et al. The association between hip morphology parameters and nineteen-year risk of end-stage osteoarthritis of the hip: a nested case-control study. *Arthritis Rheum* 2011;63(11):3392–3400