

¹⁸F-FDG Whole-Body PET-MR in Primary Hepatic Lymphoma Mimicking Focal Nodular Hyperplasia

Shyma Basheer¹ Dinesh Kumar Gauthaman¹ Padma Subramanyam¹ Shanmuga Sundaram Palaniswamy¹

¹ Department of Nuclear Medicine & Molecular imaging, Amrita Institute of Medical Sciences, Kochi, Kerala, India

World J Nuclear Med 2022;21:338-341.

Address for correspondence Shyma Basheer, MD, Department of Nuclear Medicine & Molecular Imaging, Amrita Institute of Medical Sciences (AIMS), Ponekkara, Kochi, 682 041, Kerala, India (e-mail: shyma.b@gmail.com).

Abstract

Keywords

- primary hepatic
 lymphoma
- ► PET-MR
- diffuse large B cell lymphoma

Primary hepatic lymphomas are rare hepatic malignancies which are often misdiagnosed preoperatively. Early accurate diagnosis is essential as the patients can be treated successfully with chemotherapy, eluding the need for surgery. We present a case of primary hepatic lymphoma which mimicked as focal nodular hyperplasia with normal biochemical tumor markers, and ¹⁸F-fluorodeoxyglucose (¹⁸F-FDG) whole-body positron emission magnetic resonance showed intense FDG uptake in the large hepatic lesion. The patient subsequently underwent right hepatectomy, and histopathology revealed diffuse large B cell lymphoma.

Key Messages

- Primary hepatic lymphoma can present with imaging features of focal nodular hyperplasia.
- Intense fluorodeoxyglucose uptake in solitary hepatic lesion should always raise a suspicion of malignancy and biopsy may be considered.

Introduction

The isolated primary hepatic lymphoma (PHL) constitutes approximately 0.0016% of all non-Hodgkin's lymphoma (NHL) patients. As PHL is rare, the patients may present with nonspecific symptoms and imaging characteristics. Early diagnosis can be challenging and sometimes established only postoperatively. We present a case of surgically resected primary hepatic diffuse large B cell lymphoma, in which computed tomography (CT) showed suspicion of focal nodular hyperplasia (FNH) and magnetic resonance imaging (MRI) showed suspicion for fibrolamellar carcinoma or FNH

published online September 9, 2022 DOI https://doi.org/ 10.1055/s-0042-1750398. ISSN 1450-1147. and positron emission tomography (PET) showed intense fluorodeoxyglucose (FDG) uptake in the lesion by which suspicion of possible primary hepatic malignancy was raised. Patient subsequently underwent right hepatectomy and histopathology showed diffuse large B cell lymphoma.

Case History

A 64-year-old female presented with complaints of epigastric discomfort, difficulty in eating, loss of appetite, and loss of weight. CT showed a large space-occupying lesion in the right lobe of the liver with central scar which was suspicious for FNH. AFP (alpha-fetoprotein), CEA (carcinoembryonic antigen), and CA 19–9 were within normal levels.

A dose of 6 mCi of ¹⁸F-FDG was administered intravenously to the patient in the fasting state. After 35 minutes of injection, on a 3 Tesla Siemens Biograph mMR, image acquisition was started. ¹⁸F-FDG whole-body PET-magnetic resonance (MR) showed intense FDG uptake in large lobulated tumor in the right lobe of the liver. Central portions of the

© 2022. World Association of Radiopharmaceutical and Molecular Therapy (WARMTH). All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/licenses/by-nc-nd/4.0/)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India



Fig. 1 ¹⁸F-FDG PET maximal intensity projection image showing large focus of increased abnormal uptake of FDG in the right lobe of liver. Physiological tracer uptake is seen in the brain, kidneys, and urinary bladder. Axial section of the liver on MR and fused PET-MR showing that bright central portions of the tumor are bright on T2-weighted images, consistent with the necrosis/scar with solid lesion in the periphery. MR, magnetic resonance; PET, positron emission tomography.



Fig. 2 Right hepatic mass in diffusion-weighed imaging, ADC map, postcontrast imaging, and T1-weighted volumetric interpolated breath-hold examination (VIBE). Postcontrast MR imaging shows washout of the solid components of the tumor in delayed phase and delayed enhancement of the central T2 bright scar tissue. MR, magnetic resonance.



Fig. 3 Micrograph of immunohistochemical staining of histopathology of right hepatectomy specimen showing positive staining for CD 10, bcl2, and bcl6 (IHC, \times 400).

tumor are bright on T2-weighted images and consistent with the necrosis/scar. Solid components in the periphery have a lobulated well-defined margin. Postcontrast imaging shows washout of the solid components of the tumor in the delayed phase and delayed enhancement of the central T2 bright scar tissue. Diffusion restriction was noted in the diffusionweighed imaging and apparent diffusion coefficient (ADC) mapping. Fused imaging demonstrated intense FDG uptake in the peripheral solid components of the lesion. SUVmax of the lesion was 24. MRI features were suspicious for fibrolamellar carcinoma or FNH. As the lesion showed intense FDG uptake in the isolated liver lesion with no other FDG-avid lesions on whole-body PET-MR imaging, the possibility of primary hepatic malignancy was considered. Thus, the patient underwent right hepatectomy. The histopathology of the specimen revealed dyscohesive cellular neoplasm arranged in lobules, nests, cords, and trabeculae admixed with lymphocytes, eosinophils, and foamy macrophages with ingested debris. The neoplastic cells are positive for CD 20, Pax 5, CD 10, bcl 6, and bcl 2. *K*_i index greater than 90%. The diagnosis of diffuse large B cell lymphoma was confirmed. The patient was subsequently started on rituximab, cyclophosphamide, doxorubicin, vincristine, and prednisone (R-CHOP) regimen of chemotherapy.

Discussion

PHL is defined as lymphoma confined to the liver without any involvement of other organs, or leukemic changes in the peripheral blood for at least 6 months after diagnosis. PHL represents 0.4% of extranodal lymphomas and 0.016% of NHLs.¹ Although the pathogenesis of PHL is not well understood, it is usually seen in immunocompromised patients like organ-transplant recipients and individuals with acquired immune deficiency syndrome receiving immunosuppressive therapy. Recently, a strong relation between hepatitis C infection and PHL has also been identified, proposed cause being stimulation of B lymphocytes and chronic polyclonal proliferation, resulting in hepatic lymphoma.²

PHL shows variable clinical presentation from incidental discovery in asymptomatic patients to nonspecific symptoms such as fatigue, night sweats, weight loss, evening rise of temperature, jaundice, abdominal pain, and hepatomegaly. Laboratory tests are also not helpful in diagnosis, with tumor markers like CEA, AFP, and CA 19.9 usually within normal ranges.^{3,4}

PHL may be Hodgkin's or non-Hodgkin's, with the latter being more common. B cell type is more common than the T cell type. Microscopically, PHL may show diffuse or nodular pattern of infiltration, both resulting in destruction of liver parenchyma. Among B cell types, diffuse large B cell type is the commonest followed by chronic lymphocytic lymphoma, Burkitt's lymphoma, and mantle cell lymphoma. Similar to peripheral T cell NHL, NOS is common among T cell type, followed by anaplastic large cell lymphoma and hepatosplenic gamma delta lymphomas.^{5,6}

PHL can present as a solitary lesion, multiple lesions within the liver, or as diffuse hepatic infiltration. Radiological appearance of PHL is nonspecific and can resemble a hypovascular metastasis.⁷ Fungal and tubercular infections and inflammatory pathology like sarcoidosis may also show similar imaging appearances. PHLs are usually hypoechoic in ultrasonogram, hypoattenuating with minimal or no enhancement in CT, hypodense in T1-weighted and variable in T2-weighted sequences with similar pattern of enhancement in MRI.⁸ As a result, PHLs are rarely diagnosed preoperatively and commonly misdiagnosed as other malignant or benign lesions. In this case, the MRI appearance was highly suggestive of FNH or fibrolamellar variant of hepatoma. However, FDG avidity as well as absence of elevated tumor markers should have served as an alert, to consider an alternate diagnosis. Biopsy should be considered in such atypical cases for accurate diagnosis to plan appropriate treatment.

Surgical resection, chemotherapy, and radiotherapy alone or in combination are the various treatment modalities. The CHOP regimen is the standard treatment given for the patients with diffuse large B cell lymphoma. Rituximab targeting the CD20 antigenic marker can be added, which results in an improved response rate and prolonged overall survival. Liver resection followed by adjuvant chemotherapy and/or radiation has also been shown to have favorable short-term and mid-term outcomes with a prolonged survival rate.⁹

The factors which determine the treatment outcome and prognosis are advanced age, poor performance status, bulky disease, unfavorable histology, elevated lactate dehydrogenase, comorbidities, and immunosuppressed state. Since PHL is chemosensitive, there is a distinct role of chemotherapy as an adjuvant to surgery or as a primary treatment modality.

In conclusion, PHL is extremely rare with nonspecific clinical presentation and laboratory and imaging findings. This case demonstrates that this diagnosis should be considered in any patient with atypical presentation like FDG-avid liver mass with normal tumor markers. Definite diagnosis and timely therapeutic intervention can produce better outcome in these patients than other liver tumors owing to their chemosensitivity.

Funding None.

Conflict of Interest None declared.

References

1 Dantas E, Santos J, Coelho M, et al. Primary hepatic lymphoma in a patient with cirrhosis: a case report. J Med Case Reports 2020;14 (01):168

- 2 Nasr Ben Ammar C, Chaari N, Kochbati L, Besbes M, Maalej M. Primary non-Hodgkin lymphoma of the liver: case report and review of the literature [in French]. Cancer Radiother 2006;10 (08):595–601
- ³ Park YK, Choi JE, Jung WY, Song SK, Lee JI, Chung CW. Mucosaassociated lymphoid tissue (MALT) lymphoma as an unusual cause of malignant hilar biliary stricture: a case report with literature review. World J Surg Oncol 2016;14(01):167
- 4 Ugurluer G, Miller RC, Li Y, et al. Primary hepatic lymphoma: a retrospective, multicenter rare cancer network study. Rare Tumors 2016;8(03):6502
- ⁵ Bauduer F, Marty F, Gemain MC, Dulubac E, Bordahandy R. Primary non-Hodgkin's lymphoma of the liver in a patient with hepatitis B, C, HIV infections. Am J Hematol 1997;54(03):265
- 6 Stancu M, Jones D, Vega F, Medeiros LJ. Peripheral T-cell lymphoma arising in the liver. Am J Clin Pathol 2002;118(04):574–581
- 7 Levy AD. Malignant liver tumors. Clin Liver Dis 2002;6(01): 147-164
- 8 Padhan RK, Das P, , Shalimar. Primary hepatic lymphoma. Trop Gastroenterol 2015;36(01):14–20
- 9 Yaka M, Chehab F, Allaoui M, Ait Ali A, Zentar A. Postsurgical diagnosis of an unusual case of primary hepatic LYMPHOMA presenting as liver abscess with an uncommon complication: a hepatogastric fistula. Case Rep Hematol 2021;2021:6647558