

Y. BRÄGGER, O. GREEN, B. N. BHAWAL*, B. MORANDI* (ETH ZÜRICH, SWITZERLAND, UNIVERSITY OF EDINBURGH, UK)

Late-Stage Molecular Editing Enabled by Ketone Chain-Walking Isomerization

J. Am. Chem. Soc. **2023**, *145*, 19496–19502, DOI: 10.1021/jacs.3c05680.

Ketone Chain-Walking Isomerization by Organocatalysis

Category

Organo- and Biocatalysis

Key words

isomerization

organocatalysis


chain-walking

sulfur


ketone

reversible transition


Synfact
of the
Month



Selected examples



Proposed mechanism



Significance: The Morandi group developed a novel and simple method for isomerizing cyclic ketones. To perform this reversible transition, widely available and affordable pyrrolidine and elemental sulfur are used as catalysts. The isomerization of several steroids was demonstrated to access new steroids by using this remarkable ketone chain-walking technique.

Comment: Modifying the placement of functional groups is arguably difficult since it necessitates reversible cleavage and the transposition of strong bonds. Inspired by the Willgerodt–Kindler reaction, the authors succeeded in brilliantly designing a method for the ketone chain-walking isomerization of cyclic ketones enabled by an organocatalyst in combination with elemental sulfur. This low-cost, ecologically friendly approach has the potential to be widely used in the future.

SYNFACTS Contributors: Benjamin List, Fuxing Shi
Synfacts 2023, 19(11), 1137 Published online: 17.10.2023
DOI: 10.1055/s-0042-1752030; Reg-No.: B09623SF

© 2023, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

1137