


R. L. DOROW*, D. E. GINGRICH (DUPONT PHARMACEUTICALS COMPANY, WILMINGTON, USA)


A Novel, One-Pot Preparation of *N*-Methyl- α -Amino Acid Dipeptides from Oxazolidinones and Amino Acids

Tetrahedron Lett. **1999**, *40*, 467–470, DOI: 10.1016/S0040-4039(98)02426-5.

One-Pot Synthesis of *N*-Methyl Dipeptides from Amino Acid-Derived Oxazolidinones



Selected examples:



Significance: *N*-Methyl peptides play a crucial role in peptide drug discovery. This *N*-methyl peptide is especially useful for the improvement of biological properties such as target affinity, proteolytic stability and membrane permeability as well as the alteration of the conformational rigidity and hydrophobicity of the peptides. In 1999, R. L. Dorow and co-workers developed a one-pot synthesis of *N*-methyl- α -amino acid dipeptides from oxazolidinones and amino acids.

Comment: A series of amino-acid-derived oxazolidinones reacted smoothly with the amino acid esters to produce *N*-hydroxymethyl dipeptides, which were treated with TFAA and Et₃SiH to produce the corresponding *N*-methyl dipeptides in good yields. This one-pot protocol is a practically simple and highly efficient method to produce various *N*-methyl peptides.

SYNFACTS Contributors: Hisashi Yamamoto, Isai Ramakrishna
Synfacts 2023, 19(03), 0319 Published online: 13.02.2023
DOI: 10.1055/s-0042-1753292; Reg-No.: H01823SF

© 2023, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Peptide Chemistry

Key words

oxazolidinones

N-methyl dipeptides

amino acid esters

Synfact
Classic