
E. MIGUEL-CASAÑ, M. D. DARAWSHEH, V. FARIÑA-TORRES, I. J. VITÓRICA-YREZÁBAL, E. ANDRES-GARCIA, M. FAÑANÁS-MASTRAL*, G. MÍNGEZ ESPALLARGAS* (UNIVERSIDAD DE VALENCIA, PATERNA AND UNIVERSIDADE DE SANTIAGO DE COMPOSTELA, SPAIN) Heterometallic Palladium—Iron Metal—Organic Framework as a Highly Active Catalyst for Cross-Coupling Reactions Chem. Sci. 2023, 14, 179–185, DOI: 10.1039/d2sc05192c.

Cross-coupling of Cinnamyl Bromides and Alkyl Boronates Catalyzed by a Pd-Fe MOF



Significance: A heterometallic palladium–iron metal-organic framework (**Pd-Fe-MOF**) was prepared according to equation 1. **Pd-Fe-MOF** catalyzed the allylic alkylation of cinnamyl bromides with alkylboronic pinacol esters in the presence of Cs_2CO_3 and KF to afford the corresponding coupling products in \leq 76% yield (eq. 2).

Comment: Pd-Fe-MOF was characterized by means of PXRD, single-crystal XRD, XPS, TGA, FT-IR, SEM, EDS, ICP-MS and N_2 , CO_2 , CH_4 gas adsorption isotherm analyses. In the coupling reaction between cinnamyl bromide and phenylboronic acid pinacol ester, Pd-Fe-MOF was recovered and reused three times without significant loss of its catalytic activity.

SYNFACTS Contributors: Yasuhiro Uozumi, Shusuke Hattori Synfacts 2023, 19(03), 0279 Published online: 13.02.2023 DOI: 10.1055/s-0042-1753309; Reg-No.: Y03123SF

Polymer-Supported Synthesis

Key words

heterometal

palladium

iror

metal-organic framework

