
Endoscopic mucosa-submucosal clip closure method

It is desirable to perform a prophylactic closure using endoclips in patients who are at high risk of delayed bleeding or perforation after endoscopic resection [1]. A mucosal defect after endoscopic mucosal resection can be closed with sequential endoclips as if zipping up the mucosal edges. However, it is difficult to close a large mucosal defect after endoscopic submucosal dissection (ESD) using only conventional endoclips. Several methods have been reported, such as the "slip knot clip suturing method" [2] and the "endoscopic hand-suturing" technique [3]. However, these endoscopic closure methods can be difficult, and require a string or special device, which is not always available and takes time to apply. This report describes a simpler closure method using conventional endoclips.

In the "mucosa-submucosa clip closure method", endoclips (EZ Clip, HX-610-090 L long-type; Olympus, Tokyo, Japan) are placed at the edge of the mucosal defect after colorectal ESD. Each arm of the endoclip grips the mucosa and submucosa, respectively. The direction in which the endoclip grips is parallel to the short axis of the defect. Several endoclips are applied in this way, and the mucosal defect is significantly reduced in size. Then, additional endoclips can be applied to both sides of the mucosal defect. Several endoclips are required to achieve complete closure (Video 1).

This method is simple, and greatly facilitates complete closure of a large mucosal defect simply by applying conventional endoclips.

Endoscopy_UCTN_Code_TTT_1AQ_2AC





▶ Video 1 Mucosa–submucosa clip closure method. a A mucosal defect after colonic endoscopic submucosal dissection. b Endoclips are placed at the edge of the mucosal defect. Each arm of the endoclip hooks mucosa and submucosa, respectively. The direction in which the endoclip grips is parallel to the short axis of the defect. c Several endoclips are applied to the mucosal edge in the same way, bringing the mucosal edges close to each other and enabling application of clips to both sides of the mucosa. d Additional endoclips are placed to achieve complete closure.

Competing interests

None

The Authors

Shigeo Banno¹, Toshihiro Nishizawa¹.².³, Satoshi Kinoshita¹.².³, Hideki Mori¹.², Toshio Uraoka¹.³

- Department of Gastroenterology, National Hospital Organization Tokyo Medical Center, Tokyo, Japan
- 2 Department of Gastroenterology and Hepatology, Keio University School of Medicine, Tokyo, Japan
- 3 Division of Research and Development for Minimally Invasive Treatment, Cancer Center, Keio University School of Medicine, Tokyo, Japan

Corresponding author

Toshio Uraoka, MD, PhD

Department of Gastroenterology, National Hospital Organization Tokyo Medical Center, 2-5-1 Higashigaoka, Meguro-ku, Tokyo 152-8902, Japan

Fax: +81-3-53633967 toshi_urao@yahoo.co.jp

References

- [1] Liaquat H, Rohn E, Rex DK. Prophylactic clip closure reduced the risk of delayed postpolypectomy hemorrhage: experience in 277 clipped large sessile or flat colorectal lesions and 247 control lesions. Gastrointest Endosc 2013; 77: 401 – 407
- [2] Nishizawa T, Ochiai Y, Uraoka T et al. Endoscopic slip-knot clip suturing method: prospective pilot study (with video). Gastrointest Endosc 2017; 85: 433 – 437

[3] Goto O, Sasaki M, Akimoto T et al. Endoscopic hand-suturing for defect closure after gastric endoscopic submucosal dissection: a pilot study in animals and in humans. Endoscopy 2017; 49: 792 – 797

Bibliography

DOI https://doi.org/10.1055/s-0043-119349 Published online: 9.10.2017 Endoscopy 2017; 49: E307–E308 © Georg Thieme Verlag KG Stuttgart · New York ISSN 0013-726X

ENDOSCOPY E-VIDEOS https://eref.thieme.de/e-videos



Endoscopy E-Videos is a free access online section, reporting on interesting cases and new

techniques in gastroenterological endoscopy. All papers include a high quality video and all contributions are freely accessible online.

This section has its own submission website at https://mc.manuscriptcentral.com/e-videos