


J. WENG*, S.-C. WANG, X. ZHOU, Y.-X. LI, C.-Y. ZHANG, Z.-Y. ZHANG, Y.-S. XIONG, G. LU, J. DONG (SUN YAT-SEN UNIVERSITY, GUANGZHOU, P. R. OF CHINA)
Enabling Modular Click Chemistry Library through Sequential Ligations of Carboxylic Acids and Amines
Angew. Chem. Int. Ed. **2024**, e202410699 DOI: 10.1002/anie.202410699.

High-Throughput Access to Libraries of *N*-Acylsulfamide Derivatives, a Bioisostere of Carboxylic Acids



Significance: *N*-Acylsulfamides have been established as bioisosteres of carboxylic acids, notably in the HCV NS5B polymerase inhibitor beclabuvir which was approved in Japan, the Nav1.7 inhibitor GDC-0276 and the herbicide saflufenacil. The process developed in this article is mild and high yielding, amenable to the generation of large libraries of *N*-acylsulfamide derivatives with products typically isolated by a simple filtration after acidification of the reaction mixture. The process was demonstrated with a range of carboxylic acid derivatives that explored only relatively benign substituents (for example, there was an absence of alcohols or amides in the acids), although the amines contained a broader range of functionality that included acids, tertiary amines, and alkynes. Bis-carboxylic acids could be fully or partially converted by controlling the ratio of reagents. The process has potential for late-stage functionalization of carboxylic acids, the scope of which will depend on the functionality inherent to a molecule.

Comment: Exposure of a carboxylic acid to the highly electrophilic but readily available fluorosulfonyl isocyanate in CH₃CN, determined to be the crucial solvent, at room temperature afforded an *N*-fluorosulfonyl amide in yields typically >90%. The scope of the process extended to aromatic, heteroaromatic and aliphatic carboxylic acids with limited polar functionality, although NH-sulfonamide and imide functionality were compatible, and the reaction was tolerant of steric encumbrance. Electron-deficient acids required heating to 50–60 °C in order to complete the reaction. Reaction of the *N*-fluorosulfonyl amides with aromatic and aliphatic amines occurred in H₂O with Na₂CO₃ as the base heated at 80 °C for 12 hours to afford *N*-acylsulfamides in good to excellent yields, with the products typically isolated by acidification and simple filtration to afford the product. Execution of a 13x24 library in 96 well plates generated 312 products, with 281 proceeding with >90% conversion and 97% proceeding with >80% conversion.

SYNFACTS Contributors: Antonia F. Stepan (Roche), Nicholas A. Meanwell (Baruch S. Blumberg Institute, U. Michigan, Rutgers U.)
Synfacts 2024, 20(09), 0981 Published online: 16.08.2024
DOI: 10.1055/s-0043-1775033; Reg-No.: A04424SF

© 2024, Thieme. All rights reserved.
Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Category

Innovative Drug
Discovery and
Development

Key words

bioisostere design

carboxylic acid
bioisosteres

N-acylsulfamides

library generation

Synfact
of the
Month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.