Mir Bilal<sup>1</sup> Parveen Shaheena<sup>2</sup> Ashish Chauhan<sup>10</sup> Mushfiq Syed<sup>20</sup> Majeed Tahir<sup>1</sup> Nadeem Mir<sup>3</sup> Rameez Najar<sup>3</sup> Kalpana Acharya<sup>2</sup>

Address for correspondence Mushfig Syed, MD, DM, Department of

Gastroenterology, Sher-i-Kashmir Institute of Medical Sciences,

Srinagar, Jammu and Kashmir, 190011, India

(e-mail: syedmushfiq37@gmail.com).

 $\bigcirc$   $\bigcirc$   $\bigcirc$   $\bigcirc$   $\bigcirc$   $\bigcirc$ 

<sup>1</sup>Department of Gastroenterology, Indira Gandhi Medical College and Hospital, Shimla, Himachal Pradesh, India

<sup>2</sup>Department of Gastroenterology, Sher-i-Kashmir Institute of Medical Sciences, Srinagar, Jammu and Kashmir, India

| Gastrointest Infect 2023;13:68-73.

Abstract	<b>Background</b> Cholangitis is one of the commonest emergencies encountered by
	(PCT) is an important biomarker of cholangitis with a potential to guide early
	therapeutic decision.
	Materials and Methods We did a retrospective analysis of prospectively maintained
	data of patients admitted in the hospital from March 2018 to September 2020 with a
	diagnosis of acute cholangitis based on Tokyo 18 guidelines (TG-18). All the demo-
	graphic parameters, biochemical and hematological parameters, and the result were recorded.
	Results One-hundred two patients were admitted to our hospital with a mean age of
	51.07 $\pm$ 7.99 years, among which 58.5% (59/102) were females with most common
	etiology being choledocholithiasis (64%). Fever, jaundice, and abdominal pain were
	seen in 78% (80/102), 73.3% (74/102), and 76.9% (78/102) of patients, respectively.
	Organ failure was seen in 29.3% (29/102) of patients with most common organ failure
	being acute kidney injury. Mild, moderate, and severe cholangitis as per TG-13 was seen
	in 43.9, 26.8, and 29.3% of patients, respectively. Elevated PCT levels were associated
	with severe disease (76.7 vs. 26.5%, $p < 0.05$ ). Forty-three percent (44/102) patients
	were managed with delayed biliary drainage, while the majority of remaining specifi-
Keywords	cally with raised PCT levels needed early endoscopic intervention.
<ul> <li>cholangitis</li> </ul>	<b>Conclusion</b> In this retrospective analysis, we showed that most common etiology of
<ul> <li>choledocholithiasis</li> </ul>	cholangitis in was choledocholithiasis. Raised serum PCT levels were significantly
<ul> <li>procalcitonin</li> </ul>	associated with increased severity of cholangitis.

procalcitonin

received May 22, 2023 first decision June 13, 2023 accepted September 14, 2023 article published online November 27, 2023

DOI https://doi.org/ 10.1055/s-0043-1776392. ISSN 2277-5862.

© 2023. Gastroinstestinal Infection Society of India. All rights reserved.

This is an open access article published by Thieme under the terms of the Creative Commons Attribution-NonDerivative-NonCommercial-License, permitting copying and reproduction so long as the original work is given appropriate credit. Contents may not be used for commercial purposes, or adapted, remixed, transformed or built upon. (https://creativecommons.org/ licenses/by-nc-nd/4.0/)

Thieme Medical and Scientific Publishers Pvt. Ltd., A-12, 2nd Floor, Sector 2, Noida-201301 UP, India

<sup>&</sup>lt;sup>3</sup>Department of Gastroenterology, G.M.C, Srinagar, Jammu and Kashmir, India

# Introduction

Acute cholangitis (AC) is one of the commonest, yet potentially treatable emergency encountered by gastroenterologists worldwide. The process of biliary stasis that is most commonly secondary to mechanical obstruction like choledocholithiasis, benign strictures, malignancy, and stent blockade, and increased intraductal pressure is main pathogenic event that leads to AC.<sup>1</sup> The rise in intraductal pressure causes bacterial translocation into the systemic circulation known as cholangial-venous reflux that is the basic pathophysiology behind suppurative AC.<sup>2</sup> Moreover, the biliary stasis abolishes the continuous flushing activity of bile and the bacteriostatic effect of bile salts, and ultimately both lead to severe systemic inflammatory cascade and multiorgan dysfunction. This becomes potentially lethal when sepsis sets in, and prompt triage is mandatory to define timely management strategy and achieve favorable outcome.

The severity assessment is currently guided by Tokyo Grading; however, recent data have shown serum procalcitonin (PCT) level at admission to be a useful predictor of clinical severity.<sup>3</sup> PCT is a 116 amino acid protein and precursor of calcitonin. Under physiological conditions, calcitonin is produced predominantly by C cells of the thyroid gland, and serum PCT is almost undetectable. The normal values of serum PCT are less than 0.1 ng/dL and values more than 0.25ng/dL are suggestive of infective process.<sup>4</sup> In states of systemic bacterial infection, PCT is readily produced by various tissues of the body (e.g., lung, kidney, liver, adipose cell, and muscle) immediately after disease onset (usually within 6-12h), and its serum values rise to significantly higher levels.<sup>5–8</sup> There are noninfective causes of serum PCT elevation that include malignancies (medullary thyroid carcinoma, lung carcinoma, and lymphoma), drugs (steroids, rituximab), and chronic kidney disease (CKD).<sup>9,10</sup>

Given the importance of risk stratification in AC and appropriate management, serum PCT might be a helpful biomarker. We evaluated our database to see the correlation between serum PCT and grades of cholangitis across all etiologies and whether serum PCT at admission has any role in deciding treatment strategy.

#### Methods

We did a retrospective analysis of prospectively maintained data of patients admitted in a single tertiary care center from 2019 to September 2020 with a diagnosis of AC and severity grading as mild, moderate, and severe cholangitis as per Tokyo 18 guidelines (TG-18) guidelines. Clinical parameters, organ failures, and levels of PCT were measured at baseline. We measured serum PCT levels (measured by immunoluminometric assay) in patients and compared the values between different grades of severity of cholangitis. The upper limit normal of serum PCT was taken as 0.25 ng/dL. This study was approved by the institutional ethical committee, and was conducted according to the guidelines in the Helsinki Declaration. Written informed consent was obtained from all patients.

## Patients

All patients aged more than 18 years of age with diagnosis of AC as per TG-18 were included as cases. Both referred (from peripheral healthcare facility) and direct admission patients were included. Other inclusion criteria were; if no serum PCT was done at the time of admission, patients with a history of known malignancy, patients on steroids and patients of CKD. Patient demographic characteristics including age, sex, body mass index, comorbidities were obtained. Symptoms at presentation along with hematological and biochemical variables including inflammatory markers like PCT and total leukocyte count (TLC) were recorded. All patients received fluid therapy and parenteral antibiotics. Inotropic agents were administered in fluid nonresponsive cases. The assessment of initial medical treatment response and timing and method of biliary decompression was done as per the discretion of the treating physician.

#### Statistics

All analysis was done using Statistical Package for Social Sciences (SPSS) version 21.0. Continuous variables were summarized as mean and standard deviation. Categorical variables were summarized as frequency and percentage. Chi-squared test was used to analyze the relationship between two categorical variables. Two-sided *p*-valves were reported and a *p*-valve less than 0.05 was considered statistically significant.

### Results

**Patient characteristics** The mean age of our study population (n = 102) was  $51.1 \pm 7.9$  years, among which 58.5% (59/102) patients were females. The baseline characteristics of study cohort is given in **-Table 1**. The four most common presenting symptoms were abdominal pain, jaundice, vomiting, and fever among which fever, jaundice, and abdominal pain were seen in 76.9 (78/102), 73.3 (74/102), and 78% (80/102) of patients, respectively. Charcot's triad including pain, jaundice, and fever was present in 45% (46/102) of patients, whereas two symptoms were present in 90% (92/102) of patients. The most common organ failure was renal failure, present in 60% of severe cholangitis patients followed by respiratory and hemodynamic failures

**Etiology** The spectrum of etiology included choledocholithiasis, post-cholecystectomy bile duct injury, hydatid cyst, common bile duct stent block, malignant obstruction, and others (**-Table 1**). The most common etiology in this study was choledocholithiasis (64%). Mild, moderate, and severe cholangitis was seen in 43.9, 26.8, and 29.3% of patients, respectively.

**Blood cultures** Blood culture was positive in 25% of patients among which 80% were having severe (grade-III) cholangitis. The various organisms grown on culture are shown in results (**~Fig. 1**).

**Table 1** Baseline characteristics of acute cholangitis study cohort (n = 102)

Variable		$Mean\pmSD$
Age (years)		$51.07 \pm 7.99$
Weight (kg)		$68\pm12$
Females		58%
BMI (kg/m <sup>2</sup> )		$26\pm2.8$
Comorbidity (%)	Metabolic syndrome	3
	Diabetes	5
	Hypertension	9
Presenting symptom (%)	Pain	90
	Jaundice	70
	Fever	82
	Vomiting	50
	Charcot's triad	45
Etiology (%)	Choledocholithiasis	64
	Post chole BDI	15
	Hydatid cyst	10
	CBD stent	5
	Benign biliary stricture	3
	Malignant obstruction	4
	Others	1
Blood pressure (%)	Normal	82
	Hypotension-fluid responsive	15
	Shock requiring Pressors	5
Mentation (%)	Normal	97
	Altered	3
SPO2 (%age)	Normal (>95)	75
	Mild hypoxemia (90–95)	20
	Severe hypoxemia (<90)—ARDS	7
TLC (x1000/µL)		8.7 (5.4–13.2)
Platelet (x1000/mL)		120 (90–180)
CRP (mg/dL)		4.9 (0.2–11.8)
PCT (ng/mL)		9.8 (0.4–38)
AST (U/L)		180(50–450)
ALT (U/L)		200(65–459)
ALP (U/L)		365(190-850)
Bilirubin (mg/dL)		3.5 (0.0-8.9)
INR		1.3 (0.98–1.9)
Creatinine (mg/dL)		1.57(0.78-3.2)
BUN (mg/dL)		19 (10–28)

Abbreviations: ALP, alkaline phosphatase; ALT, alanine transaminase; ARDS, acute respiratory distress syndrome; AST, aspartate transaminase; BMI, body mass index; BUN, blood urea nitrogen; CRP, C-reactive protein; INR, international normalized ratio; PCT, procalcitonin; SD, standard deviation; TLC, total leukocyte count.

## Procalcitonin and Other Inflammatory Markers for Assessment of Severity

The sensitivity and specificity of serum PCT for predicting grade of cholangitis and septic shock are plotted under

receiver operating-characteristic with area under the curve of 0.65 and 0.97, respectively (**Figs. 2** and **3**). Among TLC, C-reactive protein (CRP), and PCT, the statistically significant prediction for severe cholangitis was only seen with PCT,



Fig. 1 Various microorganisms grown on blood cultures.



**Fig. 2** Receiver operating-characteristic (ROC) curve for procalcitonin and severity of cholangitis.

with p = 0.035, p = 0.92, and p = 0.052 for PCT, TLC, and CRP, respectively (**>Table 2**).

**PCT and organ failures** Values of PCT vary significantly between the groups when patients were compared in terms of severity of cholangitis, cholangitis with or without organ failure, and patients with or without positive blood culture.

**Correlation of PCT with severity of disease, presence or absence of organ failures and blood culture** e PCT values were significantly variable in patients in terms of severity of cholangitis (mild–moderate: 0.8 [0.2–2.5] vs. severe cholangitis: 6.9 [0.6–38]) organ failure (organ failure present: 6.9 Q10 [0.6–38] vs. organ failure absent: 0.8 [0.2–2.5]) and also PCT values also differ from blood culture positive: 5.8 [0.9–38]



**Fig. 3** Receiver operating-characteristic (ROC) curve for procalcitonin and shock.

vs. negative: 1.1 [0.4–10.5]. The mean serum PCT levels were 2.9  $\pm$  1.2. Elevated PCT levels were significantly associated with severe disease compared to mild-to-moderate cholangitis (76.7 vs. 26.5%, p < 0.05), and majority of these (90%) needed early (<24 hours) endoscopic biliary drainage. Overall mortality in our studied group was 3.3%.

## Discussion

AC is an emergency where endoscopic intervention in addition to aggressive medical management can be lifesaving, and the delay in same especially in severe cases results in increased mortality. According to TG-18 guidelines, severe cholangitis is characterized by sepsis-associated life-threatening organ failure, while mild-to-moderate cholangitis is not associated with organ failures. The main aim of this study was to obtain a serum biomarker, which is easily done in laboratory, and correlate its values to predict severity of disease, organ failure, response to treatment, and mortality.

There are varied etiologies of cholangitis in our study; most common etiology of AC is choledocholithiasis. The most common organ failure observed is acute kidney injury. AC results from biliary stasis due to obstruction to bile flow and subsequent infection of static bile. While the fluid therapy and antibiotics are indicated in all patients, biliary drainage becomes cornerstone treatment in AC and as per TG-18 guidelines. In patients with mild-to-moderate cholangitis, biliary drainage should be performed within 24 to 48 hours.

 Table 2
 Levels of different serum markers across all grades of acute cholangitis based on TG-13

Marker	Grade 1 cholangitis (n = 43.9%)	Grade II cholangitis (n = 26.8%)	Grade III cholangitis (n = 29.3%)	p-Value
TLC (x1000/µL)	7.2 (4.3–12.2)	6.1(5.4–12.8)	8.5(6.5–14.2)	0.092
CRP (mg/dL)	0.8 (0.5–1.2)	3.9 (0.9–5.9)	4.6 (0.8–11.4)	0.052
PCT (ng/mL)	0.4 (0.2–1.3)	1.2 (0.5–2.5)	6.9 (0.6–38)	0.035

Abbreviations: CRP, Greactive protein; PCT, procalcitonin; TG-13, Tokyo Grading-13; TLC, total leukocyte count.

Patients with mild-to-moderate cholangitis that fail to respond to conservative management for 24 hours and patients with severe (suppurative) cholangitis require urgent (within 24 hours) biliary decompression. In our study, 58 (57%) patients needed emergency intervention, while 44 (43%) were subjected to delayed intervention.

Shinya et al<sup>11</sup> have concluded that many cases of AC with positive hemocultures or purulent bile were classified as either mild or moderate cholangitis on the basis of TG-13 grading despite the fact that these cases represent systemic septicemia and need early biliary drainage for source control to prevent imminent organ failure. This highlights the importance of serum markers to reflect impending organ failure and guide to early intervention. Serum PCT has been shown to correlate well with systemic septicemia and severity of cholangitis at the earliest and a useful serum marker to prevent delayed intervention.<sup>6</sup>

This study shows PCT levels at admission were significantly raised in severe AC irrespective of etiology, and the serum level was proportional to the increasing severity of AC. This finding translates into the ability of PCT to predict severe cases of AC at the earliest and be useful clinical guide to decide management. In this study, all patients with grade III with hemodynamic failure had higher elevation of serum PCT with none of them had PCT values below 3.5 ng/dL. This probably reflects the cutoff level where AC starts worsening and degenerating into septic shock. The results were similar to study conducted by Lee et al, where he recorded a cutoff 3.7ng/dL for deterioration into shock.<sup>12</sup> Most previous studies have reported significantly higher levels of serum procalcitonin in patients assessed as having severe AC based on the TG severity grade.<sup>12–17</sup>

We studied the microorganisms found in the bile and their association with grade of cholangitis. The majority of patients with positive blood culture (~95%) had hemodynamic compromise, and all patients had hypotension.<sup>11</sup> The low blood culture yield is probably related to containment of infection within biliary system in most patients and use of prior antibiotics before hospitalization.

Although PCT seems to be a marker for diagnosis, severity assessment, treatment response, it is important to note that PCT is not a holy grain marker for cholangitis, and its interpretation should be done in conjunction with other clinical findings and test results.

Despite small sample numbers, we showed that serum PCT levels are useful at admission when assessing the severity of AC and could prove potential guide to early intervention as backed up by some Indian studies also.<sup>18,19</sup>

Ethical Statement Not applicable.

Authors' Contribution All authors contributed equally to the article.

Data Availability Statement

There is no data associated with this work.

Funding None.

Conflict of Interest None declared.

Acknowledgments None.

#### References

- Kimura Y, Takada T, Kawarada Y, et al. Definitions, pathophysiology, and epidemiology of acute cholangitis and cholecystitis: Tokyo Guidelines. J Hepatobiliary Pancreat Surg 2007;14(01): 15–26
- 2 Huang T, Bass JA, Williams RD. The significance of biliary pressure in cholangitis. Arch Surg 1969;98(05):629–632
- 3 Kiriyama S, Takada T, Strasberg SM, et al; Tokyo Guidelines Revision Committee. TG13 guidelines for diagnosis and severity grading of acute cholangitis (with videos). J Hepatobiliary Pancreat Sci 2013;20(01):24–34
- 4 Covington EW, Roberts MZ, Dong J. Procalcitonin monitoring as a guide for antimicrobial therapy: a review of current literature. Pharmacotherapy 2018;38(05):569–581
- 5 Self WH, Grijalva CG, Williams DJ, et al. Procalcitonin as an early marker of the need for invasive respiratory or vasopressor support in adults with community-acquired pneumonia. Chest 2016; 150(04):819–828
- 6 Assicot M, Gendrel D, Carsin H, Raymond J, Guilbaud J, Bohuon C. High serum procalcitonin concentrations in patients with sepsis and infection. Lancet 1993;341(8844):515–518
- 7 Becker KL, Nylén ES, White JC, Müller B, Snider RH Jr. Clinical review 167: Procalcitonin and the calcitonin gene family of peptides in inflammation, infection, and sepsis: a journey from calcitonin back to its precursors. J Clin Endocrinol Metab 2004;89 (04):1512–1525
- 8 Becker KL, Snider R, Nylen ES. Procalcitonin assay in systemic inflammation, infection, and sepsis: clinical utility and limitations. Crit Care Med 2008;36(03):941–952
- 9 Jensen JU, Heslet L, Jensen TH, Espersen K, Steffensen P, Tvede M. Procalcitonin increase in early identification of critically ill patients at high risk of mortality. Crit Care Med 2006;34(10): 2596–2602
- 10 Simon L, Gauvin F, Amre DK, Saint-Louis P, Lacroix J. Serum procalcitonin and C-reactive protein levels as markers of bacterial infection: a systematic review and meta-analysis. Clin Infect Dis 2004;39(02):206–217
- 11 Shinya S, Sasaki T, Yamashita Y, et al. Procalcitonin as a useful biomarker for determining the need to perform emergency biliary drainage in cases of acute cholangitis. J Hepatobiliary Pancreat Sci 2014;21(10):777–785
- 12 Lee YS, Cho KB, Park KS, Lee JY, Lee YJ. Procalcitonin as a decisionsupporting marker of urgent biliary decompression in acute cholangitis. Dig Dis Sci 2018;63(09):2474–2479
- 13 Scheinpflug K, Schalk E, Grabert E, Achenbach HJ. Procalcitonin is not useful to discriminate between infectious and noninfectious CRP elevation in patients with non-small cell lung cancer. Infect Control Hosp Epidemiol 2015;36(09): 1117–1118
- 14 Grace E, Turner RM. Use of procalcitonin in patients with various degrees of chronic kidney disease including renal replacement therapy. Clin Infect Dis 2014;59(12):1761–1767
- 15 Yokoe M, Takada T, Strasberg SM, et al; Tokyo Guidelines Revision Committee. TG13 diagnostic criteria and severity grading of acute

cholecystitis (with videos). J Hepatobiliary Pancreat Sci 2013;20 (01):35–46

- 16 Umefune G, Kogure H, Hamada T, et al. Procalcitonin is a useful biomarker to predict severe acute cholangitis: a single-center prospective study. J Gastroenterol 2017;52(06):734–745
- 17 Hamano K, Noguchi O, Matsumoto Y, et al. Usefulness of procalcitonin for severity assessment in patients with acute cholangitis. Clin Lab 2013;59(1-2):177–183
- 18 Dhar J, Samanta J, Birda CL, et al. Dynamics of serum procalcitonin for predicting outcome in patients of infected pancreatic necrosis. (UEG Week 2021 Poster Presentations). United European Gastroenterol J 2021;9:740
- 19 Jain A, Jena A, Gautam V, Samanta J, Sharma V, Mandavdhare HS. Role of change in the levels of inflammatory markers post drainage in predicting outcome in acute cholangitis. Arq Gastroenterol 2022;59(02):212–218