

46 XX Ovotesticular Disorder of Sex Development with Gonadotropin-Releasing Hormone Receptor, Autosomal Recessive Heterozygous Missense Mutation and Autosomal Dominant Heterozygous Missense Mutation of the *PROKR2* Gene: A Case Report

Francesca Peranzoni¹ Roberto De Castro² Emilio Merlini³ Yen Le Nguyen⁴

¹ Department of Pediatric Surgery, Lausanne University Hospital, Lausanne, Switzerland

²Thien Nhan and Friends Association, Italy

³Department of Pediatric Surgery, Hospital of Alexandria, Italy

- ⁴Department of Pediatric Urology, Vietnam National Hospital of
- Pediatric 2, Ho Chi Minh City, Vietnam

Glob Med Genet 2024;11:220-224.

Abstract

True hermaphroditism is a disorder of sex development (DSD), accounting for less than 5% of all DSD cases, defined by the simultaneous presence of testicular tissue and ovarian tissue in the same individual. In the reported case, the patient presented two genetic mutations involved in the pathogenic pathway of the DSD condition associated with the clinical features of Kallmann syndrome (KS), a developmental disease that associates hypogonadotropic hypogonadism (HH), due to gonadotropin-releasing hormone deficiency, and anosmia, related to the absence or hypoplasia of the olfactory bulbs. Given the variable degree of hyposmia in KS, the distinction between KS and normosmic idiopathic HH is currently unclear, especially as HH patients do not always undergo detailed olfactory testing. This syndrome is very rare, with an estimated prevalence of 1:80,000 in males and 1:40,000 in females.

Keywords

- ovotestis
- ► Kallmann syndrome
- disorder of sex
 development
 affected by HH and digenic inheritance of Kallmann syndrome.

Introduction

True hermaphroditism is a disorder of sex development (DSD), accounting for less than 5% of all DSD cases, defined by the simultaneous presence of testicular tissue and ovarian tissue in the same individual. Kallmann syndrome (KS) is characterized by a developmental disease that associates hypogonadotropic hypogonadism (HH), with gonadotropin-releasing hormone (GnRH) deficiency, and anosmia,

relative to the absence or hypoplasia of the olfactory bulbs. This syndrome has an estimated prevalence of 1:80,000 in males and 1:40,000 in females.¹

Address for correspondence Francesca Peranzoni, MD, Rue du

bugnon 46, 1005 Lausanne, VD, Switzerland

(e-mail: francesca.peranzoni@chuv.ch).

Here we report the case of a patient with a 46 XX karyotype, gonadotropin-releasing hormone receptor (GnRHR), and PROKR2 mutation, diagnosed with true hermaphroditism who underwent genital reconstructive surgery during a humanitarian mission in Vietnam in autumn 2023.

DOI https://doi.org/ 10.1055/s-0044-1788060. ISSN 2699-9404. © 2024. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution License, permitting unrestricted use, distribution, and reproduction so long as the original work is properly cited. (https://creativecommons.org/licenses/by/4.0/). Georg Thieme Verlag KG, Rüdigerstraße 14, 70469 Stuttgart, Germany

Case Report

The patient, the first child of a nonconsanguineous couple, was referred to the Vietnam National Hospital of Pediatric 2 in Ho Chi Minh City immediately after birth due to ambiguous external genitalia. Family history and pregnancy anamnesis showed no particularities.

At the objective examination of the external genitalia, an ovalar structure was palpable in the left labioscrotal area. The collateral side was empty. Only one orifice, compatible with a proximal hypospadias or a urogenital sinus, was present in the perineal area. Anus was in the correct position as well as the anal sphincter (**-Fig. 1**).

An ultrasound was realized and confirmed the presence of an ovalar, vascularized structure in the left labioscrotal region, an abdominal ultrasound excluded the presence of intraabdominal Mullerian structure as well as the normality of other abdominal organs. A blood test sample showed normal electrolytes Na (138 mmol/L), K 4.75 mmol/L, Ca 2.49 mmol/L, Cl 107.5 mmol/L, and morning cortisol level between normal margins 30.7 pg/mL. The aldosterone was normal at 20.66 ng/dL, active renin was slightly higher at 43.12 ng/L, and beta-human choriogonadotropin (hCG) was normal at <1.20 mUL/mL. Testosterone was 47.26 ng/dL, which is a normal level for a female but a lower level for a male. Dehydroepiandrosterone (DHEASO4) was also normal.

A karyotype examination was realized and an endocrinological investigation was done. The child has confirmed 46 XX through a genetic karyotype analysis with the fluorescence in situ hybridization technique using the kit Cytocell SRY/DYZ1/DXZ1 to identify the SRY segment. The chromosome analysis was done through a Kit NEBNext (United States) and then the new generation sequencing was realized with the system NextSeq, Illumina (United States), the control reference DNA (hg19) was split in the region of 100 kb to detect the copy variations numbers defined according to the following database: http://decipher.sanger.ac.uk/, http://dgv.tcag.ca./, http://www.internationalgenome.org/, http://omim.org/. In conclusion, no CVN was detected. As a routine examination of the sequencing of the congenital adrenal hyperplasia, the sequencing of the CYP21A2 was realized and was normal.

Exploration laparoscopy was performed when the child was 2 years old; meanwhile, two biopsies of the ovalar structure were performed and the axis of the structure in the left labioscrotal region was realized. The absence of intraabdominal Mullerian remnant was confirmed. A few months later endoscopic examination of the urogenital sinus was realized and showed one vagina and one bladder with a common channel 3 cm long.

Biopsies, fixed in hematoxylin–eosin, revealed the presence of ovarian tissue. A new karyotype was realized. The result still confirmed a 46 XX. Genetic studies further confirmed the absence of the *SRY* gene, and the absence of mutation of the CYP21A2 gene and identified the mutation of two genes: GnRHR, autosomal recessive, heterozygous mutation missense, encoding for hypogonadism hypogonadal 7 without anosmia, and PROKR2 autosomal dominant missense mutation responsible for hypogonadism hypogonadotropic 3 with or without anosmia.

According to these considerations and the female actinglike attitude, the patient underwent a female genital reconstruction with reduction clitoridoplasty and labioplasty at 6 years of age.

The previous fixed structure in the left labia was reached through an inguinal approach, at that moment a structure compatible with ovotestis was found (**Fig. 2**). The structure was further split (**Fig. 3**) and the testis was removed and sent to anatomy pathology for histological examination. The remnant ovary was pulled into the abdominal cavity through



Fig. 1 Phenotypical appearance.



Fig. 2 Ovotestis.



Fig. 3 Splitted ovotestis, on the right the testis and the ovary in the left.

the inguinal ring. The specimen, fixed with hematoxylin and eosin, showed a testis structure with spermatic ducts and vas deferens melted with fibrotic tissue.

The patient was discharged 2 weeks after the surgery.

At 6-month follow-up, the patient had no complaints, and the neovagina presented a good healing process.

Discussion

In human fetuses during the ambisexual stage, primordial germ cells migrate from their site of origin in the allantois into the genital ridges around week 5. Sex cords are formed within the gonads in the sixth embryonic week, and the urogenital folds and labioscrotal folds begin to form in the perineum. Neurons that produce GnRH develop from the epithelium of the medial olfactory pit and move to the fetal hypothalamus by migrating along nerve fibers.² This process occurs at approximately 40 days of gestation.³ The pituitary gland develops and begins synthesizing both luteinizing hormone (LH) and follicle-stimulating hormone (FSH) at 9 weeks of gestation.⁴

The Mullerian and Wolffian ducts form around the 7th embryonic week and gonadal differentiation occurs from the 10th through 12th embryonic weeks.

The Sex-determining Region of the Y chromosome (*SRY*) gene is required to initiate signaling for male gonadal differentiation.⁵ *SRY* gene expression results in the differentiation of Sertoli cells, responsible for Mullerian-inhibiting substance production. The fetal testicles also express FSH receptors that probably control Sertoli cell proliferation, although only a few studies exist regarding the effects of FSH.^{6,7} SRY also causes the differentiation of Leydig cells.

During the first trimester of gestation, placental hCG induces the differentiation of Leydig cells and stimulates

testosterone (T) production through the activation of the LH/CG receptors expressed on their surfaces. After this period T secretion comes under the control of pituitary LH.⁸ T permits the persistence of the Wolffian duct and its differentiation into the epididymis, vas deferens, and seminal vesicles.

 5α -reductase is an enzyme that converts testosterone to dihydrotestosterone. This androgen drives the growth of the external genitalia and prostate.⁹

Additional genes involved in the gonadal genesis and normal SRY levels of expression are GATA4, FOG2,¹⁰ and SRY box-related gene 9.¹¹

The absence of the *SRY* gene results in feminine differentiation. In females, the Mullerian ducts persist and develop into the Fallopian tubes, uterus, cervix, and the cranial portion of the vagina, and the Wolffian ducts regress. The gonadotropin levels peak at midgestation and decrease toward birth and are suppressed at term.¹²

The development of the primordial follicle in the fetal ovaries begins before the 13th week of gestation. The pool of primordial follicles is approximately 680,000 at the 34th week of gestation. Subsequently, the pool remains stable.¹³ This pool relies on estrogen levels in the fetal circulation, which are mainly produced by the placenta. The roles of FSH and LH in ovarian development during pregnancy are not completely understood. It seems that hypothalamic stimulation is necessary to ensure physiological ovarian development after the 7th month of gestation.¹⁴

Additionally, female fetuses produce higher LH and FSH levels than male fetuses.^{15,16} Another marked difference between the sexes is that the LH levels overcome the FSH levels in male fetuses,¹⁷ whereas the opposite situation occurs in females.

At delivery, in healthy infants, the LH and FSH levels are low in the cord blood in both sexes.¹⁸

This complex genetic signaling pathway is not the unique determining factor of the development of sexual identity. Environmental factors also play an essential role in gonadal differentiation, and many studies have shown a link between genital malformation in humans and exposure to endocrine-disrupting chemical compounds,¹⁹ generally related to parental occupation during pregnancy.²⁰ Five-year-old Vietnamese children demonstrated disruptions in androgen, including decreases in T and DHEA levels and an increase in A-dione levels, which are associated with highly chlorinated dioxin congeners. These disorders are likely due to stimulation or inhibition of the activity of steroidogenic enzymes such as CYP17 lyase, 3β-HSD, and 17β-HSD by dioxin exposure.²¹

In the reported case, the patient presented two genetic mutations involved in the pathogenic pathway of the DSD condition associated with the clinical features of KS, a developmental disease that associates with HH, due to GnRH) deficiency, and anosmia, related to the absence or hypoplasia of the olfactory bulbs. Given the variable degree of hyposmia in KS, the distinction between KS and normosmic idiopathic HH is currently unclear, especially as HH patients do not always undergo detailed olfactory testing.²²

This syndrome is very rare, with an estimated prevalence of 1:80,000 in males and 1:40,000 in females.²³

The GNRHR, autosomal recessive heterozygous missense mutation, concerns the GNRHR gene that encodes the receptor for type 1 GnRH. This receptor is expressed on the surface of pituitary gonadotrope cells as well as lymphocytes, breast, ovary, and prostate. Following the binding of GnRHR, the receptor associates with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Activation of the receptor ultimately causes the release of gonadotropic LH and FSH. Defects in this gene are a cause of HH.^{23,24}

The patient presented also an autosomal dominant heterozygous missense mutation of the *PROKR2* gene, the protein encoded by this gene is an integral membrane protein and G protein-coupled receptor for prokinetics. Prokineticins are secreted cysteine-rich proteins that possess diverse biological activities including effects on neuronal survival, gastrointestinal smooth muscle contraction,²⁵ and circadian locomotor rhythm.²⁶ Prokineticins can bind to two different G protein-coupled receptors, prokineticin receptor-1 and -2 (PROKR1 and PROKR2), which have about 85% sequence identity. PROKR2 shows relatively localized distribution in the central nervous system.²⁵

Moreover, the mRNA expression of PROKR2 has been detected in the subventricular zone and the olfactory bulbs.

It was found that Prokr2 -/- knockout mice exhibit early hypoplasia of the olfactory bulbs and severe atrophy of the reproductive organs in both sexes, a phenotype reminiscent of the KS features. In addition, immunohistochemical analysis of these mice revealed that the neuroendocrine GnRH cells were absent from the hypothalamus.²⁷

To date, digenic inheritance of KS has been shown in few patients who had monoallelic missense mutations both in *PROKR2* or *PROK2* and in other KS genes (KAL1, FGFR1) or genes underlying normosmic congenital HH (GNRHR, KISS1R).^{28,29}

A subgroup of patients with KS can present other clinical manifestations, such as bimanual synkinesis, abnormal ocular movements, congenital ptosis, coloboma of the iris, nasal cartilage agenesis, malformation of the external ear, cleft lip or palate, unilateral or bilateral renal agenesis, agenesis of the corpus callosum, hypodontia, skeletal anomalies of the feet and hands, and obesity.^{22,30,31} Additional anomalies have so far not been reported in KS patients carrying mutations in *PROKR2* or *PROK2*, with the notable exception of a severe sleep disorder and marked obesity in one patient,³⁰ which could be related to the known function of prokineticin-2 signaling in behavioral circadian rhythms, including sleep–wake and ingestive behavior.³² In the reported case, none of these associated signs and symptoms were present.

This is the only case report concerning a patient with 46 XX true hermaphroditism affected by HH and digenic inheritance of KS.

In the diagnosis process of a patient with ambiguous genital malformation, primary importance is given to an accurate anamnesis, looking for inherited forms of genital anomalies and maternal exposure to chemical compounds.

Ultrasound as well as other complementary imaging such as magnetic resonance imaging (MRI) and computed tomog-

raphy as well as exploration laparoscopy helps to rule out associated anomalies and to better understand the internal genitalia development. Nevertheless, the usefulness of forebrain MRI in diagnosing the disease in children too young to undergo meaningful testing of olfaction or the hypothalamic-pituitary–gonadal axis should be emphasized,³³ even though normal olfactory bulb images have been reported in a few KS patients.³⁴

Endocrinology assessment is mandatory to rule out any deficiency in hormonal pathways.

Biopsies and histological studies of the surgical samples can help determine the sex of the patient; in this case, two biopsies of the left inguinoscrotal mass were carried out preoperatively, showing ovarian tissue, and the surgical specimen showed well-differentiated testis.

Moreover, genetic study and karyotype of patients with genital ambiguity are mandatory to formulate the correct diagnosis and offer appropriate therapy.

Conclusion

In the case of sexual differentiation disorders, definite gender assignment can be challenging, and it is based on phenotype, genotype, gonadal histology, rearing gender, and social factors.

Genotypes play an important role in the decision-making process and help in understanding the individual basis of the pathogenetic origin of the condition. Nevertheless, some pathogenic pathways are not yet completely understood, as the role of gonadotropin during the embryologic development of the genital tract, and further research in families carrying these mutations is necessary to allow a better understanding of the development of sexual characters.

A multidisciplinary discussion between endocrinology, pediatrician, geneticist pediatric surgeon, and urologist is necessary to assure the patient of the best possible treatment option and gender assignment.

Patient Consent

The family of the patient gave written informed consent to publish this article.

Conflict of Interest None declared.

References

- 1 The genetic aspects of primary eunuchoidism. Semantic Scholar. Accessed June 20, 2024 at: https://www.semanticscholar.org/ paper/The-genetic-aspects-of-primary-eunuchoidism-Kallmann/9f415d8d9e1a4aab4faff38d21bfa067ee4401ca
- 2 Schwanzel-Fukuda M, Pfaff DW. Origin of luteinizing hormonereleasing hormone neurons. Nature 1989;338(6211):161–164
- 3 Schwanzel-Fukuda M, Crossin KL, Pfaff DW, Bouloux PM, Hardelin JP, Petit C. Migration of luteinizing hormone-releasing hormone (LHRH) neurons in early human embryos. J Comp Neurol 1996; 366(03):547–557
- 4 Hagen C, McNeilly AS. The gonadotrophins and their subunits in foetal pituitary glands and circulation. J Steroid Biochem 1977;8 (05):537–544

- 5 Sinclair AH, Berta P, Palmer MS, et al. A gene from the human sexdetermining region encodes a protein with homology to a conserved DNA-binding motif. Nature 1990;346(6281):240–244
- 6 Huhtaniemi IT, Yamamoto M, Ranta T, Jalkanen J, Jaffe RB. Folliclestimulating hormone receptors appear earlier in the primate fetal testis than in the ovary. J Clin Endocrinol Metab 1987;65(06): 1210–1214
- 7 O'Shaughnessy PJ, Baker PJ, Monteiro A, Cassie S, Bhattacharya S, Fowler PA. Developmental changes in human fetal testicular cell numbers and messenger ribonucleic acid levels during the second trimester. J Clin Endocrinol Metab 2007;92(12):4792–4801
- 8 Tapanainen J, Kellokumpu-Lehtinen P, Pelliniemi L, Huhtaniemi I. Age-related changes in endogenous steroids of human fetal testis during early and midpregnancy. J Clin Endocrinol Metab 1981;52 (01):98–102
- 9 Wilson JD, Griffin JE, Leshin M, George FW. Role of gonadal hormones in development of the sexual phenotypes. Hum Genet 1981;58(01):78–84
- 10 Tevosian SG, Albrecht KH, Crispino JD, Fujiwara Y, Eicher EM, Orkin SH. Gonadal differentiation, sex determination and normal Sry expression in mice require direct interaction between transcription partners GATA4 and FOG2. Development 2002;129(19):4627–4634
- 11 Agrawal R, Wessely O, Anand A, Singh L, Aggarwal RK. Malespecific expression of Sox9 during gonad development of crocodile and mouse is mediated by alternative splicing of its prolineglutamine-alanine rich domain. FEBS J 2009;276(15):4184–4196
- 12 Massa G, de Zegher F, Vanderschueren-Lodeweyckx M. Serum levels of immunoreactive inhibin, FSH, and LH in human infants at preterm and term birth. Biol Neonate 1992;61(03):150–155
- 13 Forabosco A, Sforza C. Establishment of ovarian reserve: a quantitative morphometric study of the developing human ovary. Reprod Endocrinol 2007;88(03):P675–P683
- 14 Baker TG, Scrimgeour JB. Development of the gonad in normal and anencephalic human fetuses. Reproduction 1980;60(01):193–199
- 15 Clements JA, Reyes FI, Winter JS, Faiman C. Studies on human sexual development. III. Fetal pituitary and serum, and amniotic fluid concentrations of LH, CG, and FSH. J Clin Endocrinol Metab 1976;42(01):9–19
- 16 Kaplan SL, Grumbach MM. The ontogenesis of human foetal hormones. II. Luteinizing hormone (LH) and follicle stimulating hormone (FSH). Acta Endocrinol (Copenh) 1976;81(04):808–829
- 17 Beck-Peccoz P, Padmanabhan V, Baggiani AM, et al. Maturation of hypothalamic-pituitary-gonadal function in normal human fetuses: circulating levels of gonadotropins, their common alphasubunit and free testosterone, and discrepancy between immunological and biological activities of circulating follicle-stimulating hormone. J Clin Endocrinol Metab 1991;73(03):525–532
- 18 Lanciotti L, Cofini M, Leonardi A, Penta L, Esposito S. Up-to-date review about minipuberty and overview on hypothalamic-pituitary-gonadal axis activation in fetal and neonatal life. Front Endocrinol (Lausanne) 2018;9:410

- 19 Wang M-H, Baskin LS. Endocrine disruptors, genital development, and hypospadias. J Androl 2008;29(05):499–505
- 20 Nassar N, Abeywardana P, Barker A, Bower C. Parental occupational exposure to potential endocrine disrupting chemicals and risk of hypospadias in infants. Occup Environ Med 2010;67(09): 585–589
- 21 Oanh NTP, Kido T, Honma S, et al. Androgen disruption by dioxin exposure in 5-year-old Vietnamese children: decrease in serum testosterone level. Sci Total Environ 2018;640-641:466–474
- 22 Dodé C, Hardelin J-P. Kallmann syndrome. Eur J Hum Genet 2009; 17(02):139–146
- 23 McArdle C. Gonadotropin releasing hormone (GnRH) receptor. In: Enna SJ, Bylund DB, eds. xPharm: The Comprehensive Pharmacology Reference. New York: Elsevier; 2007:1–12
- 24 GNRHR gonadotropin releasing hormone receptor [*Homo sapiens* (human)]. Accessed June 20, 2024 at: https://www.ncbi.nlm. nih.gov/gene/2798
- 25 Li M, Bullock CM, Knauer DJ, Ehlert FJ, Zhou QY. Identification of two prokineticin cDNAs: recombinant proteins potently contract gastrointestinal smooth muscle. Mol Pharmacol 2001;59(04): 692–698
- 26 Cheng MY, Bullock CM, Li C, et al. Prokineticin 2 transmits the behavioural circadian rhythm of the suprachiasmatic nucleus. Nature 2002;417(6887):405–410
- 27 Matsumoto S-I, Yamazaki C, Masumoto K-H, et al. Abnormal development of the olfactory bulb and reproductive system in mice lacking prokineticin receptor PKR2. Proc Natl Acad Sci U S A 2006;103(11):4140–4145
- 28 Dodé C, Teixeira L, Levilliers J, et al. Kallmann syndrome: mutations in the genes encoding prokineticin-2 and prokineticin receptor-2. PLoS Genet 2006;2(10):e17–5
- 29 Louden ED, Poch A, Kim HG, Ben-Mahmoud A, Kim SH, Layman LC. Genetics of hypogonadotropic hypogonadism-human and mouse genes, inheritance, oligogenicity, and genetic counseling. Mol Cell Endocrinol 2021;534:111334
- 30 Hardelin J-P, Dodé C The complex genetics of Kallmann syndrome: KAL1, FGFR1, FGF8, PROKR2, PROK2, et al. Sex Dev 2008;2(4-5): 181–193
- 31 de Roux N. Isolated gonadotropic deficiency with and without anosmia: a developmental defect or a neuroendocrine regulation abnormality of the gonadotropic axis. Horm Res 2005;64 (Suppl 2):48–55
- 32 Zhou Q-Y, Cheng MY. Prokineticin 2 and circadian clock output. FEBS J 2005;272(22):5703–5709
- 33 Truwit CL, Barkovich AJ, Grumbach MM, Martini JJ. MR imaging of Kallmann syndrome, a genetic disorder of neuronal migration affecting the olfactory and genital systems. AJNR Am J Neuroradiol 1993;14(04):827–838
- 34 Quinton R, Duke VM, de Zoysa PA, et al. The neuroradiology of Kallmann's syndrome: a genotypic and phenotypic analysis. J Clin Endocrinol Metab 1996;81(08):3010–3017