



Neurophysiology combined with sonography as a useful tool in median nerve leprosy diagnosis

Neurofisiologia combinada com ultrassonografia como ferramenta útil no diagnóstico da hanseníase do nervo mediano

Oswaldo J. M. Nascimento^{1,2}

¹ Universidade Federal Fluminense, Hospital Universitário Antonio Pedro, Departamento de Neurologia, Niterói RJ, Brazil.

² Academia Nacional de Medicina, Rio de Janeiro RJ, Brazil.

Arq. Neuro-Psiquiatr. 2024;82(12):s00441801743.

Carpal tunnel syndrome (CTS) is the most common compressive peripheral neuropathy worldwide, affecting individual quality of life and work-related activities, mainly in those living in low or middle-income countries. The incidence and prevalence of CTS vary in different target populations, with a higher rate in middle-aged individuals (50-59 years) and women, as recently corroborated in a study conducted in Taiwan with an approximate annual incidence of 0.4%.¹

In California workers, a total of 139,336 CTS cases were referred (incidence of 6.3 cases per 10,000 full-time workers), being in women (8.2) 3.3 times higher than that seen among men (2.5).² Specific data from Brazil is limited but reflects the role of occupational biomechanical overload in the development of CTS.³ In Rio Grande do Sul State was observed a female-to-male ratio of 5.6:1 with a mean age of ~48.3 years. Countryside population seems to be more susceptible due to agriculture and farming-intensive jobs.⁴ Kouyoumdjian and Araujo (2006) evaluated CTS electrodiagnosis in 3.125 consecutive patients.⁵ They found 43 cases (1.38%) associated with manual milking (mean age was 44.9 years). The most interesting finding was the male prevalence (88.4%); manual milking work is practically a male job; it reflects the role of occupational biomechanical overload in CTS.⁵ Even though CTS pathophysiological mechanisms are not fully understood, the existence of an underlying subclinical neuropathy is a factor that may increase the susceptibility for the median nerve entrapment. Metabolic, hormonal disturbances, obesity, pregnancy, hereditary diseases, inflammation, and other associated conditions can be implicated.

Leprosy, a chronic infectious disease caused by *Mycobacterium leprae* (*M leprae*), is one of the most common treatable peripheral neuropathies in the world, being India and Brazil the most endemic countries. Leprosy is a neuritis of immuno-

logical origin that results in autonomic, sensory, and motor neuropathy.⁶ It can be effectively treated, including multidrug therapy, before disability appears. The risk of irreparable deficiencies is associated with late diagnosis. *M. leprae* has a strong tropism for Schwann cells, consequently resulting in progressive axonal damage.⁶ The neurological clinical presentations of leprosy have been related in 2013,⁶ including six cases of the rare CTS presentation.^{6,7} These cases of isolated median neuropathy were the first sign of leprosy, selected from patients with an exclusive median sensory complaint. Three of the patients had skin lesions, and three had pure neuritic leprosy (PNL).⁸ Clinical median nerve function impairment was confirmed by neurophysiological testing and histopathology. We highlighted the importance of early diagnosis so that the patient can receive adequate clinical treatment that consequently can avoid permanent disability.⁷

Recently, Naggapa et al. (2021) in a study with high-resolution sonography technique called attention to median nerve enlargement 2-cm proximal to the distal wrist crease in 26 patients, distinguishing leprosy from CTS.⁹ They concluded that this important discriminating sign can be used at point-of-care to identify patients with leprosy.

In Brazil, leprosy cases are also more prevalent in low-income areas, mainly in the countryside and in the Amazon region, where CTS are frequent. It should be noted that the diagnostic instruments used, the electromyography and high-resolution ultrasound devices, are expensive and require highly specialized and trained professionals, who are difficult to find in the regions with the highest prevalence of these two conditions in developing countries.

In this issue of *Arquivos de Neuro-Psiquiatria*, Alves and colleagues offer a very important and practical contribution revisiting the median nerve electrodiagnostic and high-

Address for correspondence
Oswaldo J. M. Nascimento
(email: osvaldo_nascimento@hotmail.com)

DOI <https://doi.org/10.1055/s-0044-1801743>.
ISSN 0004-282X.

received

December 12, 2024

accepted

December 15, 2024

© 2025. The Author(s).

This is an open access article published by Thieme under the terms of the Creative Commons Attribution 4.0 International License, permitting copying and reproduction so long as the original work is given appropriate credit (<https://creativecommons.org/licenses/by/4.0/>).

Thieme Revinter Publicações Ltda., Rua do Matoso 170, Rio de Janeiro, RJ, CEP 20270-135, Brazil

resolution sonography findings in patients with CTS or leprosy in the article entitled “Median nerve impairment in leprosy: how does it differ from the classic carpal tunnel syndrome?”¹⁰. The Brazilian contribution to leprosy knowledge has worldwide recognition. The importance of electrodiagnostic studies, recently associated with nerve imaging with high resolution sonography, brings a better understanding of the involvement of different nerve trunks that facilitates a more precise differential diagnosis and the best treatment window for leprosy. This is the focus of Alves and colleagues, considering the PNL modality.¹⁰ There were 29 patients included (CTS: 14; PNL: 15), with a female predominance (55.2%; 16/29). 92.8% of patients with CTS and 80.0% with PNL had bilateral impairment of the median nerve, with a total of 27 nerves affected in each group. This study is the first to evaluate the morphological and functional impairment of the median nerve in a combined and systematic way, favoring a better etiological diagnosis of PNL.¹⁰ They conclude that the presence of neural thickening and demyelinating involvement in the segments proximal to the carpal tunnel favors the diagnosis of leprosy. This methodology could be applied to other nerve trunks favoring the differential diagnosis of leprosy, including its challenging PNL form of presentation. Inflammatory chronic autoimmune neuropathies, other infectious neuropathies, inherited neuropathies, and others can be benefited with this dual methodology facing a clinical CTS case in endemic areas, as elegantly shown by Alves et al.¹⁰

Conflict of Interest

There is no conflict of interest to declare.

References

- 1 Yang C, Chen HH, Lee MC, et al. Risk Factors of Carpal Tunnel Syndrome in Taiwan: A Population-Based Cohort Study. *Ann Plast Surg* 2022;88(01):74–78
- 2 Jackson R, Beckman J, Frederick M, Musolin K, Harrison R. Rates of Carpal Tunnel Syndrome in a State Workers' Compensation Information System, by Industry and Occupation - California, 2007-2014. *MMWR Morb Mortal Wkly Rep* 2018;67(39):1094–1097
- 3 da Silva Pontes N, Costa de Assis SJ, de Oliveira GS, et al. Social determinants and work-related musculoskeletal disorders in Brazil. *PLoS One* 2024;19(07):e0306840. Doi: 10.1371/journal.pone.0306840
- 4 Nora DB, Becker J, Ehlers JA, Gomes I. Clinical features of 1039 patients with neurophysiological diagnosis of carpal tunnel syndrome. *Clin Neurol Neurosurg* 2004;107(01):64–69. Doi: 10.1016/j.clineuro.2004.08.003
- 5 Kouyoumdjian JA. [Carpal tunnel syndrome: clinical and epidemiological study in 668 cases]. *Arq Neuropsiquiatr* 1999;57(2A):202–207
- 6 Nascimento OJ. Leprosy neuropathy: clinical presentations. *Arq Neuropsiquiatr* 2013;71(9B):661–666
- 7 Vital RT, Illarramendi X, Antunes SL, et al. Isolated median neuropathy as the first symptom of leprosy. *Muscle Nerve* 2013;48(02):179–184
- 8 Tomaselli PJ, Dos Santos DF, Dos Santos ACJ, et al. Primary neural leprosy: clinical, neurophysiological and pathological presentation and progression. *Brain* 2022;145(04):1499–1506
- 9 Nagappa M, Pujar GS, Keshavan AH, et al. Sonographic pattern of median nerve enlargement in Hansen's neuropathy. *Acta Neurol Scand* 2021;144(02):155–160
- 10 Alves PHSC, Cirino FO, Garcia LP, et al. Median nerve impairment in leprosy: how does it differ from the classic carpal tunnel syndrome? *Arq Neuropsiquiatr* 2024;82(12):1–6. Doi: 10.1055/s-0044-1792092