Images in Focus

A Signet-Ring Cell Carcinoma of the Stomach Endoscopically Recognized as a Tiny Discolored Spot

M. Sanaka ¹, Y. Kuyama ¹, M. Yamanaka ¹, Y. Tamura ²


- ¹ Dept. of Internal Medicine, Teikyo University, School of Medicine, Tokyo, Japan
- ² First Dept. of Internal Medicine, Tokyo Medical and Dental University, School of Medicine, Tokyo, Japan


References

- Oohara T, Aono G, Ukawa S, et al. Clinical diagnosis of minute gastric cancer less than 5 mm in diameter. Cancer 1984; 53: 162-5.
- Hioki K, Nakane Y, Yamamoto M. Surgical strategy for early gastric cancer. Br J Surg 1990; 77: 1330 – 4.

Corresponding Author

M. Sanaka, M.D.
Dept. of Internal Medicine
Teikyo University School of Medicine
Kaga 2-11-1
Itabashi-ku
Tokyo 173
Japan
Fax: +81-3-3964-7094





A 4-mm diameter well demarcated slightly depressed discolored area (arrow) was seen in the anterior wall of the antrum in a 29-year-old woman (Figure 1) (1). It was biopsied twice. Since signet-ring cells were histologically identified (Figure 2, arrow), gastrectomy was carried out (2). Pathological examination showed that there was no residual cancer in the resected stomach. The signet-ring cell carcinoma was tiny, and had been completely removed by the two biopsies.