

SYNLETT Spotlight 232

(Dichloroiodo)benzene (PhICl₂)

Compiled by Zhou Wu



This feature focuses on a reagent chosen by a postgraduate, highlighting the uses and preparation of the reagent in current research

Zhou Wu was born in Hebei Province, P. R. of China. She attended Hebei University of Science and Technology for her B.Sc. (2001) and Hebei University of Technology for her M.Sc. (2005) in chemical engineering. She is currently pursuing her Ph.D. under the supervision of Prof. Shende Jiang at Tianjin University. Her research is focused on the synthesis of fluorine-18 labelled radiopharmaceuticals for Positron Emission Tomography in molecular imaging and nuclear medicine.


School of Pharmaceutical Science and Technology, Tianjin University, Tianjin 300072, P. R. of China
E-mail: zwu@tju.edu.cn

Introduction

(Dichloroiodo)benzene (PhICl₂) was first prepared in 1886; its chemistry then became an intensely studied topic because of its wide application as an effective chlorinating or oxidizing agent in various organic reactions.¹ As a chlorinating agent, (dichloroiodo)benzene can be applied not only to the addition of chlorine to alkenes, but also to the chlorination with other substrates, such as alkanes, ethers, esters, thioethers, ketones, sulfoxides, steroids, etc., with moderate to high yields.^{1,2} (Dichloroiodo)benzene can oxidize aldoximes and ketoximes to nitrile oxides³ and ketones,⁴ respectively in good yields. In this spotlight, recent applications using PhICl₂ in organic syntheses are summarized.

Preparation

(Dichloroiodo)benzene was prepared by bubbling chlorine gas through a solution of iodobenzene in chlorinated solvents,⁵ which had been applied to an industrial-scale preparation (Scheme 1).⁶




Scheme 1

Properties

(Dichloroiodo)benzene exists as yellow crystalline solid with mp 110–112 °C. It is not stable on long standing, which means that it is normally prepared and used fresh for subsequent applications.


Abstracts

(A) Alkynes can be effectively thiocyanated by the combination of PhICl₂/Pb(SCN)₂ in dichloromethane to yield the corresponding alkynes with high *E/Z* ratio.⁷




R¹ = Ph, *p*-MeC₆H₄, *n*-Pr, Me(CH₂)_{*n*}, (*n* = 3, 5, 7)
HO(CH₂)₄, 1-cyclohexenyl; R² = H, Me, Ph, TMS

(B) A combination of PhICl₂/KSCN can thiocyanate β-dicarbonyl compounds, phenols as well as anilines to the corresponding thiocyanato derivatives in dichloromethane.⁸




R = Ph, *p*-ClC₆H₄, *p*-MeC₆H₄,
p-MeOC₆H₄, *p*-O₂NC₆H₄




R¹ = NH₂, R² = H, OH, COOH, R³ = H or
R¹ = OH, R² = H, OH, R³ = H


(C) PhICl_2 can be used for the chloromethoxylation, iodomethoxylation, and iodohydroxylation, and iodochlorination of alkenes under different conditions with good yields.^{9,10}




(D) Aliphatic primary and secondary alcohols can be oxidized to the corresponding aldehydes and ketones in good to excellent yields by the combination of PhICl_2 /2,2,6,6-tetramethylpiperidin-1-yloxy (TEMPO)/pyridine in CHCl_3 . This oxidizing system has preference to oxidize secondary alcohols over primary ones.¹¹



(E) Reactions of PhICl_2 with 4-hydroxypyridin-2-ones in aq Na_2CO_3 gave an aryliodonium ylide that rearranged to 3-iodo-4-phenoxy-pyridinones. This operation can install both the iodine and aryloxy groups to the pyridinone nucleus.¹²



(F) PhICl_2 is a good reagent for the oxidative chlorination of various metal complexes.¹³ For example, treatment of dinuclear platinum(II) complexes as a mixture of isomers (the half-lantern and lantern complexes) with PhICl_2 in dichloromethane gave the dichlorodiplatinum(III) lantern complex in excellent yield.¹⁴



References

- (1) (a) Stang, P. J.; Zhdankin, V. V. *Chem. Rev.* **1996**, *96*, 1123. (b) Zhdankin, V. V.; Stang, P. J. *Chem. Rev.* **2002**, *102*, 2523.
- (2) *Topics in Current Chemistry: Hypervalent Iodine Chemistry, Modern Developments in Organic Synthesis*; Wirth, T., Ed.; Springer: Berlin, **2002**.
- (3) Radhakrishna, A. S.; Sivaprakash, K.; Singh, B. B. *Synth. Commun.* **1991**, *21*, 1625.
- (4) Radhakrishna, A. S.; Augustine, B.; Sivaprakash, K.; Singh, B. B. *Synth. Commun.* **1991**, *21*, 1473.
- (5) Lucas, H. J.; Kennedy, E. R. *Org. Synth., Coll. Vol. III* **1955**, 482.
- (6) Zanka, A.; Takeuchi, H.; Kubota, A. *Org. Process Res. Dev.* **1998**, *2*, 270.
- (7) Prakash, O.; Sharma, V.; Batra, H.; Moriarty, R. M. *Tetrahedron Lett.* **2001**, *42*, 553.
- (8) Prakash, O.; Kaur, H.; Pundeer, R.; Dhillon, R. S.; Singh, S. P. *Synth. Commun.* **2003**, *33*, 4037.
- (9) Yusubov, M. S.; Yusubova, R. J.; Filimonov, V. D.; Chi, K. W. *Synth. Commun.* **2004**, *34*, 443.
- (10) Yusubov, M. S.; Drygunova, L. A.; Tkachev, A. V.; Zhdankin, V. V. *ARKIVOC* **2005**, (iv), 179.
- (11) Zhao, X. F.; Zhang, C. *Synthesis* **2007**, 551.
- (12) Benjahad, A.; Guillemont, J.; Andries, K.; Nguyen, C. H.; Grierson, D. S. *Bioorg. Med. Chem. Lett.* **2003**, *13*, 4309.
- (13) (a) Bachmann, J.; Hodgkiss, J. M.; Young, E. R.; Nocera, D. G. *Inorg. Chem.* **2007**, *46*, 607. (b) Cotton, F. A.; Koshevoy, I. O.; Lahuerta, P.; Murillo, C. A.; Sanaú, M.; Ubeda, M. A.; Zhao, Q. *J. Am. Chem. Soc.* **2006**, *128*, 13674. (c) Jin, L.; Yin, J.; Chen, L.; Guo, C.; Chen, Q. *Synlett* **2005**, 2893. (d) Martins, A. M.; Branquinho, R.; Cui, J.; Dias, A. R.; Duarte, M. T.; Fernandes, J.; Rodrigues, S. S. *J. Organomet. Chem.* **2004**, *689*, 2368.
- (14) Bennett, M. A.; Bhargava, S. K.; Bond, A. M.; Edwards, A. J.; Guo, S.; Privér, S. H.; Rae, D.; Willis, A. C. *Inorg. Chem.* **2004**, *43*, 7752.