


N. UMEBAYASHI, Y. HAMASHIMA, D. HASHIZUME, M. SODEOKA* (RIKEN, WAKO AND SAITAMA UNIVERSITY, JAPAN)


Catalytic Enantioselective Aldol-type Reaction of β -Ketoesters with Acetals

Angew. Chem. Int. Ed. **2008**, *47*, 4196-4199.

Pd-Catalyzed Aldol Reaction for Formation of Chiral Quaternary Centers



Selected examples:



Significance: The low nucleophilicity of 1,3-dicarbonyl metal enolates and the ready retro-aldol reaction have stalled the development of metal catalysts for asymmetric aldol reactions. The protocol uses overall neutral conditions where only the metal center acts as Lewis acid. Using BINAP as the chiral ligand, the reaction yields protected alcohols directly with excellent enantioselectivity.

Comment: Recognizing acetals as oxocarbenium ion precursors is the key for the success of the reaction. Competitive retro-aldol reactions are prevented. Furthermore, the strategy affords highly functionalized chiral products bearing chemodifferentiated synthetic handles (ketone, alkene, ester and protected alcohol). In some cases, a palladium catalyst was not optimal and necessitated the use of a more stable platinum catalyst (i.e. **3e**). Intriguingly, when an unsubstituted acrylate acetal was used, it reacted via a 1,4-addition (**6**).

SYNFACTS Contributors: Mark Lautens, Frédéric Ménard
Synfacts 2008, 8, 0853-0853 Published online: 23.07.2008
DOI: 10.1055/s-2008-1077917; Reg-No.: L08808SF

2008 © THIEME STUTTGART • NEW YORK

Category

Metal-Catalyzed
Asymmetric
Synthesis and
Stereoselective
Reactions

Key words

palladium

quaternary centers

catalytic aldol
reaction

SYNFACTS
of the month

This document was downloaded for personal use only. Unauthorized distribution is strictly prohibited.

853